
Marcella Mun-San Kwan

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/3984129/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Factors associated with appointment nonâ€attendance at a medical imaging department in regional Australia: a retrospective cohort analysis. Journal of Medical Radiation Sciences, 2018, 65, 192-199.	1.5	16
2	Effect of Exercise and Cognitive Training on Falls and Fall-Related Factors in Older Adults With Mild Cognitive Impairment: A Systematic Review. Archives of Physical Medicine and Rehabilitation, 2017, 98, 2079-2096.	0.9	61
3	The rural pipeline to longer-term rural practice: General practitioners and specialists. PLoS ONE, 2017, 12, e0180394.	2.5	55
4	Fall risk in <scp>C</scp> hinese communityâ€dwelling older adults: A physiological profile assessment study. Geriatrics and Gerontology International, 2016, 16, 259-265.	1.5	11
5	Comparison of faecal microbiota in Blastocystis-positive and Blastocystis-negative irritable bowel syndrome patients. Microbiome, 2016, 4, 47.	11.1	77
6	Blastocystis specific serum immunoglobulin in patients with irritable bowel syndrome (IBS) versus healthy controls. Parasites and Vectors, 2015, 8, 453.	2.5	15
7	Effectiveness of Exergaming Training in Reducing Risk and Incidence of Falls in Frail Older Adults With a History of Falls. Archives of Physical Medicine and Rehabilitation, 2015, 96, 2096-2102.	0.9	78
8	Development and validation of a Chinese version of the Falls Efficacy Scale International. Archives of Gerontology and Geriatrics, 2013, 56, 169-174.	3.0	55
9	Increased Concern Is Protective for Falls in Chinese Older People: The Chopstix Fall Risk Study. Journals of Gerontology - Series A Biological Sciences and Medical Sciences, 2013, 68, 946-953.	3.6	38
10	Depressive symptoms in addition to visual impairment, reduced strength and poor balance predict falls in older Taiwanese people. Age and Ageing, 2012, 41, 606-612.	1.6	31
11	Sensorimotor function, balance abilities and pain influence Timed Up and Go performance in older community-living people. Aging Clinical and Experimental Research, 2011, 23, 196-201.	2.9	59
12	Minimal Chair Height Standing Ability Is Independently Associated With Falls in Taiwanese Older People. Archives of Physical Medicine and Rehabilitation, 2011, 92, 1080-1085.	0.9	11
13	Falls Incidence, Risk Factors, and Consequences in Chinese Older People: A Systematic Review. Journal of the American Geriatrics Society, 2011, 59, 536-543.	2.6	196
14	Effect on falls of providing single lens distance vision glasses to multifocal glasses wearers: VISIBLE randomised controlled trial. BMJ: British Medical Journal, 2010, 340, c2265-c2265.	2.3	104
15	Preventing falls in older multifocal glasses wearers by providing single-lens distance glasses: the protocol for the VISIBLE randomised controlled trial. BMC Geriatrics, 2009, 9, 10.	2.7	18
16	Group exercise can improve participants' mobility in an outpatient rehabilitation setting: a randomized controlled trial. Clinical Rehabilitation, 2008, 22, 493-502.	2.2	39
17	A novel weight-bearing strengthening program during rehabilitation of older people is feasible and improves standing up more than a non-weight-bearing strengthening program: a randomised trial. Australian Journal of Physiotherapy, 2007, 53, 147-153.	0.9	37
18	Foot pain in community-dwelling older people: an evaluation of the Manchester Foot Pain and Disability Index. Rheumatology, 2006, 45, 863-867.	1.9	126

#	Article	IF	CITATIONS
19	Reliability of the GAITRite® walkway system for the quantification of temporo-spatial parameters of gait in young and older people. Gait and Posture, 2004, 20, 20-25.	1.4	577
20	Reliability of Clinical Tests of Foot and Ankle Characteristics in Older People. Journal of the American Podiatric Medical Association, 2003, 93, 380-387.	0.3	80