B Babbette Lamarca

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/3995261/publications.pdf Version: 2024-02-01

| | | 76196 | 110170 |
|----------|----------------|--------------|----------------|
| 123 | 4,445 | 40 | 64 |
| papers | citations | h-index | g-index |
| | | | |
| | | | |
| | | | |
| 123 | 123 | 123 | 3255 |
| all docs | docs citations | times ranked | citing authors |
| | | | |

R RABBETTE LAMADCA

| # | Article | IF | CITATIONS |
|----|---|-----|-----------|
| 1 | The role of inflammation in the pathology of preeclampsia. Clinical Science, 2016, 130, 409-419. | 1.8 | 379 |
| 2 | Hypertension in Response to Autoantibodies to the Angiotensin II Type I Receptor (AT1-AA) in Pregnant Rats. Hypertension, 2009, 54, 905-909. | 1.3 | 185 |
| 3 | Pathophysiology and Current Clinical Management of Preeclampsia. Current Hypertension Reports, 2017, 19, 61. | 1.5 | 175 |
| 4 | A model of preeclampsia in rats: the reduced uterine perfusion pressure (RUPP) model. American Journal of Physiology - Heart and Circulatory Physiology, 2012, 303, H1-H8. | 1.5 | 163 |
| 5 | Autoantibodies to the Angiotensin Type I Receptor in Response to Placental Ischemia and Tumor Necrosis Factor α in Pregnant Rats. Hypertension, 2008, 52, 1168-1172. | 1.3 | 153 |
| 6 | Hypertension in Response to Chronic Reductions in Uterine Perfusion in Pregnant Rats. Hypertension, 2008, 52, 1161-1167. | 1.3 | 150 |
| 7 | The Effect of Immune Factors, Tumor Necrosis Factor-Â, and Agonistic Autoantibodies to the Angiotensin II Type I Receptor on Soluble fms-Like Tyrosine-1 and Soluble Endoglin Production in Response to Hypertension During Pregnancy. American Journal of Hypertension, 2010, 23, 911-916. | 1.0 | 129 |
| 8 | CD4 ⁺ T-Helper Cells Stimulated in Response to Placental Ischemia Mediate Hypertension During Pregnancy. Hypertension, 2011, 57, 949-955. | 1.3 | 118 |
| 9 | Preeclampsia: long-term consequences for vascular health. Vascular Health and Risk Management, 2015, 11, 403. | 1.0 | 116 |
| 10 | IL-17-mediated oxidative stress is an important stimulator of AT1-AA and hypertension during pregnancy. American Journal of Physiology - Regulatory Integrative and Comparative Physiology, 2012, 303, R353-R358. | 0.9 | 114 |
| 11 | Role of Mitochondrial Dysfunction and Reactive Oxygen Species in Mediating Hypertension in the Reduced Uterine Perfusion Pressure Rat Model of Preeclampsia. Hypertension, 2018, 72, 703-711. | 1.3 | 112 |
| 12 | Hypertension in Response to Placental Ischemia During Pregnancy. Hypertension, 2011, 57, 865-871. | 1.3 | 107 |
| 13 | Administration of Interleukin-17 Soluble Receptor C Suppresses T _H 17 Cells, Oxidative Stress, and Hypertension in Response to Placental Ischemia During Pregnancy. Hypertension, 2013, 62, 1068-1073. | 1.3 | 99 |
| 14 | Endothelin-1, Oxidative Stress, and Endogenous Angiotensin II. Hypertension, 2013, 62, 886-892. | 1.3 | 82 |
| 15 | Angiotensin <scp>II</scp> Type 1 Receptor Autoantibody (<scp>AT</scp> 1â€ <scp>AA</scp>)â€Mediated Pregnancy Hypertension. American Journal of Reproductive Immunology, 2013, 69, 413-418. | 1.2 | 81 |
| 16 | Magnesium Sulfate Treatment Reverses Seizure Susceptibility and Decreases Neuroinflammation in a Rat Model of Severe Preeclampsia. PLoS ONE, 2014, 9, e113670. | 1.1 | 81 |
| 17 | Elucidating Immune Mechanisms Causing Hypertension During Pregnancy. Physiology, 2013, 28, 225-233. | 1.6 | 78 |
| 18 | Preventing Autoimmunity Protects Against the Development of Hypertension and Renal Injury. Hypertension, 2014, 64, 792-800. | 1.3 | 75 |

| # | Article | IF | CITATIONS |
|----|---|-----|-----------|
| 19 | Identifying immune mechanisms mediating the hypertension during preeclampsia. American Journal of Physiology - Regulatory Integrative and Comparative Physiology, 2016, 311, R1-R9. | 0.9 | 74 |
| 20 | IL-10 supplementation increases Tregs and decreases hypertension in the RUPP rat model of preeclampsia. Hypertension in Pregnancy, 2015, 34, 291-306. | 0.5 | 68 |
| 21 | An increased population of regulatory T cells improves the pathophysiology of placental ischemia in a rat model of preeclampsia. American Journal of Physiology - Regulatory Integrative and Comparative Physiology, 2015, 309, R884-R891. | 0.9 | 68 |
| 22 | Hypertension in Response to AT1-AA: Role of Reactive Oxygen Species in Pregnancy-Induced Hypertension. American Journal of Hypertension, 2011, 24, 835-840. | 1.0 | 67 |
| 23 | Activating autoantibodies to the angiotensin II type I receptor play an important role in mediating hypertension in response to adoptive transfer of CD4 ⁺ T lymphocytes from placental ischemic rats. American Journal of Physiology - Regulatory Integrative and Comparative Physiology, 2012. 302. R1197-R1201. | 0.9 | 65 |
| 24 | Role of angiotensin II type I receptor agonistic autoantibodies (AT1-AA) in preeclampsia. Current Opinion in Pharmacology, 2011, 11, 175-179. | 1.7 | 64 |
| 25 | Reduced uterine perfusion pressure T-helper 17 cells cause pathophysiology associated with preeclampsia during pregnancy. American Journal of Physiology - Regulatory Integrative and Comparative Physiology, 2016, 311, R1192-R1199. | 0.9 | 61 |
| 26 | Hypertension in response to IL-6 during pregnancy: role of AT1-receptor activation. International Journal of Interferon, Cytokine and Mediator Research, 2011, 2011, 65. | 1.1 | 59 |
| 27 | IL-6-induced pathophysiology during pre-eclampsia: potential therapeutic role for magnesium sulfate?. International Journal of Interferon, Cytokine and Mediator Research, 2011, 2011, 59. | 1.1 | 59 |
| 28 | Inflammatory mediators: a causal link to hypertension during preeclampsia. British Journal of Pharmacology, 2019, 176, 1914-1921. | 2.7 | 59 |
| 29 | Cytochrome P450 Subfamily 2J Polypeptide 2 Expression and Circulating Epoxyeicosatrienoic Metabolites in Preeclampsia. Circulation, 2012, 126, 2990-2999. | 1.6 | 57 |
| 30 | AT1-AA (Angiotensin II Type 1 Receptor Agonistic Autoantibody) Blockade Prevents Preeclamptic Symptoms in Placental Ischemic Rats. Hypertension, 2018, 71, 886-893. | 1.3 | 56 |
| 31 | Endothelin type A receptor antagonist attenuates placental ischemia–induced hypertension and uterine vascular resistance. American Journal of Obstetrics and Gynecology, 2011, 204, 330.e1-330.e4. | 0.7 | 55 |
| 32 | Placental Ischemia and Resultant Phenotype in Animal Models of Preeclampsia. Current Hypertension Reports, 2016, 18, 38. | 1.5 | 52 |
| 33 | Risk of cardiovascular disease, end-stage renal disease, and stroke in postpartum women and their fetuses after a hypertensive pregnancy. American Journal of Physiology - Regulatory Integrative and Comparative Physiology, 2018, 315, R521-R528. | 0.9 | 52 |
| 34 | 17-Hydroxyprogesterone Caproate Significantly Improves Clinical Characteristics of Preeclampsia in the Reduced Uterine Perfusion Pressure Rat Model. Hypertension, 2015, 65, 225-231. | 1.3 | 51 |
| 35 | The Role of Agonistic Autoantibodies to the Angiotensin II Type 1 Receptor (AT1-AA) in Pathophysiology of Preeclampsia. Current Pharmaceutical Biotechnology, 2018, 19, 781-785. | 0.9 | 49 |
| 36 | Effects of Reduced Uterine Perfusion Pressure on Blood Pressure and Metabolic Factors in Pregnant Rats. American Journal of Hypertension, 2007, 20, 686-691. | 1.0 | 48 |

| # | Article | IF | CITATIONS |
|----|---|-----|-----------|
| 37 | Agonistic Autoantibodies to the Angiotensin II Type 1 Receptor Enhance Angiotensin II–Induced Renal Vascular Sensitivity and Reduce Renal Function During Pregnancy. Hypertension, 2016, 68, 1308-1313. | 1.3 | 44 |
| 38 | Natural killer cells mediate pathophysiology in response to reduced uterine perfusion pressure. Clinical Science, 2017, 131, 2753-2762. | 1.8 | 44 |
| 39 | Agonistic Autoantibodies to the Angiotensin II Type I Receptor Cause Pathophysiologic Characteristics of Preeclampsia. Gender Medicine, 2012, 9, 139-146. | 1.4 | 42 |
| 40 | Effects of 17-Hydroxyprogesterone on Tumor Necrosis Factor-Â-Induced Hypertension During Pregnancy. American Journal of Hypertension, 2009, 22, 1120-1125. | 1.0 | 41 |
| 41 | Hypertension in response to CD4+ T cells from reduced uterine perfusion pregnant rats is associated with activation of the endothelin-1 system. American Journal of Physiology - Regulatory Integrative and Comparative Physiology, 2012, 303, R144-R149. | 0.9 | 40 |
| 42 | Progesterone blunts vascular endothelial cell secretion ofÂendothelin-1 in response to placental ischemia. American Journal of Obstetrics and Gynecology, 2013, 209, 44.e1-44.e6. | 0.7 | 39 |
| 43 | Endothelin, the kidney, and hypertension. Current Hypertension Reports, 2006, 8, 298-303. | 1.5 | 38 |
| 44 | <scp>l</scp> -Arginine supplementation abolishes the blood pressure and endothelin response to chronic increases in plasma sFlt-1 in pregnant rats. American Journal of Physiology - Regulatory Integrative and Comparative Physiology, 2012, 302, R259-R263. | 0.9 | 38 |
| 45 | CD4 ⁺ T Cells Are Important Mediators of Oxidative Stress That Cause Hypertension in Response to Placental Ischemia. Hypertension, 2014, 64, 1151-1158. | 1.3 | 37 |
| 46 | Role of Reactive Oxygen Species During Hypertension in Response to Chronic Antiangiogenic Factor (sFlt-1) Excess in Pregnant Rats. American Journal of Hypertension, 2011, 24, 110-113. | 1.0 | 34 |
| 47 | Vitamin D supplementation improves pathophysiology in a rat model of preeclampsia. American Journal of Physiology - Regulatory Integrative and Comparative Physiology, 2016, 310, R346-R354. | 0.9 | 33 |
| 48 | Renal natural killer cell activation and mitochondrial oxidative stress; new mechanisms in AT1-AA mediated hypertensive pregnancy. Pregnancy Hypertension, 2019, 15, 72-77. | 0.6 | 32 |
| 49 | Placental ischemia-stimulated T-helper 17 cells induce preeclampsia-associated cytolytic natural killer cells during pregnancy. American Journal of Physiology - Regulatory Integrative and Comparative Physiology, 2018, 315, R336-R343. | 0.9 | 31 |
| 50 | Serelaxin improves the pathophysiology of placental ischemia in the reduced uterine perfusion pressure rat model of preeclampsia. American Journal of Physiology - Regulatory Integrative and Comparative Physiology, 2016, 311, R1158-R1163. | 0.9 | 30 |
| 51 | CD4+ T Cells Play a Critical Role in Mediating Hypertension in Response to Placental Ischemia. Journal of Hypertension: Open Access, 2013, 02, . | 0.2 | 28 |
| 52 | Progesterone supplementation attenuates hypertension andÂthe autoantibody to the angiotensin II type I receptor inÂresponse to elevated interleukin-6 during pregnancy. American Journal of Obstetrics and Gynecology, 2014, 211, 158.e1-158.e6. | 0.7 | 26 |
| 53 | Preeclampsia: Linking Placental Ischemia with Maternal Endothelial and Vascular Dysfunction. , 2020, 11, 1315-1349. | | 26 |
| 54 | RNA interference therapeutics targeting angiotensinogen ameliorate preeclamptic phenotype in rodent models. Journal of Clinical Investigation, 2020, 130, 2928-2942. | 3.9 | 25 |

| # | Article | IF | CITATIONS |
|----|--|-----|-----------|
| 55 | Interleukin-4 supplementation improves the pathophysiology of hypertension in response to placental ischemia in RUPP rats. American Journal of Physiology - Regulatory Integrative and Comparative Physiology, 2019, 316, R165-R171. | 0.9 | 24 |
| 56 | Enodthelin 1 Is Elevated in Plasma and Explants From Patients Having Uterine Leiomyomas. Reproductive Sciences, 2014, 21, 1196-1205. | 1.1 | 23 |
| 57 | Proliferation of endogenous regulatory T cells improve the pathophysiology associated with placental ischaemia of pregnancy. American Journal of Reproductive Immunology, 2017, 78, e12724. | 1.2 | 22 |
| 58 | Arachidonic acid metabolites of CYP4A and CYP4F are altered in women with preeclampsia. Prostaglandins and Other Lipid Mediators, 2018, 136, 15-22. | 1.0 | 22 |
| 59 | Continued Investigation Into 17-OHPC. Hypertension, 2017, 70, 1250-1255. | 1.3 | 20 |
| 60 | 17-hydroxyprogesterone blunts the hypertensive response associated with reductions in uterine perfusion pressure in pregnant rats. American Journal of Obstetrics and Gynecology, 2009, 201, 324.e1-324.e6. | 0.7 | 19 |
| 61 | Tumor necrosis factor alpha (TNF-α) blockade improves natural killer cell (NK) activation, hypertension, and mitochondrial oxidative stress in a preclinical rat model of preeclampsia. Hypertension in Pregnancy, 2020, 39, 399-404. | 0.5 | 19 |
| 62 | Blockade of CD40 ligand for intercellular communication reduces hypertension, placental oxidative stress, and AT ₁ -AA in response to adoptive transfer of CD4 ⁺ T lymphocytes from RUPP rats. American Journal of Physiology - Regulatory Integrative and Comparative Physiology, 2015, 309, R1243-R1250. | 0.9 | 17 |
| 63 | 17-Hydroxyprogesterone caproate improves T cells and NK cells in response to placental ischemia; new mechanisms of action for an old drug. Pregnancy Hypertension, 2020, 19, 226-232. | 0.6 | 16 |
| 64 | Vitamin D supplementation reduces some AT ₁ -AA-induced downstream targets implicated in preeclampsia including hypertension. American Journal of Physiology - Regulatory Integrative and Comparative Physiology, 2017, 312, R125-R131. | 0.9 | 15 |
| 65 | CD4+ T cells cause renal and placental mitochondrial oxidative stress as mechanisms of hypertension in response to placental ischemia. American Journal of Physiology - Renal Physiology, 2021, 320, F47-F54. | 1.3 | 15 |
| 66 | Natural killer cells contribute to mitochondrial dysfunction in response to placental ischemia in reduced uterine perfusion pressure rats. American Journal of Physiology - Regulatory Integrative and Comparative Physiology, 2019, 316, R441-R447. | 0.9 | 14 |
| 67 | Progesterone-induced blocking factor improves blood pressure, inflammation, and pup weight in response to reduced uterine perfusion pressure (RUPP). American Journal of Physiology - Regulatory Integrative and Comparative Physiology, 2021, 320, R719-R727. | 0.9 | 14 |
| 68 | Placental CD4+ T cells isolated from preeclamptic women cause preeclampsia-like symptoms in pregnant nude-athymic rats. Pregnancy Hypertension, 2019, 15, 7-11. | 0.6 | 13 |
| 69 | Characterization of Mitochondrial Bioenergetics in Preeclampsia. Journal of Clinical Medicine, 2021, 10, 5063. | 1.0 | 13 |
| 70 | Vascular endothelial mitochondrial oxidative stress in response to preeclampsia: a role for angiotension II type 1 autoantibodies. American Journal of Obstetrics & Gynecology MFM, 2021, 3, 100275. | 1.3 | 10 |
| 71 | Endothelin-1 is not a Mechanism of IL-17 Induced Hypertension during Pregnancy. Medical Journal of Obstetrics and Gynecology, 2013, 1, . | 0.2 | 10 |
| 72 | Vitamin D Supplementation Suppresses Hypoxia-Stimulated Placental Cytokine Secretion, Hypertension and CD4 T Cell Stimulation in Response to Placental Ischemia. Medical Journal of Obstetrics and Gynecology, 2013, 1, . | 0.2 | 8 |

| # | Article | IF | CITATIONS |
|----|--|-----|-----------|
| 73 | Angiotensin II type 1 receptor autoantibody blockade improves cerebral blood flow autoregulation and hypertension in a preclinical model of preeclampsia. Hypertension in Pregnancy, 2020, 39, 451-460. | 0.5 | 7 |
| 74 | Progesterone Induced Blocking Factor Reduces Hypertension and Placental Mitochondrial Dysfunction in Response to sFlt-1 during Pregnancy. Cells, 2021, 10, 2817. | 1.8 | 7 |
| 75 | 17-Hydroxyprogesterone caproate improves hypertension and renal endothelin-1 in response to sFlt-1 induced hypertension in pregnant rats. Pregnancy Hypertension, 2020, 22, 151-155. | 0.6 | 6 |
| 76 | Investigation of interleukin-2-mediated changes in blood pressure, fetal growth restriction, and innate immune activation in normal pregnant rats and in a preclinical rat model of preeclampsia. Biology of Sex Differences, 2021, 12, 4. | 1.8 | 6 |
| 77 | Placental CD4+ T cells from preeclamptic patients cause autoantibodies to the angiotensin II type I receptor and hypertension in a pregnant rat model of preeclampsia. Exploration of Medicine, 0, , 99-111. | 1.5 | 6 |
| 78 | Selective inhibition of 20-hydroxyeicosatetraenoic acid lowers blood pressure in a rat model of preeclampsia. Prostaglandins and Other Lipid Mediators, 2018, 134, 108-113. | 1.0 | 5 |
| 79 | Low Dose of IL-2 Normalizes Hypertension and Mitochondrial Function in the RUPP Rat Model of Placental Ischemia. Cells, 2021, 10, 2797. | 1.8 | 4 |
| 80 | Is Mitochondrial Oxidative Stress a Viable Therapeutic Target in Preeclampsia?. Antioxidants, 2022, 11, 210. | 2.2 | 4 |
| 81 | The Renin-Angiotensin System, its Autoantibodies, and Body Fluid Volume in Preeclampsia. , 2015, , 315-334. | | 3 |
| 82 | Effects of reduced uterine perfusion pressure on blood pressure and metabolic factors in the pregnant rat. FASEB Journal, 2007, 21, A894. | 0.2 | 3 |
| 83 | Novel treatment avenues for uterine leiomyoma: a new implication for endothelin?. Clinical Science, 2018, 132, 2261-2267. | 1.8 | 2 |
| 84 | Progesterone inhibits trophoblast TNF alpha production FASEB Journal, 2010, 24, 793.13. | 0.2 | 2 |
| 85 | 233: T Lymphocyte induced AT1-AAs cause hypertension in response to placental ischemia. American Journal of Obstetrics and Gynecology, 2011, 204, S100. | 0.7 | 1 |
| 86 | Letter to the Editor: Importance of B cells in response to placental ischemia. American Journal of Physiology - Heart and Circulatory Physiology, 2020, 318, H723-H725. | 1.5 | 1 |
| 87 | The role of angiotensin II type I receptor activation in mediating TNF alphaâ€induced hypertension in the pregnant rat FASEB Journal, 2007, 21, A592. | 0.2 | 1 |
| 88 | Humoral immune system activation promotes the development of hypertension. FASEB Journal, 2013, 27, 906.4. | 0.2 | 1 |
| 89 | The Role of Interleukinâ€2 (ILâ€2) in Natural Killer Cell (NK) Activation and Hypertension in a Preclinical Rat Model of Preeclampsia. FASEB Journal, 2018, 32, 911.1. | 0.2 | 1 |
| 90 | Salt, Aldosterone, and the Renin–Angiotensin System in Pregnancy. , 2022, , 335-353. | | 1 |

6

| # | Article | IF | CITATIONS |
|-----|---|-----|-----------|
| 91 | The Role of B Cells in Mediating Hypertension in Preeclampsia or COVIDâ€19 Infection During Pregnancy. FASEB Journal, 2022, 36, . | 0.2 | 1 |
| 92 | B Cell Depletion During Pregnancy Improves Hypertension, Natural Killer Cell Activation, and May Not Worsen Fetal Outcomes in Response to Placental Ischemia. FASEB Journal, 2021, 35, . | 0.2 | 0 |
| 93 | ILâ€17 causes hypertension and multiâ€organ tissue dysfunction which is attenuated with blockade of agonistic autoantibodies to the angiotensin II type I (AT1â€AA) receptor during pregnancy. FASEB Journal, 2021, 35, . | 0.2 | 0 |
| 94 | Progesterone and PIBF: new insights into treatment options for preeclampsia. FASEB Journal, 2021, 35, . | 0.2 | 0 |
| 95 | The Importance of B Cells in Causing Hypertension During Pregnancy; to B or Not to B. FASEB Journal, 2021, 35, . | 0.2 | 0 |
| 96 | Soluble fmsâ€like tyrosineâ€1 (sFltâ€1) production in response to placental ischemia and hypoxia in placenta of pregnant rats. FASEB Journal, 2008, 22, 969.19. | 0.2 | 0 |
| 97 | Endothelial cell activation in response to placental ischemia in pregnant rats is mediated by agonistic autoantibodies to the angiotensin type I receptor. FASEB Journal, 2008, 22, . | 0.2 | 0 |
| 98 | Agonistic autoantibodies to the angiotensin II type 1 receptor increases Angiotensin IIâ€induced ETâ€1 production. FASEB Journal, 2008, 22, 969.18. | 0.2 | 0 |
| 99 | Soluble fmsâ€like tyrosineâ€1 (sFltâ€1) is enhanced in response to chronic tumor necrosis factor―alpha excess during pregnancy. FASEB Journal, 2009, 23, 805.4. | 0.2 | 0 |
| 100 | PLACENTAL ISCHEMIA TRIGGERS IMMUNE ACTIVATION AS LEUKOCYTE OVERPRODUCTION OF SFltâ€1: A STEP IN THE PATHOGENESIS OF PREECLAMPSIA?. FASEB Journal, 2010, 24, 793.12. | 0.2 | 0 |
| 101 | Hypertension in response to agonistic autoantibodies to the angiotensin II type I receptor (AT1â€AA): role of reactive oxygen species (ROS). FASEB Journal, 2010, 24, 1025.6. | 0.2 | 0 |
| 102 | Blunting Circulating TH17 cells Decreases Hypertension and Oxidative Stress in Response to Placental Ischemia. FASEB Journal, 2013, 27, 1115.4. | 0.2 | 0 |
| 103 | Cerebral White Matter CD4+ T cell Infiltration is Associated with Hypertension in a rat model of HELLP Syndrome. FASEB Journal, 2013, 27, 691.12. | 0.2 | 0 |
| 104 | Hypertension in a rat model of HELLP Syndrome is associated with Increased TNFâ€alpha, ILâ€6 and CD4+ T cell activation. FASEB Journal, 2013, 27, 1115.7. | 0.2 | 0 |
| 105 | ILâ€10 supplementation suppressed hypertension and inflammation in response to placental ischemia during pregnancy (860.17). FASEB Journal, 2014, 28, 860.17. | 0.2 | 0 |
| 106 | T Cellâ€Dependent B Cell Activation Mediates Pathophysiology in Reponse to CD4 + T Cells from Reduced Uterine Perfusion Pregnant Rats. FASEB Journal, 2015, 29, 810.4. | 0.2 | 0 |
| 107 | Agonistic Autoantibodies to the Angiotensin II Type 1 Receptor Enhance ANGII Binding on Vascular Endothelial Cells. FASEB Journal, 2015, 29, 810.12. | 0.2 | 0 |
| 108 | Early Administration of 17â€Hydroxyprogesterone Caproate to Reduced Uterine Perfusion Pressure (RUPP) Rat Model of Preeclampsia Improves Inflammation, Uterine artery Vasoconstriction and Blood Pressure During Pregnancy. FASEB Journal, 2015, 29, 810.6. | 0.2 | 0 |

| # | Article | IF | CITATIONS |
|-----|--|-----|-----------|
| 109 | Serelaxin Improves Blood Pressure and Uterine Artery Resistance in the Reduced Uterine Perfusion Pressure (RUPP) Rat Model of Preeclampsia. FASEB Journal, 2015, 29, 810.8. | 0.2 | ο |
| 110 | Early Development of Glomerular Injury in Dahl Salt‧ensitive (SS) Rats with Metabolic Syndrome Independent of Diabetes and Hypertension. FASEB Journal, 2015, 29, 964.8. | 0.2 | 0 |
| 111 | Placental Ischemiaâ€Induced T H 17 Cells Mediate the Pathophysiology Associated with Preeclampsia. FASEB Journal, 2015, 29, 667.6. | 0.2 | Ο |
| 112 | Progesterone induced blocking factor improves fetal growth restriction possibly by reducing inflammation and placental cytolytic NK cells in response to placental ischemia during pregnancy. FASEB Journal, 2018, 32, 729.5. | 0.2 | 0 |
| 113 | Placental Ischemia‧timulated T H 17 Cells Induce Preeclampsiaâ€Associated Cytolytic Natural Killer Cells During Pregnancy. FASEB Journal, 2018, 32, 729.6. | 0.2 | 0 |
| 114 | The role of T cells on the elevated blood pressure of female and male PCOS offspring. FASEB Journal, 2019, 33, 593.5. | 0.2 | 0 |
| 115 | Interleukinâ€4 supplementation improves the proinflammatory cell ratios, autoantibodies and blood pressure in response to placental ischemia. FASEB Journal, 2019, 33, 865.18. | 0.2 | Ο |
| 116 | Angiotensin II Type I Receptor Agonistic Autoantibody Blockade Improves Cerebral Blood Flow Autoregulation, Blood Brain Barrier Permeability, and Hypertension in the Preâ€Clinical Rat Model of Preeclampsia. FASEB Journal, 2020, 34, 1-1. | 0.2 | 0 |
| 117 | CD4+ T Cells from RUPP rat model activate NK cells and cause mitochondrial oxidative stress and hypertension in normal pregnant rats. FASEB Journal, 2020, 34, 1-1. | 0.2 | 0 |
| 118 | Prevention of T Cell Activation in Response to Placental Ischemia Improves Hypertension and Natural Killer Cell Number During Pregnancy. FASEB Journal, 2020, 34, 1-1. | 0.2 | 0 |
| 119 | Progesterone induced blocking factor improves blood pressure, mitochondrial dysfunction and reactive oxygen species in response to sFltâ€l induced hypertension during pregnancy. FASEB Journal, 2020, 34, 1-1. | 0.2 | 0 |
| 120 | Maternal B Cell Depletion Reduces Blood Pressure and Improves Fetal Weights in Male Offspring of a Rat Model of Preeclampsia. FASEB Journal, 2022, 36, . | 0.2 | 0 |
| 121 | CD4+T Cells cause increased glucose, mitochondrial dysfunction, and hypertension in a Novel Pregnant Rodent Model of Gestational Diabetes Mellitus. FASEB Journal, 2022, 36, . | 0.2 | Ο |
| 122 | IL17 administration in the Absence of T cells Results in Hypertension, NK cell Activation, and Reduced Pup Weight at Birth, but No Changes in Blood Pressure or Weight at Maturation of Offspring. FASEB Journal, 2022, 36, . | 0.2 | 0 |
| 123 | Progesterone prolongs time to delivery and attenuates blood pressure possibly by improving inflammation and endothelial function in response to preeclampsia. FASEB Journal, 2022, 36, . | 0.2 | 0 |