

# Xose Souto

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3997163/publications.pdf>

Version: 2024-02-01

27  
papers

841  
citations

623734

14  
h-index

610901

24  
g-index

27  
all docs

27  
docs citations

27  
times ranked

990  
citing authors

#	ARTICLE	IF	CITATIONS
1	Effect of phenolic compounds on the germination of six weeds species. <i>Plant Growth Regulation</i> , 1999, 28, 83-88.	3.4	152
2	Title is missing!. <i>Journal of Chemical Ecology</i> , 2000, 26, 2025-2034.	1.8	77
3	Allelopathic effects of <i>Acacia melanoxylon</i> R.Br. phyllodes during their decomposition. <i>Forest Ecology and Management</i> , 1995, 77, 53-63.	3.2	60
4	Critical environmental and genotypic factors for <i>Fusarium verticillioides</i> infection, fungal growth and fumonisin contamination in maize grown in northwestern Spain. <i>International Journal of Food Microbiology</i> , 2014, 177, 63-71.	4.7	59
5	Comparative analysis of allelopathic effects produced by four forestry species during decomposition process in their soils in Galicia (NW Spain). <i>Journal of Chemical Ecology</i> , 1994, 20, 3005-3015.	1.8	57
6	Allelopathic Effects of Tree Species on Some Soil Microbial Populations and Herbaceous Plants. <i>Biologia Plantarum</i> , 2001, 44, 269-275.	1.9	53
7	Allelopathic Effects of Humus Phenolics on Growth and Respiration of Mycorrhizal Fungi. <i>Journal of Chemical Ecology</i> , 2000, 26, 2015-2023.	1.8	50
8	Putative Role of Pith Cell Wall Phenylpropanoids in <i>Sesamia nonagrioides</i> (Lepidoptera: Noctuidae) Resistance. <i>Journal of Agricultural and Food Chemistry</i> , 2006, 54, 2274-2279.	5.2	49
9	Ecophysiological responses of three native herbs to phytotoxic potential of invasive <i>Acacia melanoxylon</i> R. Br.. <i>Agroforestry Systems</i> , 2011, 83, 149-166.	2.0	42
10	Allelopathy in Northern Temperate and Boreal Semi-Natural Woodland. <i>Critical Reviews in Plant Sciences</i> , 1999, 18, 637-652.	5.7	34
11	Diferulate Content of Maize Sheaths Is Associated with Resistance to the Mediterranean Corn Borer <i>Sesamia nonagrioides</i> (Lepidoptera: Noctuidae). <i>Journal of Agricultural and Food Chemistry</i> , 2006, 54, 9140-9144.	5.2	33
12	Free Phenols in Maize Pith and Their Relationship with Resistance to <i>Sesamia nonagrioides</i> (Lepidoptera: Noctuidae) Attack. <i>Journal of Economic Entomology</i> , 2005, 98, 1349-1356.	1.8	31
13	Relationship Between Maize Stem Structural Characteristics and Resistance to Pink Stem Borer (Lepidoptera: Noctuidae) Attack. <i>Journal of Economic Entomology</i> , 2003, 96, 1563-1570.	1.8	24
14	Assessing white maize resistance to fumonisin contamination. <i>European Journal of Plant Pathology</i> , 2014, 138, 283-292.	1.7	18
15	Water-soluble phenolic acids and flavonoids involved in the bioherbicidal potential of <i>Ulex europaeus</i> and <i>Cytisus scoparius</i> . <i>South African Journal of Botany</i> , 2020, 133, 201-211.	2.5	17
16	Allelopathy in forest ecosystems. , 2000, , 183-193.		16
17	Role of Hydroxycinnamic Acids in the Infection of Maize Silks by <i>Fusarium graminearum</i> Schwabe. <i>Molecular Plant-Microbe Interactions</i> , 2011, 24, 1020-1026.	2.6	15
18	Methods for Determining Cell Wall-Bound Phenolics in Maize Stem Tissues. <i>Journal of Agricultural and Food Chemistry</i> , 2018, 66, 1279-1284.	5.2	14

#	ARTICLE	IF	CITATIONS
19	Relationship Between Maize Stem Structural Characteristics and Resistance to Pink Stem Borer (Lepidoptera: Noctuidae) Attack. <i>Journal of Economic Entomology</i> , 2003, 96, 1563-1570.	1.8	13
20	Effect of Maize Pith Free Phenols on Larval Growth and Development of <i>Sesamia nonagrioides</i> (Lepidoptera: Noctuidae). <i>Journal of Entomology</i> , 2006, 3, 281-289.	0.2	7
21	Predictive phytotoxic value of water-soluble allelochemicals in plant extracts for choosing a cover crop or mulch for specific weed control. <i>Italian Journal of Agronomy</i> , 2021, 16, .	1.0	6
22	Chemical Changes during Maize Tissue Aging and Its Relationship with Mediterranean Corn Borer Resistance. <i>Journal of Agricultural and Food Chemistry</i> , 2017, 65, 9180-9185.	5.2	5
23	Rye ( <i>Secale cereale</i> L.) and squarrose clover ( <i>Trifolium squarrosum</i> L.) cover crops can increase their allelopathic potential for weed control when used mixed as dead mulch. <i>Italian Journal of Agronomy</i> , 0, , .	1.0	3
24	Elucidating the multifunctional role of the cell wall components in the maize exploitation. <i>BMC Plant Biology</i> , 2021, 21, 251.	3.6	2
25	Feedback mechanism in the chemical ecology of plants: role of soil microorganisms. , 2002, , 89-97.		2
26	Allelopathic Effects of Exotic Tree Species on Microorganisms and Plants in Galicia (Spain). <i>Forestry Sciences</i> , 1998, , 293-300.	0.4	1
27	Identification of single nucleotide polymorphisms (SNPs) for maize cell wall hydroxycinnamates using a multi-parent advanced generation intercross (MAGIC) population. <i>Phytochemistry</i> , 2022, 193, 113002.	2.9	1