Patrizia Giannatempo

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/4003828/publications.pdf Version: 2024-02-01



#	Article	lF	CITATIONS
1	Pembrolizumab as Neoadjuvant Therapy Before Radical Cystectomy in Patients With Muscle-Invasive Urothelial Bladder Carcinoma (PURE-01): An Open-Label, Single-Arm, Phase II Study. Journal of Clinical Oncology, 2018, 36, 3353-3360.	1.6	474
2	ESMO Consensus Conference on testicular germ cell cancer: diagnosis, treatment and follow-up. Annals of Oncology, 2018, 29, 1658-1686.	1.2	228
3	Updated Results of PURE-01 with Preliminary Activity of Neoadjuvant Pembrolizumab in Patients with Muscle-invasive Bladder Carcinoma with Variant Histologies. European Urology, 2020, 77, 439-446.	1.9	228
4	Pazopanib in advanced and platinum-resistant urothelial cancer: an open-label, single group, phase 2 trial. Lancet Oncology, The, 2012, 13, 810-816.	10.7	130
5	Impact of Molecular Subtyping and Immune Infiltration on Pathological Response and Outcome Following Neoadjuvant Pembrolizumab in Muscle-invasive Bladder Cancer. European Urology, 2020, 77, 701-710.	1.9	128
6	Second-line single-agent versus doublet chemotherapy as salvage therapy for metastatic urothelial cancer: a systematic review and meta-analysis. Annals of Oncology, 2016, 27, 49-61.	1.2	108
7	The Relationship between Characteristics of Inguinal Lymph Nodes and Pelvic Lymph Node Involvement in Penile Squamous Cell Carcinoma: A Single Institution Experience. Journal of Urology, 2014, 191, 977-982.	0.4	75
8	Multiparametric Magnetic Resonance Imaging as a Noninvasive Assessment of Tumor Response to Neoadjuvant Pembrolizumab in Muscle-invasive Bladder Cancer: Preliminary Findings from the PURE-01 Study. European Urology, 2020, 77, 636-643.	1.9	75
9	Radiotherapy or chemotherapy for clinical stage IIA and IIB seminoma: a systematic review and meta-analysis of patient outcomes. Annals of Oncology, 2015, 26, 657-668.	1.2	71
10	Treatment and Clinical Outcomes of Patients with Teratoma with Somatic-Type Malignant Transformation: An International Collaboration. Journal of Urology, 2016, 196, 95-100.	0.4	70
11	Firstâ€line therapy with dacomitinib, an orally available panâ€HER tyrosine kinase inhibitor, for locally advanced or metastatic penile squamous cell carcinoma: results of an openâ€label, singleâ€arm, singleâ€centre, phase 2 study. BJU International, 2018, 121, 348-356.	2.5	70
12	Brain Metastases in Patients With Germ Cell Tumors: Prognostic Factors and Treatment Options—An Analysis From the Global Germ Cell Cancer Group. Journal of Clinical Oncology, 2016, 34, 345-351.	1.6	69
13	A Combination of Cisplatin and 5-Fluorouracil With a Taxane in Patients Who Underwent Lymph Node Dissection for Nodal Metastases From Squamous Cell Carcinoma of the Penis: Treatment Outcome and Survival Analyses in Neoadjuvant and Adjuvant Settings. Clinical Genitourinary Cancer, 2016, 14, 323-330	1.9	59
14	Comparison of the Prognostic Value of Assessing Tumor Diameter Versus Tumor Volume at Diagnosis or in Response to Initial Chemotherapy in Rhabdomyosarcoma. Journal of Clinical Oncology, 2010, 28, 1322-1328.	1.6	58
15	An Open-label Randomized Phase 2 study of Durvalumab Alone or in Combination with Tremelimumab in Patients with Advanced Germ Cell Tumors (APACHE): Results from the First Planned Interim Analysis. European Urology, 2019, 75, 201-203.	1.9	54
16	PF-03446962, a fully-human monoclonal antibody against transforming growth-factor β (TGFβ) receptor ALK1, in pre-treated patients with urothelial cancer: an open label, single-group, phase 2 trial. Investigational New Drugs, 2014, 32, 555-560.	2.6	50
17	Efficacy and Safety of Gemcitabine Plus Either Taxane or Carboplatin in the First-Line Setting of Metastatic Urothelial Carcinoma: A Systematic Review and Meta-Analysis. Clinical Genitourinary Cancer, 2017, 15, 23-30.e2.	1.9	50
18	Clinical Outcome in Testicular Sex Cord Stromal Tumors: Testis Sparing vs Radical Orchiectomy and Management of Advanced Disease. Urology, 2015, 85, 402-406.	1.0	47

#	Article	IF	CITATIONS
19	Panitumumab Treatment for Advanced Penile Squamous Cell Carcinoma When Surgery and Chemotherapy Have Failed. Clinical Genitourinary Cancer, 2016, 14, 231-236.	1.9	38
20	Clinical Outcomes of Perioperative Chemotherapy in Patients With Locally Advanced Penile Squamous-Cell Carcinoma: Results of a Multicenter Analysis. Clinical Genitourinary Cancer, 2017, 15, 548-555.e3.	1.9	37
21	Patients Selection for Immunotherapy in Solid Tumors: Overcome the NaÃ⁻ve Vision of a Single Biomarker. BioMed Research International, 2019, 2019, 1-15.	1.9	37
22	Persistent CD30 Expression by Embryonal Carcinoma in the Treatment Time Course: Prognostic Significance of a Worthwhile Target for Personalized Treatment. Journal of Urology, 2013, 190, 1919-1924.	0.4	36
23	Pazopanib in advanced germ cell tumors after chemotherapy failure: results of the open-label, single-arm, phase 2 Pazotest trial. Annals of Oncology, 2017, 28, 1346-1351.	1.2	34
24	Predicting the Pathologic Complete Response After Neoadjuvant Pembrolizumab in Muscle-Invasive Bladder Cancer. Journal of the National Cancer Institute, 2021, 113, 48-53.	6.3	30
25	Prognostic Factors of Adjuvant Taxane, Cisplatin, and 5-Fluorouracil Chemotherapy for Patients With Penile Squamous Cell Carcinoma After Regional Lymphadenectomy. Clinical Genitourinary Cancer, 2016, 14, 518-523.	1.9	28
26	Combination of Paclitaxel, Cisplatin, and Gemcitabine (TPG) for Multiple Relapses or Platinum-Resistant Germ Cell Tumors: Long-Term Outcomes. Clinical Genitourinary Cancer, 2014, 12, 63-69.e1.	1.9	27
27	Clinical experience with temsirolimus in the treatment of advanced renal cell carcinoma. Therapeutic Advances in Urology, 2015, 7, 152-161.	2.0	27
28	A Prognostic Model Including Pre- and Postsurgical Variables to Enhance Risk Stratification of Primary Mediastinal Nonseminomatous Germ Cell Tumors: The 27-Year Experience of a Referral Center. Clinical Genitourinary Cancer, 2015, 13, 87-93.e1.	1.9	27
29	Effect of Bleomycin Administration on the Development of Pulmonary Toxicity in Patients With Metastatic Germ Cell Tumors Receiving First-Line Chemotherapy: A Meta-Analysis of Randomized Studies. Clinical Genitourinary Cancer, 2017, 15, 213-220.e5.	1.9	27
30	Impact of the Number of Cycles of Platinum Based First Line Chemotherapy for Advanced Urothelial Carcinoma. Journal of Urology, 2018, 200, 1207-1214.	0.4	26
31	The Impact of Adding Taxanes to Gemcitabine and Platinum Chemotherapy for the First-Line Therapy of Advanced or Metastatic Urothelial Cancer: A Systematic Review and Meta-analysis. European Urology, 2016, 69, 624-633.	1.9	25
32	A Suggested Prognostic Reclassification of Intermediate and Poor-Risk Nonseminomatous Germ Cell Tumors. Clinical Genitourinary Cancer, 2017, 15, 306-312.e3.	1.9	25
33	Interim Fluorine-18 Fluorodeoxyglucose Positron Emission Tomography for Early Metabolic Assessment of Therapeutic Response to Chemotherapy for Metastatic Transitional Cell Carcinoma. Clinical Genitourinary Cancer, 2014, 12, 433-439.	1.9	24
34	Relationship between lymph node ratio and cancer-specific survival in a contemporary series of patients with penile cancer and lymph node metastases. BJU International, 2015, 116, 727-733.	2.5	23
35	Can Patients with Muscle-invasive Bladder Cancer and Fibroblast Growth Factor Receptor-3 Alterations Still Be Considered for Neoadjuvant Pembrolizumab? A Comprehensive Assessment from the Updated Results of the PURE-01 Study. European Urology Oncology, 2021, 4, 1001-1005.	5.4	23
36	Brentuximab Vedotin in CD30-Expressing Germ Cell Tumors After Chemotherapy Failure. Clinical Genitourinary Cancer, 2016, 14, 261-264.e4.	1.9	22

Patrizia Giannatempo

#	Article	IF	CITATIONS
37	Association Between Human Papillomavirus Infection and Outcome of Perioperative Nodal Radiotherapy for Penile Carcinoma. European Urology Oncology, 2021, 4, 802-810.	5.4	22
38	An open-label, single-arm, phase 2 study of the Aurora kinase A inhibitor alisertib in patients with advanced urothelial cancer. Investigational New Drugs, 2016, 34, 236-242.	2.6	21
39	Laparoscopic Retroperitoneal Lymph Node Dissection for Clinical Stage I Nonseminomatous Germ Cell Tumors of the Testis: Safety and Efficacy Analyses at a High Volume Center. Journal of Urology, 2018, 199, 741-747.	0.4	21
40	Laparoscopic Postchemotherapy Retroperitoneal Lymph-Node Dissection Can Be a Standard Option in Defined Nonseminomatous Germ Cell Tumor Patients. Journal of Endourology, 2016, 30, 1112-1119.	2.1	20
41	The Value of Multiparametric Magnetic Resonance Imaging Sequences to Assist in the Decision Making of Muscle-invasive Bladder Cancer. European Urology Oncology, 2021, 4, 829-833.	5.4	20
42	Palliative treatment of germ cell cancer. Cancer Treatment Reviews, 2018, 71, 102-107.	7.7	19
43	Gene Expression Profiling of Advanced Penile Squamous Cell Carcinoma Receiving Cisplatin-based Chemotherapy Improves Prognostication and Identifies Potential Therapeutic Targets. European Urology Focus, 2018, 4, 733-736.	3.1	18
44	Postchemotherapy Lymphadenectomy in Patients With Metastatic Urothelial Carcinoma: Long-Term Efficacy and Implications for Trial Design. Clinical Genitourinary Cancer, 2015, 13, 80-86.e1.	1.9	17
45	High-dose sequential chemotherapy (HDS) versus PEB chemotherapy as first-line treatment of patients with poor prognosis germ-cell tumors: mature results of an Italian randomized phase II study. Annals of Oncology, 2015, 26, 167-172.	1.2	17
46	Erdafitinib for the treatment of urothelial cancer. Expert Review of Anticancer Therapy, 2019, 19, 835-846.	2.4	17
47	Analysis of plasma cytokines and angiogenic factors in patients with pretreated urothelial cancer receiving Pazopanib: the role of circulating interleukin-8 to enhance the prognostic accuracy. British Journal of Cancer, 2014, 110, 26-33.	6.4	16
48	Outcome of patients with advanced upper tract urothelial carcinoma treated with immune checkpoint inhibitors: A systematic review and meta-analysis. Critical Reviews in Oncology/Hematology, 2021, 159, 103241.	4.4	16
49	Long-Term Efficacy and Safety Outcomes of Modified (Simplified) MVAC (Methotrexate/Vinblastine/Doxorubicin/Cisplatin) as Frontline Therapy for Unresectable or Metastatic Urothelial Cancer. Clinical Genitourinary Cancer, 2014, 12, 203-209.e1.	1.9	15
50	Cisplatin-Based First-Line Therapy for Advanced Urothelial Carcinoma After Previous Perioperative Cisplatin-Based Therapy. Clinical Genitourinary Cancer, 2015, 13, 178-184.	1.9	15
51	Modified cisplatin, etoposide, and ifosfamide (PEI) salvage therapy for male germ cell tumors: long-term efficacy and safety outcomes. Annals of Oncology, 2013, 24, 2887-2892.	1.2	14
52	Clinical Significance of Early Changes inÂCirculating Tumor Cells from Patients Receiving First-Line Cisplatin-Based Chemotherapy for Metastatic Urothelial Carcinoma1. Bladder Cancer, 2016, 2, 395-403.	0.4	13
53	Epidemiology and unmet needs of bladder cancer in Italy: a critical review. Minerva Urologica E Nefrologica = the Italian Journal of Urology and Nephrology, 2020, 72, 1-12.	3.9	13
54	Prognostic reclassification of patients with intermediate-risk metastatic germ cell tumors: Implications for clinical practice, trial design, and molecular interrogation. Urologic Oncology: Seminars and Original Investigations, 2015, 33, 332.e19-332.e24.	1.6	12

#	Article	IF	CITATIONS
55	Interim 18F-Fluorodeoxyglucose Positron Emission Tomography for Early Metabolic Assessment of Response to Cisplatin, Etoposide, and Bleomycin Chemotherapy for Metastatic Seminoma: Clinical Value and Future Directions. Clinical Genitourinary Cancer, 2016, 14, 249-254.	1.9	12
56	Cisplatin- Versus Non–Cisplatin-based First-Line Chemotherapy for Advanced Urothelial Carcinoma Previously Treated With Perioperative Cisplatin. Clinical Genitourinary Cancer, 2016, 14, 331-340.	1.9	12
57	Association of Androgen Receptor Expression on Tumor Cells and PD-L1 Expression in Muscle-Invasive and Metastatic Urothelial Carcinoma: Insights for Clinical Research. Clinical Genitourinary Cancer, 2018, 16, e403-e410.	1.9	11
58	Neoadjuvant Chemotherapy or Immunotherapy for Clinical T2N0 Muscle-invasive Bladder Cancer: Time to Change the Paradigm?. European Urology Oncology, 2021, 4, 1006-1010.	5.4	11
59	An open-label, single-group, phase 2 study of brentuximab vedotin as salvage therapy for males with relapsed germ-cell tumors (GCT): Results at the end of first stage (FM12GCT01) Journal of Clinical Oncology, 2016, 34, 480-480.	1.6	11
60	[18F]Fluoro-Deoxy-Glucose positron emission tomography to evaluate lymph node involvement in patients with muscle-invasive bladder cancer receiving neoadjuvant pembrolizumab. Urologic Oncology: Seminars and Original Investigations, 2021, 39, 235.e15-235.e21.	1.6	10
61	Immunohistochemistry to Enhance Prognostic Allocation and Guide Decision-Making of Patients With Advanced Urothelial Cancer Receiving First-Line Chemotherapy. Clinical Genitourinary Cancer, 2015, 13, 171-177.e1.	1.9	9
62	Association of an aurora kinase a (AURKA) gene polymorphism with progression-free survival in patients with advanced urothelial carcinoma treated with the selective aurora kinase a inhibitor alisertib. Investigational New Drugs, 2017, 35, 524-528.	2.6	9
63	Exceptional response to olaparib in BRCA2-altered urothelial carcinoma after PD-L1 inhibitor and chemotherapy failure. European Journal of Cancer, 2018, 96, 128-130.	2.8	9
64	Neoadjuvant sorafenib, gemcitabine, and cisplatin administration preceding cystectomy in patients with muscle-invasive urothelial bladder carcinoma: An open-label, single-arm, single-center, phase 2 study. Urologic Oncology: Seminars and Original Investigations, 2018, 36, 8.e1-8.e8.	1.6	8
65	Prognostic Effect of FGFR Mutations or Gene Fusions in Patients with Metastatic Urothelial Carcinoma Receiving First-line Platinum-based Chemotherapy: Results from a Large, Single-institution Cohort. European Urology Focus, 2019, 5, 853-856.	3.1	8
66	Prognostic impact of progression to induction chemotherapy and prior paclitaxel therapy in patients with germ cell tumors receiving salvage high-dose chemotherapy in the last 10 years: a study of the European Society for Blood and Marrow Transplantation Solid Tumors Working Party. Bone Marrow Transplantation, 2016, 51, 384-390.	2.4	7
67	Treatment of Carcinoma In Situ of the Glans Penis With Topical Imiquimod Followed by Carbon Dioxide Laser Excision. Clinical Genitourinary Cancer, 2017, 15, e483-e487.	1.9	7
68	Quality of Life and Pain Control following Laparoscopic Retroperitoneal Lymph Node Dissection in Early-stage Nonseminoma. Tumori, 2015, 101, 650-656.	1.1	6
69	The Changing Landscape of Intermediate- and Poor-Risk Germ Cell Tumors: Do We Need to Reclassify Patients With Metastatic Germ Cell Tumors?. Clinical Genitourinary Cancer, 2016, 14, 1-4.	1.9	6
70	Immunotherapy for metastatic urothelial carcinoma. Current Opinion in Urology, 2018, 28, 1-7.	1.8	6
71	Apache: An open label, randomized, phase 2 study of durvalumab (Durva), alone or in combination with tremelimumab (Treme), in patients (pts) with advanced germ cell tumors (GCT): Results at the end of first stage Journal of Clinical Oncology, 2018, 36, 4547-4547.	1.6	6
72	Secondary malignancies after high-dose chemotherapy in germ cell tumor patients: a 34-year retrospective study of the European Society for Blood and Marrow Transplantation (EBMT). Bone Marrow Transplantation, 2018, 53, 722-728.	2.4	5

#	Article	IF	CITATIONS
73	Recommendations for surveillance and follow-up of men with testicular germ cell tumors: a multidisciplinary consensus conference by the Italian Germ cell cancer Group and the Associazione Italiana di Oncologia Medica. Critical Reviews in Oncology/Hematology, 2019, 137, 154-164.	4.4	5
74	Prognostic Role of Early Interim Fluorodeoxyglucose Positron Emission Tomography in Patients With Advanced Seminoma Undergoing Standard Treatment. Clinical Genitourinary Cancer, 2021, 19, 237-245.e2.	1.9	5
75	Activity of pazopanib in chemo-resistant patients with germ cell tumors (GCT): First results of the open-label, single-group, phase II PAZOTEST-01 trial Journal of Clinical Oncology, 2014, 32, 376-376.	1.6	5
76	Nausea and Vomiting during the First 3 Intercycle Periods in Chemo-naive Cancer Patients Receiving Moderately/Highly Emetogenic Therapy. Tumori, 2015, 101, 692-696.	1.1	4
77	Clinical Outcomes of Metastatic Poor Prognosis Germ Cell Tumors: Current Perspective From a Referral Center. Clinical Genitourinary Cancer, 2015, 13, 385-391.e1.	1.9	4
78	Molecular Signature of Response to Pazopanib Salvage Therapy for Urothelial Carcinoma. Clinical Genitourinary Cancer, 2016, 14, e81-e90.	1.9	4
79	Etoposide, Methotrexate, and Dactinomycin Alternating With Cyclophosphamide and Vincristine (EMACO) for Male Patients With HCG-expressing, Chemoresistant Germ Cell Tumors. American Journal of Clinical Oncology: Cancer Clinical Trials, 2017, 40, 60-65.	1.3	4
80	Incidence and Clinical Impact of Inflammatory Fluorodeoxyglucose Positron Emission Tomography Uptake After Neoadjuvant Pembrolizumab in Patients with Organ-confined Bladder Cancer Undergoing Radical Cystectomy. European Urology Focus, 2021, 7, 1092-1099.	3.1	4
81	Administration of high-dose chemotherapy with stem cell support in patients 40 years of age or older with advanced germ cell tumours: a retrospective study from the European Society for Blood and Marrow Transplantation database. Bone Marrow Transplantation, 2017, 52, 1218-1220.	2.4	4
82	Immunotherapy and Sonpavde score validation in advanced upper tract urothelial carcinoma: a retrospective study by the Italian Network for Research in Urologic-Oncology (Meet-URO group). Immunotherapy, 2022, 14, 107-114.	2.0	4
83	Erdafitinib in locally advanced/metastatic urothelial carcinoma with certain <i>FGFR</i> genetic alterations. Future Oncology, 2022, 18, 2455-2464.	2.4	4
84	Predictors of CD34+ Cell Mobilization and Collection in Adult Men With Germ Cell Tumors: Implications for the Salvage Treatment Strategy. Clinical Genitourinary Cancer, 2014, 12, 196-202.e1.	1.9	3
85	Salvage High-Dose Chemotherapy for Relapsed Pure Seminoma in the Last 10 Years: Results From the European Society for Blood and Marrow Transplantation Series 2002-2012. Clinical Genitourinary Cancer, 2017, 15, 163-167.	1.9	3
86	An evaluation of UGN-101, a sustained-release hydrogel polymer-based formulation containing mitomycin-C, for the treatment of upper urothelial carcinomas. Expert Opinion on Pharmacotherapy, 2020, 21, 2199-2204.	1.8	3
87	Etoposide, methotrexate, and dactinomycin alternating with cyclophosphamide and vincristine (EMA/CO) for males with HCG-expressing, chemorefractory germ cell tumors (GCT): Long-term efficacy and safety outcomes Journal of Clinical Oncology, 2014, 32, 4561-4561.	1.6	2
88	Biological Therapeutic Advances for the Treatment of Advanced Urothelial Cancers. Biologics: Targets and Therapy, 2021, Volume 15, 441-450.	3.2	2
89	Impact of Prior Platinum-Based Therapy on Patients Receiving Salvage Systemic Treatment for Advanced Urothelial Carcinoma. Clinical Genitourinary Cancer, 2016, 14, 494-498.	1.9	1
90	From the Uncertainties to the Evidence: A Brief History of Immunotherapy as Salvage Therapy for Advanced Bladder Cancer Through a Meta-analysis. Clinical Genitourinary Cancer, 2017, 15, 509-512.e9.	1.9	1

#	Article	IF	CITATIONS
91	Nivolumab and its use in the second-line treatment of metastatic urothelial cancer. Future Oncology, 2018, 14, 2683-2690.	2.4	1
92	Comparison of Fibroblast Growth-factor Receptor Gene Alterations at the DNA versus Messenger RNA Level in Advanced Urothelial Cancer: Insights for Clinical Research. European Urology Focus, 2019, 5, 689-692.	3.1	1
93	Prostatic metastases from testicular nonseminomatous germ cell cancer: two case reports and a review of the literature. Tumori, 2013, 99, e203-7.	1.1	1
94	New Directions for Biologic Targets in Urothelial Carcinoma – Letter. Molecular Cancer Therapeutics, 2012, 11, 2306-2306.	4.1	0
95	Role of Neoadjuvant and Adjuvant Chemotherapy in Penile Cancer. , 2018, , 1-6.		0
96	Role of Neoadjuvant and Adjuvant Chemotherapy in Penile Cancer. , 2019, , 845-850.		0