

# Hyung Jik Kim

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4013674/publications.pdf>

Version: 2024-02-01

42  
papers

678  
citations

759233

12  
h-index

580821

25  
g-index

42  
all docs

42  
docs citations

42  
times ranked

1148  
citing authors

#	ARTICLE	IF	CITATIONS
1	Serum cystatin C to creatinine ratio is associated with sarcopenia in non-dialysis-dependent chronic kidney disease. <i>Kidney Research and Clinical Practice</i> , 2022, 41, 580-590.	2.2	10
2	Comparison of dynamic arterial and venous pressure between metal needles and plastic cannulas in incident hemodialysis patients with arteriovenous graft. <i>Journal of Vascular Access</i> , 2021, 22, 42-47.	0.9	4
3	Neutrophil extracellular traps and heparin-induced antibodies contribute to vascular access thrombosis in hemodialysis patients. <i>Kidney Research and Clinical Practice</i> , 2021, 40, 712-723.	2.2	1
4	Late stage 3 chronic kidney disease is an independent risk factor for sarcopenia, but not proteinuria. <i>Scientific Reports</i> , 2021, 11, 18472.	3.3	16
5	Prognostic role of circulating neutrophil extracellular traps levels for long-term mortality in new end-stage renal disease patients. <i>Clinical Immunology</i> , 2020, 210, 108263.	3.2	21
6	Incidence of malignancy and related mortality after kidney transplantation: a nationwide, population-based cohort study in Korea. <i>Scientific Reports</i> , 2020, 10, 21398.	3.3	14
7	P1580LONGITUDINAL CHANGES IN BODY COMPOSITION ARE ASSOCIATED WITH ALL-CAUSE MORTALITY IN PATIENTS ON PERITONEAL DIALYSIS. <i>Nephrology Dialysis Transplantation</i> , 2020, 35, .	0.7	0
8	Role of peribrachial fat as a key determinant of brachial artery dilatation for successful arteriovenous fistula maturation in hemodialysis patients. <i>Scientific Reports</i> , 2020, 10, 3841.	3.3	0
9	Impact of sarcopenia on long-term mortality and cardiovascular events in patients undergoing hemodialysis. <i>Korean Journal of Internal Medicine</i> , 2019, 34, 599-607.	1.7	79
10	SP407Serum protein carbonyl levels are associated with sarcopenia and long-term outcomes in hemodialysis patients. <i>Nephrology Dialysis Transplantation</i> , 2019, 34, .	0.7	0
11	FP728LOW LEAN TISSUE AND HIGH FAT TISSUE INDEX ARE ASSOCIATED WITH DECREASED QUALITY OF LIFE IN DIALYSIS PATIENTS. <i>Nephrology Dialysis Transplantation</i> , 2019, 34, .	0.7	0
12	Custom-made Doppler ultrasound flow simulator for dialysis access using continuous renal replacement therapy machine. <i>Journal of Vascular Access</i> , 2019, 20, 701-705.	0.9	1
13	Effects of Excessive Body Fat Accumulation on Long-Term Outcomes during Peritoneal Dialysis. <i>Peritoneal Dialysis International</i> , 2019, 39, 268-275.	2.3	13
14	Anatomical variants of upper arm veins on preoperative mapping venography for hemodialysis access in Korean adults. <i>Journal of Vascular Access</i> , 2019, 20, 270-275.	0.9	7
15	Serum alkaline phosphatase and $\hat{1}^3$ -glutamyl transferase in acute pyelonephritis. <i>Kidney Research and Clinical Practice</i> , 2019, 38, 205-211.	2.2	2
16	Outcomes of vascular access in hemodialysis patients: Analysis based on the Korean National Health Insurance Database from 2008 to 2016. <i>Kidney Research and Clinical Practice</i> , 2019, 38, 391-398.	2.2	13
17	FP505THE SIGNIFICANCE OF REPEATED FLUID STATUS MEASUREMENTS BY BIOIMPEDANCE ANALYSIS FOR PREDICTING PROGNOSIS OF PERITONELA DIALYSIS PATIENTS. <i>Nephrology Dialysis Transplantation</i> , 2018, 33, i207-i207.	0.7	0
18	Comparison of tolvaptan treatment between patients with the SIADH and congestive heart failure: a single-center experience. <i>Korean Journal of Internal Medicine</i> , 2018, 33, 561-567.	1.7	7

#	ARTICLE	IF	CITATIONS
19	The relationship between autophagy, increased neutrophil extracellular traps formation and endothelial dysfunction in chronic kidney disease. <i>Clinical Immunology</i> , 2018, 197, 189-197.	3.2	16
20	Fat-to-Lean Mass Ratio Can Predict Cardiac Events and All-Cause Mortality in Patients Undergoing Hemodialysis. <i>Annals of Nutrition and Metabolism</i> , 2018, 73, 241-249.	1.9	21
21	Long-term prognosis of end-stage renal disease patients with normal myocardial perfusion as determined by single photon emission computed tomography. <i>Korean Journal of Internal Medicine</i> , 2018, 33, 148-156.	1.7	3
22	Current treatment status and medical costs for hemodialysis vascular access based on analysis of the Korean Health Insurance Database. <i>Korean Journal of Internal Medicine</i> , 2018, 33, 1160-1168.	1.7	11
23	Impact of rapid ultrafiltration rate on changes in the echocardiographic left atrial volume index in patients undergoing haemodialysis: a longitudinal observational study. <i>BMJ Open</i> , 2017, 7, e013990.	1.9	8
24	Vitamin D: a possible modifying factor linking obesity to vascular calcification in hemodialysis patients. <i>Nutrition and Metabolism</i> , 2017, 14, 27.	3.0	7
25	Increased Neutrophil Extracellular Trap Formation in Uremia Is Associated with Chronic Inflammation and Prevalent Coronary Artery Disease. <i>Journal of Immunology Research</i> , 2017, 2017, 1-10.	2.2	28
26	Acute kidney injury can predict in-hospital and long-term mortality in elderly patients undergoing hip fracture surgery. <i>PLoS ONE</i> , 2017, 12, e0176259.	2.5	28
27	SP475RAPID ULTRAFILTRATION RATE IS ASSOCIATED WITH THE INCREASE OF ECHOCARDIOGRAPHIC LEFT ATRIAL VOLUME INDEX AMONG HEMODIALYSIS PATIENTS. <i>Nephrology Dialysis Transplantation</i> , 2016, 31, i251-i252.	0.7	0
28	SP622LOW VITAMIN D AS A MODIFYING FACTOR IN THE RELATIONSHIP BETWEEN OBESITY AND VASCULAR CALCIFICATION IN HEMODIALYSIS PATIENTS. <i>Nephrology Dialysis Transplantation</i> , 2016, 31, i302-i303.	0.7	0
29	Leptin, pre-existing vascular disease, and increased arteriovenous fistula maturation failure in dialysis patients. <i>Journal of Vascular Surgery</i> , 2016, 64, 402-410.e1.	1.1	10
30	SP483EXCESS WEIGHT GAIN DURING THE FIRST YEAR OF PERITONEAL DIALYSIS IS A RISK FACTOR FOR A RAPID DECREASE IN RESIDUAL RENAL FUNCTION. <i>Nephrology Dialysis Transplantation</i> , 2015, 30, iii539-iii539.	0.7	0
31	Therapeutic Plasma Exchange Using the Spectra Optia Cell Separator Compared With the COBE Spectra. <i>Annals of Laboratory Medicine</i> , 2015, 35, 506-509.	2.5	11
32	A Low Baseline Glomerular Filtration Rate Predicts Poor Clinical Outcome at 3 Months after Acute		

#	ARTICLE	IF	CITATIONS
37	Increased body fat rather than body weight has harmful effects on 4-year changes of renal function in the general elderly population with a normal or mildly impaired renal function. <i>Clinical Interventions in Aging</i> , 2014, 9, 1277.	2.9	9
38	B-type natriuretic peptide predicts an ischemic etiology of acute heart failure in patients with stage 4-5 chronic kidney disease. <i>Clinical Biochemistry</i> , 2014, 47, 344-348.	1.9	7
39	Prevalence of and factors associated with sarcopenia in elderly patients with end-stage renal disease. <i>Clinical Nutrition</i> , 2014, 33, 64-68.	5.0	212
40	A Case of Hemocholecystitis in a Patient with Microscopic Polyangiitis. <i>Korean Journal of Medicine</i> , 2012, 83, 127.	0.3	0
41	Could Solutions Low in Glucose Degradation Products Preserve Residual Renal Function in Incident Peritoneal Dialysis Patients? A 1-Year Multicenter Prospective Randomized Controlled Trial (Balnet) <i>Tj ETQq1 1 0.7843 14 rgB74Overlock</i>	0.7843	14
42	The relationship between plasma leptin and nutritional status in chronic hemodialysis patients. <i>Journal of Korean Medical Science</i> , 1999, 14, 546.	2.5	6