

Satoshi Gando

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4083361/publications.pdf>

Version: 2024-02-01

241
papers

12,313
citations

36691

53
h-index

34195

103
g-index

245
all docs

245
docs citations

245
times ranked

11806
citing authors

#	ARTICLE	IF	CITATIONS
1	Pathomechanisms Underlying Hypoxemia in Two COVID-19-Associated Acute Respiratory Distress Syndrome Phenotypes: Insights From Thrombosis and Hemostasis. <i>Shock</i> , 2022, 57, 1-6.	1.0	9
2	Age-related differences in the survival benefit of the administration of antithrombin, recombinant human thrombomodulin, or their combination in sepsis. <i>Scientific Reports</i> , 2022, 12, .	1.6	7
3	Current spectrum of causative pathogens in sepsis: A prospective nationwide cohort study in Japan. <i>International Journal of Infectious Diseases</i> , 2021, 103, 343-351.	1.5	20
4	Thromboplasmininflammation in COVID-19 coagulopathy. <i>Japanese Journal of Thrombosis and Hemostasis</i> , 2021, 32, 406-409.	0.1	0
5	Predictors of severe sepsis-related in-hospital mortality based on a multicenter cohort study. <i>Medicine (United States)</i> , 2021, 100, e24844.	0.4	1
6	Coagulopathy Induced by Veno-Arterial Extracorporeal Membrane Oxygenation Is Associated With a Poor Outcome in Patients With Out-of-Hospital Cardiac Arrest. <i>Frontiers in Medicine</i> , 2021, 8, 651832.	1.2	5
7	Hyperoxemia during resuscitation of trauma patients and increased intensive care unit length of stay: inverse probability of treatment weighting analysis. <i>World Journal of Emergency Surgery</i> , 2021, 16, 19.	2.1	8
8	Intensive care unit model and in-hospital mortality among patients with severe sepsis and septic shock. <i>Medicine (United States)</i> , 2021, 100, e26132.	0.4	1
9	Incidence and Impact of Dysglycemia in Patients with Sepsis Under Moderate Glycemic Control. <i>Shock</i> , 2021, 56, 507-513.	1.0	4
10	Disseminated intravascular coagulation immediately after trauma predicts a poor prognosis in severely injured patients. <i>Scientific Reports</i> , 2021, 11, 11031.	1.6	13
11	Thromboplasmininflammation in COVID-19 Coagulopathy: Three Viewpoints for Diagnostic and Therapeutic Strategies. <i>Frontiers in Immunology</i> , 2021, 12, 649122.	2.2	34
12	Pathophysiology of Coagulopathy Induced by Traumatic Brain Injury Is Identical to That of Disseminated Intravascular Coagulation With Hyperfibrinolysis. <i>Frontiers in Medicine</i> , 2021, 8, 767637.	1.2	16
13	Newly Proposed Sepsis-Induced Coagulopathy Precedes International Society on Thrombosis and Haemostasis Overt-Disseminated Intravascular Coagulation and Predicts High Mortality. <i>Journal of Intensive Care Medicine</i> , 2020, 35, 643-649.	1.3	60
14	Complementary Role of Hypothermia Identification to the Quick Sequential Organ Failure Assessment Score in Predicting Patients With Sepsis at High Risk of Mortality: A Retrospective Analysis From a Multicenter, Observational Study. <i>Journal of Intensive Care Medicine</i> , 2020, 35, 502-510.	1.3	12
15	Demographics, Treatments, and Outcomes of Acute Respiratory Distress Syndrome: the Focused Outcomes Research in Emergency Care in Acute Respiratory Distress Syndrome, Sepsis, and Trauma (FORECAST) Study. <i>Shock</i> , 2020, 53, 544-549.	1.0	13
16	Identifying Septic Shock Populations Benefitting From Polymyxin B Hemoperfusion: A Prospective Cohort Study Incorporating a Restricted Cubic Spline Regression Model. <i>Shock</i> , 2020, 54, 667-674.	1.0	7
17	Characteristics and outcomes of frail patients with suspected infection in intensive care units: a descriptive analysis from a multicenter cohort study. <i>BMC Geriatrics</i> , 2020, 20, 485.	1.1	3
18	Underlying disorders of disseminated intravascular coagulation: Communication from the ISTH SSC Subcommittees on Disseminated Intravascular Coagulation and Perioperative and Critical Care Thrombosis and Hemostasis. <i>Journal of Thrombosis and Haemostasis</i> , 2020, 18, 2400-2407.	1.9	16

#	ARTICLE	IF	CITATIONS
19	A multicenter prospective validation study on disseminated intravascular coagulation in trauma-induced coagulopathy. <i>Journal of Thrombosis and Haemostasis</i> , 2020, 18, 2232-2244.	1.9	22
20	The SIRS criteria have better performance for predicting infection than qSOFA scores in the emergency department. <i>Scientific Reports</i> , 2020, 10, 8095.	1.6	29
21	History of diabetes may delay antibiotic administration in patients with severe sepsis presenting to emergency departments. <i>Medicine (United States)</i> , 2020, 99, e19446.	0.4	0
22	ISTH interim guidance on recognition and management of coagulopathy in COVID-19. <i>Journal of Thrombosis and Haemostasis</i> , 2020, 18, 1023-1026.	1.9	1,513
23	Impact of blood glucose abnormalities on outcomes and disease severity in patients with severe sepsis: An analysis from a multicenter, prospective survey of severe sepsis. <i>PLoS ONE</i> , 2020, 15, e0229919.	1.1	28
24	Defining trauma-induced coagulopathy with respect to future implications for patient management: Communication from the SSC of the ISTH. <i>Journal of Thrombosis and Haemostasis</i> , 2020, 18, 740-747.	1.9	56
25	Significance of body temperature in elderly patients with sepsis. <i>Critical Care</i> , 2020, 24, 387.	2.5	37
26	Characteristics and outcomes of bacteremia among ICU-admitted patients with severe sepsis. <i>Scientific Reports</i> , 2020, 10, 2983.	1.6	21
27	Risk modifiers of acute respiratory distress syndrome in patients with non-pulmonary sepsis: a retrospective analysis of the FORECAST study. <i>Journal of Intensive Care</i> , 2020, 8, 7.	1.3	11
28	The significance of disseminated intravascular coagulation on multiple organ dysfunction during the early stage of acute respiratory distress syndrome. <i>Thrombosis Research</i> , 2020, 191, 15-21.	0.8	24
29	Type and dose of heparin in Covid-19: Reply. <i>Journal of Thrombosis and Haemostasis</i> , 2020, 18, 2063-2064.	1.9	19
30	DOACs and newer hemophilia therapies in COVID-19: Reply. <i>Journal of Thrombosis and Haemostasis</i> , 2020, 18, 1795-1796.	1.9	17
31	Laboratory haemostasis monitoring in COVID-19. <i>Journal of Thrombosis and Haemostasis</i> , 2020, 18, 2058-2060.	1.9	25
32	Identifying Sepsis Populations Benefitting from Anticoagulant Therapy: A Prospective Cohort Study Incorporating a Restricted Cubic Spline Regression Model. <i>Thrombosis and Haemostasis</i> , 2019, 119, 1740-1751.	1.8	21
33	Clinical features of patients with candidemia in sepsis. <i>Journal of General and Family Medicine</i> , 2019, 20, 161-163.	0.3	1
34	Significance of plasma fibrinogen level and antithrombin activity in sepsis: A multicenter cohort study using a cubic spline model. <i>Thrombosis Research</i> , 2019, 181, 17-23.	0.8	47
35	Trauma-induced coagulopathy: The past, present, and future: A comment. <i>Journal of Thrombosis and Haemostasis</i> , 2019, 17, 1567-1569.	1.9	3
36	Trends in sepsis care in Japan: comparison of two sepsis cohort studies conducted by the Japanese Association for Acute Medicine. <i>Acute Medicine & Surgery</i> , 2019, 6, 425-427.	0.5	0

#	ARTICLE	IF	CITATIONS
37	In-hospital mortality associated with the misdiagnosis or unidentified site of infection at admission. <i>Critical Care</i> , 2019, 23, 202.	2.5	28
38	Prognostic Accuracy of Quick SOFA is different according to the severity of illness in infectious patients. <i>Journal of Infection and Chemotherapy</i> , 2019, 25, 943-949.	0.8	5
39	Effect of a Recombinant Human Soluble Thrombomodulin on Mortality in Patients With Sepsis-Associated Coagulopathy. <i>JAMA - Journal of the American Medical Association</i> , 2019, 321, 1993.	3.8	221
40	Disseminated intravascular coagulation in cardiac arrest and resuscitation. <i>Journal of Thrombosis and Haemostasis</i> , 2019, 17, 1205-1216.	1.9	33
41	Variations in infection sites and mortality rates among patients in intensive care units with severe sepsis and septic shock in Japan. <i>Journal of Intensive Care</i> , 2019, 7, 28.	1.3	44
42	Nighttime and non-business days are not associated with increased risk of in-hospital mortality in patients with severe sepsis in intensive care units in Japan: The JAAM FORECAST study. <i>Journal of Critical Care</i> , 2019, 52, 97-102.	1.0	9
43	Role of disseminated intravascular coagulation in severe sepsis. <i>Thrombosis Research</i> , 2019, 178, 182-188.	0.8	72
44	Implementation of earlier antibiotic administration in patients with severe sepsis and septic shock in Japan: a descriptive analysis of a prospective observational study. <i>Critical Care</i> , 2019, 23, 360.	2.5	35
45	Impact of Body Temperature Abnormalities on the Implementation of Sepsis Bundles and Outcomes in Patients With Severe Sepsis: A Retrospective Sub-Analysis of the Focused Outcome Research on Emergency Care for Acute Respiratory Distress Syndrome, Sepsis and Trauma Study. <i>Critical Care Medicine</i> , 2019, 47, 691-699.	0.4	40
46	A case of pneumococcal pneumonia with purulent pericarditis and thoracic empyema. <i>Journal of the Japanese Society of Intensive Care Medicine</i> , 2019, 26, 189-190.	0.0	0
47	The roles of activated protein C in experimental trauma models. <i>Chinese Journal of Traumatology - English Edition</i> , 2018, 21, 311-315.	0.7	5
48	Characteristics, management, and in-hospital mortality among patients with severe sepsis in intensive care units in Japan: the FORECAST study. <i>Critical Care</i> , 2018, 22, 322.	2.5	89
49	Activated protein C plays no major roles in the inhibition of coagulation or increased fibrinolysis in acute coagulopathy of trauma-shock: a systematic review. <i>Thrombosis Journal</i> , 2018, 16, 13.	0.9	24
50	The case of extracorporeal membrane oxygenation (ECMO) catheter misplacement in the ascending lumbar vein. <i>Journal of the Japanese Society of Intensive Care Medicine</i> , 2018, 25, 145-146.	0.0	1
51	Efficacy and Bleeding Risk of Antithrombin Supplementation in Patients With Septic Disseminated Intravascular Coagulation: A Third Survey. <i>Clinical and Applied Thrombosis/Hemostasis</i> , 2017, 23, 422-428.	0.7	14
52	Should all patients with sepsis receive anticoagulation? Yes. <i>Intensive Care Medicine</i> , 2017, 43, 452-454.	3.9	21
53	Differences in coagulofibrinolytic changes between post-cardiac arrest syndrome of cardiac causes and hypoxic insults: a pilot study. <i>Acute Medicine & Surgery</i> , 2017, 4, 371-372.	0.5	7
54	Fibrin/fibrinogen degradation products (FDP) at hospital admission predict neurological outcomes in out-of-hospital cardiac arrest patients. <i>Resuscitation</i> , 2017, 111, 62-67.	1.3	24

#	ARTICLE	IF	CITATIONS
55	Assessment of mortality by qSOFA in patients with sepsis outside ICU: A post hoc subgroup analysis by the Japanese Association for Acute Medicine Sepsis Registry Study Group. <i>Journal of Infection and Chemotherapy</i> , 2017, 23, 757-762.	0.8	20
56	The qSOFA requires validation as a promptly applicable clinical criterion. <i>Acute Medicine & Surgery</i> , 2017, 4, 225-226.	0.5	0
57	Early evaluation of severity in patients with severe sepsis: a comparison with "septic shock" subgroup analysis of the Japanese Association for Acute Medicine Sepsis Registry (<sc>JAAM</sc> <sc>SR</sc>). <i>Acute Medicine & Surgery</i> , 2017, 4, 426-431.	0.5	5
58	Effects of combination therapy using antithrombin and thrombomodulin for sepsis-associated disseminated intravascular coagulation. <i>Annals of Intensive Care</i> , 2017, 7, 110.	2.2	22
59	Disseminated intravascular coagulation with increased fibrinolysis during the early phase of isolated traumatic brain injury. <i>Critical Care</i> , 2017, 21, 219.	2.5	37
60	Pharmacokinetics of recombinant human soluble thrombomodulin in disseminated intravascular coagulation patients with acute renal dysfunction. <i>Thrombosis and Haemostasis</i> , 2017, 117, 851-859.	1.8	19
61	Recent advances in diagnosis and treatment of disseminated intravascular coagulation in sepsis and trauma. <i>Japanese Journal of Thrombosis and Hemostasis</i> , 2017, 28, 492-501.	0.1	0
62	HMGB1 Promotes Intraoral Palatal Wound Healing through RAGE-Dependent Mechanisms. <i>International Journal of Molecular Sciences</i> , 2016, 17, 1961.	1.8	26
63	Infection site is predictive of outcome in acute lung injury associated with severe sepsis and septic shock. <i>Respirology</i> , 2016, 21, 898-904.	1.3	37
64	The response time threshold for predicting favourable neurological outcomes in patients with bystander-witnessed out-of-hospital cardiac arrest. <i>Resuscitation</i> , 2016, 107, 65-70.	1.3	34
65	Disseminated intravascular coagulation with the fibrinolytic phenotype predicts the outcome of patients with out-of-hospital cardiac arrest. <i>Thrombosis Journal</i> , 2016, 14, 43.	0.9	26
66	Antithrombin supplementation and risk of bleeding in patients with sepsis-associated disseminated intravascular coagulation. <i>Thrombosis Research</i> , 2016, 145, 46-50.	0.8	16
67	Disseminated intravascular coagulation. <i>Nature Reviews Disease Primers</i> , 2016, 2, 16037.	18.1	367
68	Activated protein C does not increase in the early phase of trauma with disseminated intravascular coagulation: comparison with acute coagulopathy of trauma-shock. <i>Journal of Intensive Care</i> , 2016, 4, 1.	1.3	28
69	Pathophysiology of Trauma-Induced Coagulopathy and Management of Critical Bleeding Requiring Massive Transfusion. <i>Seminars in Thrombosis and Hemostasis</i> , 2016, 42, 155-165.	1.5	64
70	What's new in the diagnostic criteria of disseminated intravascular coagulation?. <i>Intensive Care Medicine</i> , 2016, 42, 1062-1064.	3.9	24
71	Impact of serum glucose levels on disease severity and outcome in patients with severe sepsis: an analysis from a multicenter, prospective survey of severe sepsis. <i>Acute Medicine & Surgery</i> , 2015, 2, 21-28.	0.5	5
72	Effects of prehospital epinephrine administration on neurological outcomes in patients with out-of-hospital cardiac arrest. <i>Journal of Intensive Care</i> , 2015, 3, 29.	1.3	16

#	ARTICLE	IF	CITATIONS
73	Pharmacokinetics and the optimal regimen for levofloxacin in critically ill patients receiving continuous hemodiafiltration. <i>Journal of Intensive Care</i> , 2015, 3, 22.	1.3	3
74	Local hemostasis, immunothrombosis, and systemic disseminated intravascular coagulation in trauma and traumatic shock. <i>Critical Care</i> , 2015, 19, 72.	2.5	100
75	Rapid Evaluation of Fibrinogen Levels Using the CG02N Whole Blood Coagulation Analyzer. <i>Seminars in Thrombosis and Hemostasis</i> , 2015, 41, 267-271.	1.5	21
76	Hemostasis and Thrombosis in Trauma Patients. <i>Seminars in Thrombosis and Hemostasis</i> , 2015, 41, 026-034.	1.5	24
77	Fibrinogen Level Deteriorates before Other Routine Coagulation Parameters and Massive Transfusion in the Early Phase of Severe Trauma: A Retrospective Observational Study. <i>Seminars in Thrombosis and Hemostasis</i> , 2015, 41, 035-042.	1.5	62
78	The usefulness of antithrombin activity monitoring during antithrombin supplementation in patients with sepsis-associated disseminated intravascular coagulation. <i>Thrombosis Research</i> , 2015, 135, 897-901.	0.8	27
79	Noble-Collip Drum Trauma Induces Disseminated Intravascular Coagulation But Not Acute Coagulopathy of Trauma-Shock. <i>Shock</i> , 2015, 43, 261-267.	1.0	18
80	Should laryngeal tubes or masks be used for out-of-hospital cardiac arrest patients?. <i>American Journal of Emergency Medicine</i> , 2015, 33, 1360-1363.	0.7	13
81	Two cases of hyperthermia induced by zonisamide. <i>Journal of the Japanese Society of Intensive Care Medicine</i> , 2015, 22, 519-522.	0.0	1
82	A multicenter, prospective evaluation of quality of care and mortality in Japan based on the Surviving Sepsis Campaign guidelines. <i>Journal of Infection and Chemotherapy</i> , 2014, 20, 115-120.	0.8	37
83	Effects of Rikkunshito (traditional Japanese medicine) on enteral feeding and the plasma ghrelin level in critically ill patients: a pilot study. <i>Journal of Intensive Care</i> , 2014, 2, 53.	1.3	10
84	Post-marketing surveillance data of thrombomodulin alfa: sub-analysis in patients with sepsis-induced disseminated intravascular coagulation. <i>Journal of Intensive Care</i> , 2014, 2, 30.	1.3	29
85	Epidemiology of severe sepsis in Japanese intensive care units: A prospective multicenter study. <i>Journal of Infection and Chemotherapy</i> , 2014, 20, 157-162.	0.8	88
86	Effects of protease activated receptor (PAR)2 blocking peptide on endothelin-1 levels in kidney tissues in endotoxemic rat mode. <i>Life Sciences</i> , 2014, 102, 127-133.	2.0	13
87	Hemostasis during the early stages of trauma: comparison with disseminated intravascular coagulation. <i>Critical Care</i> , 2014, 18, R61.	2.5	48
88	The role of angiogenic factors and their soluble receptors in acute lung injury (ALI)/ acute respiratory distress syndrome (ARDS) associated with critical illness. <i>Journal of Inflammation</i> , 2013, 10, 6.	1.5	47
89	Coagulofibrinolytic changes in patients with disseminated intravascular coagulation associated with post-cardiac arrest syndrome—Fibrinolytic shutdown and insufficient activation of fibrinolysis lead to organ dysfunction. <i>Thrombosis Research</i> , 2013, 132, e64-e69.	0.8	45
90	Journal of Intensive Care: a new journal for all intensive care physicians. <i>Journal of Intensive Care</i> , 2013, 1, 1.	1.3	0

#	ARTICLE	IF	CITATIONS
91	A randomized, controlled, multicenter trial of the effects of antithrombin on disseminated intravascular coagulation in patients with sepsis. <i>Critical Care</i> , 2013, 17, R297.	2.5	132
92	A multicenter, prospective validation study of the Japanese Association for Acute Medicine disseminated intravascular coagulation scoring system in patients with severe sepsis. <i>Critical Care</i> , 2013, 17, R111.	2.5	156
93	The Dynamics of Angiogenic Factors and Their Soluble Receptors in Relation to Organ Dysfunction in Disseminated Intravascular Coagulation Associated with Sepsis. <i>Inflammation</i> , 2013, 36, 186-196.	1.7	18
94	Normal prothrombinase activity, increased systemic thrombin activity, and lower antithrombin levels in patients with disseminated intravascular coagulation at an early phase of trauma: Comparison with acute coagulopathy of trauma-shock. <i>Surgery</i> , 2013, 154, 48-57.	1.0	54
95	Effectiveness of end-expiratory lung volume measurements during the lung recruitment maneuver for patients with atelectasis. <i>Journal of Critical Care</i> , 2013, 28, 534.e1-534.e5.	1.0	1
96	Differentiating disseminated intravascular coagulation (DIC) with the fibrinolytic phenotype from coagulopathy of trauma and acute coagulopathy of trauma-shock (COT/ACOTS). <i>Journal of Thrombosis and Haemostasis</i> , 2013, 11, 826-835.	1.9	110
97	Role of Fibrinolysis in Sepsis. <i>Seminars in Thrombosis and Hemostasis</i> , 2013, 39, 392-399.	1.5	81
98	Effects of epinephrine administration in out-of-hospital cardiac arrest based on a propensity analysis. <i>Journal of Intensive Care</i> , 2013, 1, 12.	1.3	5
99	Massive Amounts of Tissue Factor Induce Fibrinolysis Without Tissue Hypoperfusion in Rats. <i>Shock</i> , 2013, 39, 514-519.	1.0	31
100	A Randomized, Controlled, Multicenter Trial of the Effects of Antithrombin on Disseminated Intravascular Coagulation in Patients With Sepsis. <i>Chest</i> , 2013, 144, 418A.	0.4	0
101	227. <i>Critical Care Medicine</i> , 2013, 41, A51.	0.4	0
102	The impact of body temperature abnormalities on the disease severity and outcome in patients with severe sepsis: an analysis from a multicenter, prospective survey of severe sepsis. <i>Critical Care</i> , 2013, 17, R271.	2.5	139
103	Disseminated Intravascular Coagulation (DIC) at an Early Phase of Trauma Continuously Proceeds to DIC at a Late Phase of Trauma. <i>Clinical and Applied Thrombosis/Hemostasis</i> , 2012, 18, 364-369.	0.7	11
104	Pharmacokinetics and Pharmacodynamics of Recombinant Soluble Thrombomodulin in Disseminated Intravascular Coagulation Patients With Renal Impairment. <i>Shock</i> , 2012, 37, 569-573.	1.0	10
105	The Utility of a Diagnostic Scoring System for Disseminated Intravascular Coagulation. <i>Critical Care Clinics</i> , 2012, 28, 373-388.	1.0	17
106	Synbiotic Therapy Reduces the Pathological Gram-Negative Rods Caused by an Increased Acetic Acid Concentration in the Gut. <i>Digestive Diseases and Sciences</i> , 2012, 57, 2642-2649.	1.1	17
107	Using angiogenic factors and their soluble receptors to predict organ dysfunction in patients with disseminated intravascular coagulation associated with severe trauma. <i>Critical Care</i> , 2012, 16, R63.	2.5	12
108	Angiogenic factors and their soluble receptors predict organ dysfunction and mortality in post-cardiac arrest syndrome. <i>Critical Care</i> , 2012, 16, R171.	2.5	29

#	ARTICLE	IF	CITATIONS
109	C0249 Elastase-mediated fibrinolysis in disseminated intravascular coagulation (DIC) associated with sepsis. <i>Thrombosis Research</i> , 2012, 130, S184.	0.8	0
110	The Siset incorrectly cited the JAAM DIC scoring system. <i>Thrombosis Research</i> , 2012, 129, 660.	0.8	1
111	A low TAFI activity and insufficient activation of fibrinolysis by both plasmin and neutrophil elastase promote organ dysfunction in disseminated intravascular coagulation associated with sepsis. <i>Thrombosis Research</i> , 2012, 130, 906-913.	0.8	27
112	Time-Dependent Alterations of VEGF and Its Signaling Molecules in Acute Lung Injury in a Rat Model of Sepsis. <i>Inflammation</i> , 2012, 35, 484-500.	1.7	36
113	The importance of early treatment with tranexamic acid in bleeding trauma patients: an exploratory analysis of the CRASH-2 randomised controlled trial. <i>Lancet, The</i> , 2011, 377, 1096-1101.e2.	6.3	950
114	Time-dependent expression of endothelin-1 in lungs and the effects of TNF- α blocking peptide on acute lung injury in an endotoxemic rat model. <i>Biomedical Research</i> , 2011, 32, 9-17.	0.3	21
115	Improved Detection of Heat Stroke-Induced Brain Injury by High B-Value Diffusion-Weighted Imaging. <i>Journal of Computer Assisted Tomography</i> , 2011, 35, 498-500.	0.5	10
116	Trauma, Shock, and Disseminated Intravascular Coagulation. <i>Annals of Surgery</i> , 2011, 254, 10-19.	2.1	149
117	Disseminated intravascular coagulation at an early phase of trauma is associated with consumption coagulopathy and excessive fibrinolysis both by plasmin and neutrophil elastase. <i>Surgery</i> , 2011, 149, 221-230.	1.0	96
118	Dramatic Changes of the Gut Flora Immediately After Severe and Sudden Insults. <i>Digestive Diseases and Sciences</i> , 2011, 56, 2361-2365.	1.1	138
119	Imbalance Between Macrophage Migration Inhibitory Factor and Cortisol Induces Multiple Organ Dysfunction in Patients with Blunt Trauma. <i>Inflammation</i> , 2011, 34, 193-197.	1.7	6
120	A case of severe multiple injuries due to a traffic accident with an intracerebral hemorrhage possibly caused by the rupture of an angiographically occult arteriovenous malformation. <i>Nihon Kyukyū Igakukai Zasshi</i> , 2011, 22, 62-69.	0.0	0
121	Microvascular thrombosis and multiple organ dysfunction syndrome. <i>Critical Care Medicine</i> , 2010, 38, S35-S42.	0.4	277
122	Invasive group A streptococcal infection in pregnancy. <i>Journal of Infection</i> , 2010, 60, 417-424.	1.7	23
123	Modified non-overly DIC diagnostic criteria predict the early phase of overt DIC. <i>American Journal of Hematology</i> , 2010, 85, 691-694.	2.0	35
124	SIVELESTAT (SELECTIVE NEUTROPHIL ELASTASE INHIBITOR) IMPROVES THE MORTALITY RATE OF SEPSIS ASSOCIATED WITH BOTH ACUTE RESPIRATORY DISTRESS SYNDROME AND DISSEMINATED INTRAVASCULAR COAGULATION PATIENTS. <i>Shock</i> , 2010, 33, 14-18.	1.0	74
125	IL-6 and IFN- γ from dsRNA-stimulated dendritic cells control expansion of regulatory T cells. <i>Biochemical and Biophysical Research Communications</i> , 2010, 391, 1421-1426.	1.0	10
126	Expert consensus for the treatment of disseminated intravascular coagulation in Japan. <i>Thrombosis Research</i> , 2010, 125, 6-11.	0.8	222

#	ARTICLE	IF	CITATIONS
127	Frequency and hemostatic abnormalities in pre-DIC patients. <i>Thrombosis Research</i> , 2010, 126, 74-78.	0.8	26
128	Coagulation and fibrinolytic responses at an early phase of trauma: The main issues in the world are reviewed and discussed. <i>Nihon Kyukyu Igakukai Zasshi</i> , 2010, 21, 765-778.	0.0	1
129	The relationship between high level of FDP (fibrin/fibrinogen degradation products) on the admission and massive bleeding in patients with blunt trauma. <i>Nihon Kyukyu Igakukai Zasshi</i> , 2010, 21, 165-171.	0.0	1
130	Effects of Antithrombin III in Patients With Disseminated Intravascular Coagulation Diagnosed by Newly Developed Diagnostic Criteria for Critical Illness. <i>Clinical and Applied Thrombosis/Hemostasis</i> , 2009, 15, 561-566.	0.7	17
131	Disseminated intravascular coagulation (DIC) diagnosed based on the Japanese Association for Acute Medicine criteria is a dependent continuum to overt DIC in patients with sepsis. <i>Thrombosis Research</i> , 2009, 123, 715-718.	0.8	70
132	The expression of 4 protease-activated receptors is associated with increased levels of TNF- α , tissue factor, and fibrin in the frontal cortex of endotoxemic rats. <i>Thrombosis Research</i> , 2009, 124, 498-501.	0.8	2
133	Disseminated intravascular coagulation with a fibrinolytic phenotype at an early phase of trauma predicts mortality. <i>Thrombosis Research</i> , 2009, 124, 608-613.	0.8	163
134	Application of the Japanese Association for Acute Medicine disseminated intravascular coagulation diagnostic criteria for patients at an early phase of trauma. <i>Thrombosis Research</i> , 2009, 124, 706-710.	0.8	42
135	Shortening of cardiopulmonary resuscitation time before the defibrillation worsens the outcome in out-of-hospital VF patients. <i>American Journal of Emergency Medicine</i> , 2009, 27, 470-474.	0.7	539
136	Acute Coagulopathy of Trauma Shock and Coagulopathy of Trauma: A Rebuttal. You Are Now Going Down the Wrong Path. <i>Journal of Trauma</i> , 2009, 67, 381-383.	2.3	57
137	The Administration of Ciprofloxacin During Continuous Renal Replacement Therapy: Pilot Study. <i>ASAIO Journal</i> , 2009, 55, 243-245.	0.9	9
138	PROTEASE-ACTIVATED RECEPTOR 2 BLOCKING PEPTIDE COUNTERACTS ENDOTOXIN-INDUCED INFLAMMATION AND COAGULATION AND AMELIORATES RENAL FIBRIN DEPOSITION IN A RAT MODEL OF ACUTE RENAL FAILURE. <i>Shock</i> , 2009, 32, 626-632.	1.0	19
139	Clinical course and outcome of disseminated intravascular coagulation diagnosed by Japanese Association for Acute Medicine criteria. <i>Thrombosis and Haemostasis</i> , 2008, 100, 1099-1105.	1.8	51
140	Natural history of disseminated intravascular coagulation diagnosed based on the newly established diagnostic criteria for critically ill patients: Results of a multicenter, prospective survey*. <i>Critical Care Medicine</i> , 2008, 36, 145-150.	0.4	205
141	THE RESPONSE OF ANTITHROMBIN III ACTIVITY AFTER SUPPLEMENTATION DECREASES IN PROPORTION TO THE SEVERITY OF SEPSIS AND LIVER DYSFUNCTION. <i>Shock</i> , 2008, 30, 649-652.	1.0	16
142	INSUFFICIENT PRODUCTION OF URINARY TRYPSIN INHIBITOR FOR NEUTROPHIL ELASTASE RELEASE AFTER CARDIAC ARREST. <i>Shock</i> , 2008, 29, 549-552.	1.0	6
143	A Prospective Comparative Study of Three Sets of Criteria for Disseminated Intravascular Coagulation: ISTH Criteria vs Japanese Criteria. <i>Clinical and Applied Thrombosis/Hemostasis</i> , 2007, 13, 65-72.	0.7	10
144	SIRS-ASSOCIATED COAGULOPATHY AND ORGAN DYSFUNCTION IN CRITICALLY ILL PATIENTS WITH THROMBOCYTOPENIA. <i>Shock</i> , 2007, 28, 411-417.	1.0	63

#	ARTICLE	IF	CITATIONS
145	Predicting the Severity of Systemic Inflammatory Response Syndrome (SIRS)-Associated Coagulopathy With Hemostatic Molecular Markers and Vascular Endothelial Injury Markers. <i>Journal of Trauma</i> , 2007, 63, 1093-1098.	2.3	43
146	An alternative pathway for fibrinolysis is activated in patients who have undergone cardiopulmonary bypass surgery and major abdominal surgery. <i>Thrombosis Research</i> , 2007, 120, 87-93.	0.8	13
147	The activation of neutrophil elastase-mediated fibrinolysis is not sufficient to overcome the fibrinolytic shutdown of disseminated intravascular coagulation associated with systemic inflammation. <i>Thrombosis Research</i> , 2007, 121, 67-73.	0.8	19
148	A Prospective Comparison of New Japanese Criteria for Disseminated Intravascular Coagulation. <i>Clinical and Applied Thrombosis/Hemostasis</i> , 2007, 13, 172-181.	0.7	17
149	Differential Expression, Time Course and Distribution of Four PARs in Rats with Endotoxin-induced Acute Lung Injury. <i>Inflammation</i> , 2007, 30, 14-27.	1.7	25
150	High Macrophage Migration Inhibitory Factor Levels in Disseminated Intravascular Coagulation Patients with Systemic Inflammation. <i>Inflammation</i> , 2007, 30, 118-124.	1.7	24
151	Chronological expression of Endothelin-1 and TNF- α in Acute Liver Injury and its amelioration by PAR2 Blockade in a septic Rat Model. <i>FASEB Journal</i> , 2007, 21, .	0.2	0
152	Non-Operative Management of a Blunt Traumatic Intraperitoneal Bladder Rupture as Damage Control after a Severe Pelvic Fracture. <i>Nihon Kyukyu Igakukai Zasshi</i> , 2007, 18, 23-26.	0.0	2
153	Time-dependent expression of renal vaso-regulatory molecules in LPS-induced endotoxemia in rat. <i>Peptides</i> , 2006, 27, 2258-2270.	1.2	49
154	Evaluation of haemostatic molecular markers for diagnosis of disseminated intravascular coagulation in patients with infections. <i>Thrombosis and Haemostasis</i> , 2006, 95, 282-287.	1.8	40
155	Chronological expression of PAR isoforms in acute liver injury and its amelioration by PAR2 blockade in a rat model of sepsis. <i>Thrombosis and Haemostasis</i> , 2006, 96, 830-838.	1.8	14
156	Pharmacokinetics and the Most Suitable Regimen of Panipenem/Beta Mipron in Critically Ill Patients Receiving Continuous Renal Replacement Therapy: A Pilot Study. <i>ASAIO Journal</i> , 2006, 52, 398-403.	0.9	10
157	A multicenter, prospective validation of disseminated intravascular coagulation diagnostic criteria for critically ill patients: Comparing current criteria*. <i>Critical Care Medicine</i> , 2006, 34, 625-631.	0.4	512
158	RESULTS OF TUBERCULIN SKIN TEST IN 1,172 CASES OF BACTIVITY AFTER SUPPLEMENTATION HAVE A PREDICTIVE VALUE IN CRITICALLY ILL PATIENTS. <i>Chest</i> , 2006, 130, 206S.	0.4	0
159	First day dynamic changes in antithrombin III activity after supplementation have a predictive value in critically ill patients. <i>American Journal of Hematology</i> , 2006, 81, 907-914.	2.0	19
160	Tissue Factor in Trauma and Organ Dysfunction. <i>Seminars in Thrombosis and Hemostasis</i> , 2006, 32, 048-053.	1.5	30
161	Altered expression of endothelin, vascular endothelial growth factor, and its receptor in hepatic tissue in endotoxemic rat. <i>Experimental Biology and Medicine</i> , 2006, 231, 1182-6.	1.1	6
162	PREVENTION OF SEPSIS-INDUCED ACUTE LUNG INJURY BY SMALL INTERFERING RNA FOR INHIBITION OF TISSUE FACTOR EXPRESSION.. <i>Critical Care Medicine</i> , 2005, 33, A7.	0.4	0

#	ARTICLE	IF	CITATIONS
163	Evaluation of New Japanese Diagnostic Criteria for Disseminated Intravascular Coagulation in Critically Ill Patients. <i>Clinical and Applied Thrombosis/Hemostasis</i> , 2005, 11, 71-76.	0.7	77
164	Nuclear Factor- κ B Decoy Oligodeoxynucleotides Prevent Acute Lung Injury in Mice with Cecal Ligation and Puncture-Induced Sepsis. <i>Molecular Pharmacology</i> , 2005, 67, 1018-1025.	1.0	92
165	Serial changes in neutrophil endothelial activation markers during the course of sepsis associated with disseminated intravascular coagulation. <i>Thrombosis Research</i> , 2005, 116, 91-100.	0.8	52
166	Rhabdomyolysis due to water intoxication. <i>Journal of the Japanese Society of Intensive Care Medicine</i> , 2005, 12, 219-222.	0.0	0
167	HIGH MACROPHAGE MIGRATION INHIBITORY FACTOR LEVELS IN DISSEMINATED INTRAVASCULAR COAGULATION PATIENTS WITH SYSTEMIC INFLAMMATORY RESPONSE SYNDROME AND SEPSIS.. <i>Critical Care Medicine</i> , 2005, 33, A162.	0.4	0
168	SEPSIS-INDUCED CHANGES IN THE SIGNALING MECHANISMS FOR GLUCOSE TRANSPORT 4 TRANSLOCATION TO THE MEMBRANE.. <i>Critical Care Medicine</i> , 2005, 33, A132.	0.4	0
169	Temporal changes in pulmonary expression of key procoagulant molecules in rabbits with endotoxin-induced acute lung injury: elevated expression levels of protease-activated receptors. <i>Thrombosis and Haemostasis</i> , 2004, 92, 966-979.	1.8	38
170	Histamine H1 and H2 Receptor Gene and Protein Levels Are Differentially Expressed in the Hearts of Rodents and Humans. <i>Journal of Pharmacology and Experimental Therapeutics</i> , 2004, 309, 786-795.	1.3	67
171	Therapeutic effect of in vivo transfection of transcription factor decoy to NF- κ B on septic lung in mice. <i>American Journal of Physiology - Lung Cellular and Molecular Physiology</i> , 2004, 287, L1248-L1255.	1.3	59
172	Systemic Inflammation and Disseminated Intravascular Coagulation in Early Stage of ALI and ARDS: Role of Neutrophil and Endothelial Activation. <i>Inflammation</i> , 2004, 28, 237-244.	1.7	81
173	Early Diagnosis of Toxic Shock like Syndrome by Magnetic Resonance Imaging and Histopathology. <i>Nihon Kyukyu Igakukai Zasshi</i> , 2004, 15, 563-568.	0.0	2
174	Pharmacokinetics of teicoplanin in critically ill patients undergoing continuous hemodiafiltration. <i>Intensive Care Medicine</i> , 2003, 29, 2094-2095.	3.9	27
175	Pharmacokinetics and the most suitable dosing regimen of fluconazole in critically ill patients receiving continuous hemodiafiltration. <i>Intensive Care Medicine</i> , 2003, 29, 1844-1848.	3.9	54
176	Normalization by edaravone, a free radical scavenger, of irradiation-reduced endothelial nitric oxide synthase expression. <i>European Journal of Pharmacology</i> , 2003, 476, 131-137.	1.7	26
177	Imbalances between the levels of tissue factor and tissue factor pathway inhibitor in ARDS patients. <i>Thrombosis Research</i> , 2003, 109, 119-124.	0.8	94
178	Contractions to Histamine in Pulmonary and Mesenteric Arteries from Endotoxemic Rabbits: Modulation by Vascular Expressions of Inducible Nitric-Oxide Synthase and Histamine H1-Receptors. <i>Journal of Pharmacology and Experimental Therapeutics</i> , 2003, 307, 175-181.	1.3	22
179	The Sympathomimetic Actions of l-Ephedrine and d-Pseudoephedrine: Direct Receptor Activation or Norepinephrine Release?. <i>Anesthesia and Analgesia</i> , 2003, 97, 1239-1245.	1.1	59
180	Tissue factor production not balanced by tissue factor pathway inhibitor in sepsis promotes poor prognosis*. <i>Critical Care Medicine</i> , 2002, 30, 1729-1734.	0.4	96

#	ARTICLE	IF	CITATIONS
181	Combined Activation of Coagulation and Inflammation has an Important Role in Multiple Organ Dysfunction and Poor Outcome after Severe Trauma. <i>Thrombosis and Haemostasis</i> , 2002, 88, 943-949.	1.8	89
182	Hemodynamic significance of histamine synthesis and histamine H ₁ - and H ₂ -receptor gene expression during endotoxemia. <i>Naunyn-Schmiedeberg's Archives of Pharmacology</i> , 2002, 366, 513-521.	1.4	24
183	Abdominal compartment syndrome and intrahepatic portal venous gas: a possible complication of endoscopy. <i>Intensive Care Medicine</i> , 2002, 28, 1680-1681.	3.9	10
184	Ventricular fibrillation after pseudo-Bartter's syndrome. <i>Annals of Emergency Medicine</i> , 2002, 39, 205-206.	0.3	0
185	Combined activation of coagulation and inflammation has an important role in multiple organ dysfunction and poor outcome after severe trauma. <i>Thrombosis and Haemostasis</i> , 2002, 88, 943-9.	1.8	32
186	Another point of view on the mechanism of thrombin generation during cardiopulmonary bypass: Role of tissue factor pathway inhibitor. <i>Journal of Cardiothoracic and Vascular Anesthesia</i> , 2001, 15, 60-64.	0.6	42
187	Systematic Elucidation of Effects of Tranexamic Acid on Fibrinolysis and Bleeding During and After Cardiopulmonary Bypass Surgery. <i>Thrombosis Research</i> , 2001, 104, 301-307.	0.8	26
188	Tissue factor pathway inhibitor response does not correlate with tissue factor-induced disseminated intravascular coagulation and multiple organ dysfunction syndrome in trauma patients. <i>Critical Care Medicine</i> , 2001, 29, 262-266.	0.4	54
189	Serial Measurement of Arterial Lactate Concentrations as a Prognostic Indicator in Relation to the Incidence of Disseminated Intravascular Coagulation in Patients with Systemic Inflammatory Response Syndrome. <i>Surgery Today</i> , 2001, 31, 853-859.	0.7	23
190	Systemic Macrophage Migration Inhibitory Factor Release Following Hepatic Resection. <i>Surgery Today</i> , 2001, 31, 605-609.	0.7	8
191	Predictors of Mortality in Patients Treated with Continuous Hemodiafiltration for Acute Renal Failure in an Intensive Care Setting. <i>ASAIO Journal</i> , 2001, 47, 86-91.	0.9	34
192	Disseminated Intravascular Coagulation in Trauma Patients. <i>Seminars in Thrombosis and Hemostasis</i> , 2001, 27, 585-592.	1.5	153
193	The Effect of Calcium Channel Blockers on Cerebral Oxygenation During Tracheal Extubation. <i>Anesthesia and Analgesia</i> , 2000, 91, 347-352.	1.1	1
194	Pentobarbital inhibits apoptosis in neuronal cells. <i>Critical Care Medicine</i> , 2000, 28, 1899-1904.	0.4	20
195	Impairment of Cardiac β_2 -Adrenoceptor Cellular Signaling by Decreased Expression of Gs α in Septic Rabbits. <i>Anesthesiology</i> , 2000, 93, 1465-1473.	1.3	65
196	Phosphodiesterase III Inhibitor Olprinone Chlorate Is Not Significantly Removed by Continuous Venovenous Hemodiafiltration. <i>ASAIO Journal</i> , 2000, 46, 635.	0.9	2
197	Diminished function and expression of the cardiac Na ⁺ + Ca ²⁺ exchanger in diabetic rats: implication in Ca ²⁺ overload. <i>Journal of Physiology</i> , 2000, 527, 85-94.	1.3	125
198	An increase in macrophage migration inhibitory factor release in patients with cardiopulmonary bypass surgery. <i>Surgery Today</i> , 2000, 30, 689-694.	0.7	20

#	ARTICLE	IF	CITATIONS
199	Hypokalemic coma with tetany following pseudo-Bartter syndrome. <i>Intensive Care Medicine</i> , 2000, 26, 1711-1711.	3.9	1
200	Therapeutic strategy of perioperative use of percutaneous cardiopulmonary bypass support (PCPS) for adult cardiac surgery. <i>General Thoracic and Cardiovascular Surgery</i> , 1999, 47, 20-26.	0.4	8
201	Diabetes-induced down-regulation of I^{21} -adrenoceptor mRNA expression in rat heart. <i>Biochemical Pharmacology</i> , 1999, 58, 881-885.	2.0	52
202	Tissue Factor and Tissue Factor Pathway Inhibitor Levels during and after Cardiopulmonary Resuscitation. <i>Thrombosis Research</i> , 1999, 96, 107-113.	0.8	46
203	Disseminated Intravascular Coagulation and Sustained Systemic Inflammatory Response Syndrome Predict Organ Dysfunctions After Trauma. <i>Annals of Surgery</i> , 1999, 229, 121-127.	2.1	110
204	Coagulofibrinolytic Changes after Isolated Head Injury Are Not Different from Those in Trauma Patients without Head Injury. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 1999, 46, 1070-1076.	1.1	65
205	Systemic Activation of Tissue-Factor Dependent Coagulation Pathway in Evolving Acute Respiratory Distress Syndrome in Patients with Trauma and Sepsis. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 1999, 47, 719.	1.1	68
206	Significant Correlations between Tissue Factor and Thrombin Markers in Trauma and Septic Patients with Disseminated Intravascular Coagulation. <i>Thrombosis and Haemostasis</i> , 1998, 79, 1111-1115.	1.8	138
207	Pharmacokinetics and clearance of ganciclovir during continuous hemodiafiltration. <i>Critical Care Medicine</i> , 1998, 26, 184-187.	0.4	15
208	Activation of the extrinsic coagulation pathway in patients with severe sepsis and septic shock. <i>Critical Care Medicine</i> , 1998, 26, 2005-2009.	0.4	100
209	Protein C activation during cardiopulmonary resuscitation following out-of-hospital cardiac arrest. <i>Journal of Anesthesia</i> , 1997, 11, 239-241.	0.7	2
210	Massive Fibrin Formation with Consecutive Impairment of Fibrinolysis in Patients with Out-of-Hospital Cardiac Arrest. <i>Thrombosis and Haemostasis</i> , 1997, 77, 278-282.	1.8	118
211	Participation of tissue factor and thrombin in posttraumatic systemic inflammatory syndrome. <i>Critical Care Medicine</i> , 1997, 25, 1820-1826.	0.4	99
212	Increased Neutrophil Elastase, Persistent Intravascular Coagulation, and Decreased Fibrinolytic Activity in Patients with Posttraumatic Acute Respiratory Distress Syndrome. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 1997, 42, 1068-1072.	1.1	26
213	A case of severe colchicine intoxication progressing to multiple organ dysfunction syndrome: Case report. <i>Journal of the Japanese Society of Intensive Care Medicine</i> , 1997, 4, 215-219.	0.0	0
214	Serial Studies of Protein C in Trauma Patients. <i>Japanese Journal of Thrombosis and Hemostasis</i> , 1996, 7, 312-318.	0.1	5
215	Diminishment of Contractions Associated with Depolarization-Evoked Activation of Ca^{2+} Channels in Diabetic Rat Aorta. <i>Journal of Vascular Research</i> , 1996, 33, 454-462.	0.6	5
216	Disseminated Intravascular Coagulation Is a Frequent Complication of Systemic Inflammatory Response Syndrome. <i>Thrombosis and Haemostasis</i> , 1996, 75, 224-228.	1.8	169

#	ARTICLE	IF	CITATIONS
217	Fibrinolytic shutdown after cardiopulmonary bypass surgery is caused by circulating cytokines during operation, accompanied by endothelial injury. <i>Journal of Anesthesia</i> , 1995, 9, 17-21.	0.7	1
218	Cytokines, soluble thrombomodulin and disseminated intravascular coagulation in patients with systemic inflammatory response syndrome. <i>Thrombosis Research</i> , 1995, 80, 519-526.	0.8	50
219	Cytokines and plasminogen activator inhibitor-1 in posttrauma disseminated intravascular coagulation. <i>Critical Care Medicine</i> , 1995, 23, 1835-1842.	0.4	171
220	Soluble Thrombomodulin Increases in Patients with Disseminated Intravascular Coagulation and in Those with Multiple Organ Dysfunction Syndrome after Trauma. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 1995, 39, 660-664.	1.1	53
221	Two Cases of Thrombotic Thrombocytopenic Purpura Secondary to Sepsis. <i>Journal of the Japanese Society of Intensive Care Medicine</i> , 1995, 2, 53-53.	0.0	0
222	Differential Diagnostic Value of Fibrinopeptides and FDP in Patients with Pulmonary Thromboembolism and Acute Myocardial Infarction. <i>Journal of the Japanese Society of Intensive Care Medicine</i> , 1995, 2, 81-85.	0.0	0
223	The effects of massive transfusion and haptoglobin therapy on hemolysis in trauma patients. <i>Surgery Today</i> , 1994, 24, 785-790.	0.7	27
224	Attenuated contractile response of diabetic rat aorta to caffeine but not to noradrenaline in Ca ²⁺ -free medium. <i>European Journal of Pharmacology</i> , 1994, 256, 215-219.	1.7	16
225	Diagnostic and prognostic value of fibrinopeptides in patients with clinically suspected pulmonary embolism. <i>Thrombosis Research</i> , 1994, 75, 195-202.	0.8	6
226	Histamine Receptors Mediating a Positive Inotropic Effect in Guinea Pig and Rabbit Ventricular Myocardium: Distribution of the Receptors and Their Possible Intracellular Coupling Processes. <i>The Japanese Journal of Pharmacology</i> , 1994, 65, 327-336.	1.2	13
227	Diagnostic value of plasmin antiplasmin complex(PAC) and D-dimer in patients with clinically suspected pulmonary thromboembolism.. <i>Nihon Kyukyu Igakukai Zasshi</i> , 1994, 5, 271-277.	0.0	0
228	Altered cardiac adrenergic neurotransmission in streptozotocin-induced diabetic rats. <i>British Journal of Pharmacology</i> , 1993, 109, 1276-1281.	2.7	21
229	An application of medical decision making to DIC diagnosis in trauma patients. Selection of new molecular markers and determining their optimal cut-off points.. <i>Nihon Kyukyu Igakukai Zasshi</i> , 1993, 4, 211-217.	0.0	0
230	Polymerase chain reaction for rapid detection of human cytomegalovirus. Clinical use in the ICU patient.. <i>Nihon Kyukyu Igakukai Zasshi</i> , 1993, 4, 1-6.	0.0	0
231	Posttrauma coagulation and fibrinolysis. <i>Critical Care Medicine</i> , 1992, 20, 594-600.	0.4	150
232	Methamphetamine intoxication with submassive hepatic cell necrosis.. <i>Nihon Kyukyu Igakukai Zasshi</i> , 1992, 3, 30-34.	0.0	0
233	Identification and characterization of histamine H ₁ - and H ₂ -receptors in guinea-pig left atrial membranes by [³ H]mepyramine and [³ H]tiotidine binding. <i>British Journal of Pharmacology</i> , 1991, 103, 1573-1579.	2.7	18
234	Characterization of histamine receptors modulating inotropic and biochemical activities in rabbit left atria. <i>European Journal of Pharmacology</i> , 1991, 196, 29-36.	1.7	20

#	ARTICLE	IF	CITATIONS
235	Massive Gastrointestinal Hemorrhage Caused by Cytomegalovirus Infection.. Nihon Kyukyu Igakukai Zasshi, 1991, 2, 949-954.	0.0	0
236	A comparison of serum ionized calcium in arterial and mixed venous blood during CPR. Annals of Emergency Medicine, 1990, 19, 850-856.	0.3	12
237	End-tidal carbon dioxide concentration during cardiopulmonary resuscitation(CPR). Is this effective as an indicator of CPR success?. Nihon Kyukyu Igakukai Zasshi, 1990, 1, 19-24.	0.0	0
238	Analysis of Blood Coagulation and Fibrinolysis During the Acute Period of Severe Subarachnoid Hemorrhage Using Newly Designed Molecular Markers.. Nihon Kyukyu Igakukai Zasshi, 1990, 1, 145-152.	0.0	0
239	Plateau Waves due to Carbon Dioxide Production in a Patient with Hypoxic Encephalopathy. Critical Care Medicine, 1989, 17, 1083.	0.4	2
240	Variation in serum ionized calcium on cardiopulmonary resuscitation. Journal of Anesthesia, 1988, 2, 154-160.	0.7	20
241	Association of Histones With Coagulofibrinolytic Responses and Organ Dysfunction in Adult Post-cardiac Arrest Syndrome. Frontiers in Cardiovascular Medicine, 0, 9, .	1.1	2