

Joshua T Anderson

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4090993/publications.pdf>

Version: 2024-02-01

40
papers

615
citations

623734

14
h-index

610901

24
g-index

40
all docs

40
docs citations

40
times ranked

657
citing authors

#	ARTICLE	IF	CITATIONS
1	Chronic Opioid Therapy After Lumbar Fusion Surgery for Degenerative Disc Disease in a Workersâ€™™ Compensation Setting. <i>Spine</i> , 2015, 40, 1775-1784.	2.0	92
2	Clinical Depression Is a Strong Predictor of Poor Lumbar Fusion Outcomes Among Workersâ€™¼ Compensation Subjects. <i>Spine</i> , 2015, 40, 748-756.	2.0	85
3	Preoperative Opioid Use is a Predictor of Poor Return to Work in Workersâ€™™ Compensation Patients After Lumbar Discectomy. <i>Spine</i> , 2018, 43, 594-602.	2.0	48
4	Prolonged Preoperative Opioid Therapy Associated With Poor Return to Work Rates After Single-Level Cervical Fusion for Radiculopathy for Patients Receiving Workersâ€™™ Compensation Benefits. <i>Spine</i> , 2017, 42, E104-E110.	2.0	46
5	Can Internet Information on Vertebroplasty be a Reliable Means of Patient Self-education?. <i>Clinical Orthopaedics and Related Research</i> , 2014, 472, 1597-1604.	1.5	38
6	Single-Level Lumbar Fusion for Degenerative Disc Disease Is Associated With Worse Outcomes Compared With Fusion for Spondylolisthesis in a Workersâ€™¼ Compensation Setting. <i>Spine</i> , 2015, 40, 323-331.	2.0	35
7	Return to Work After Diskogenic Fusion in Workers' Compensation Subjects. <i>Orthopedics</i> , 2015, 38, e1065-72.	1.1	28
8	Preoperative Opioid Use. <i>Clinical Spine Surgery</i> , 2018, 31, E19-E24.	1.3	26
9	Workers' Compensation, Return to Work, and Lumbar Fusion for Spondylolisthesis. <i>Orthopedics</i> , 2016, 39, e1-8.	1.1	24
10	Prolonged Preoperative Opioid Therapy in Patients With Degenerative Lumbar Stenosis in a Workersâ€™™ Compensation Setting. <i>Spine</i> , 2017, 42, E1140-E1146.	2.0	22
11	Treatment of Recurrent Lumbar Disc Herniation With or Without Fusion in Workersâ€™™ Compensation Subjects. <i>Spine</i> , 2017, 42, E864-E870.	2.0	21
12	Neck Pain, Preoperative Opioids, and Functionality After Cervical Fusion. <i>Orthopedics</i> , 2017, 40, 25-32.	1.1	21
13	Decompression Versus Decompression and Fusion for Degenerative Lumbar Stenosis in a Workersâ€™™ Compensation Setting. <i>Spine</i> , 2017, 42, 1017-1023.	2.0	20
14	Return to Work Rates After Single-level Cervical Fusion for Degenerative Disc Disease Compared With Fusion for Radiculopathy in a Workersâ€™™ Compensation Setting. <i>Spine</i> , 2016, 41, 1160-1166.	2.0	17
15	Surgical and Functional Outcomes After Multilevel Cervical Fusion for Degenerative Disc Disease Compared With Fusion for Radiculopathy. <i>Spine</i> , 2017, 42, 700-706.	2.0	12
16	A Multidisciplinary Transitional Pain Management Program Is Associated With Reduced Opioid Dependence After Primary Total Joint Arthroplasty. <i>Journal of Arthroplasty</i> , 2022, 37, 1048-1053.	3.1	11
17	Return to Work and Multilevel Versus Single-Level Cervical Fusion for Radiculopathy in a Workersâ€™™ Compensation Setting. <i>Spine</i> , 2017, 42, E111-E116.	2.0	9
18	The Timing of Surgery Affects Return to Work Rates in Patients With Degenerative Lumbar Stenosis in a Workersâ€™™ Compensation Setting. <i>Clinical Spine Surgery</i> , 2017, 30, E1444-E1449.	1.3	8

#	ARTICLE	IF	CITATIONS
19	Analysis of Internet information on the controversial X-Stop device. Spine Journal, 2014, 14, 2412-2419.	1.3	7
20	NKAP Regulates Senescence and Cell Death Pathways in Hematopoietic Progenitors. Frontiers in Cell and Developmental Biology, 2019, 7, 214.	3.7	7
21	Vertebroplasty and Return to Work for Thoracolumbar Fractures Within the Workers' Compensation Population. Spine, 2017, 42, 1024-1030.	2.0	6
22	Displaced Femoral Neck Fractures in Workers' Compensation Patients Aged 45-65 Years: Is It Best to Fix the Fracture or Replace the Joint?. Journal of Arthroplasty, 2020, 35, 3195-3203.	3.1	5
23	Lumbar Diskography and Failed Back Syndrome in Patients Receiving Workers' Compensation. Orthopedics, 2015, 38, e951-8.	1.1	5
24	Shorter Time to Surgery Is Associated With Better Outcomes for Spondylolisthesis in the Workers' Compensation Population. Orthopedics, 2020, 43, 154-160.	1.1	5
25	Multilevel Lumbar Fusion Is a Risk Factor for Lower Return to Work Rates Among Workers' Compensation Subjects With Degenerative Disc Disease. Journal of Surgical Orthopaedic Advances, 2018, 27, 209-218.	0.1	5
26	Prolonged Preoperative Opioid Therapy in Patients with Degenerative Lumbar Stenosis in a Workers' Compensation Setting. Spine Journal, 2017, 17, S203.	1.3	3
27	Time to Surgery Affects Return to Work Rates for Workers' Compensation Patients With Single-Level Lumbar Disk Herniation. Orthopedics, 2021, 44, e43-e49.	1.1	2
28	Lumbar Discography Is Associated With Poor Return to Work Status Following Lumbar Fusion Surgery in a Workers' Compensation Setting. Journal of Surgical Orthopaedic Advances, 2018, 27, 25-32.	0.1	2
29	Evaluation of Factors Affecting Return to Work Following Carpal Tunnel Release: A Statewide Cohort Study of Workers' Compensation Subjects. Journal of Hand Surgery, 2022, 47, 544-553.	1.6	2
30	Return-to-Work Rates after Single Level Cervical Fusion Surgery for Degenerative Disc Disease (DDD) Compared to Fusion for Radiculopathy in Workers' Compensation (WC) Setting. Spine Journal, 2015, 15, S106.	1.3	1
31	Prognostic Factors for Return to Work Status following Lumbar Fusion among Workers' Compensation Subjects with Spondylolisthesis. Spine Journal, 2016, 16, S367.	1.3	1
32	Kyphoplasty vs Orthosis for Treatment of Thoracolumbar Vertebral Fractures in the Workers' Compensation Population. Spine Journal, 2017, 17, S217-S218.	1.3	1
33	Presurgical Patient Evaluation with Lumbar Discography is Associated with Significantly More Days Absent from Work following Lumbar Fusion in a Workers' Compensation Setting. Spine Journal, 2014, 14, S102-S103.	1.3	0
34	Lumbar Fusion for Degenerative Disc Disease is Associated with Significantly Higher Rates of Failed Back Surgery Syndrome Compared to Fusion for Spondylolisthesis in a Workers' Compensation Setting. Spine Journal, 2014, 14, S71.	1.3	0
35	Clinical Depression is a Strong Risk Factor for Poor Return to Work Rates Following Lumbar Fusion Surgery in a Workers Compensation Setting. Spine Journal, 2014, 14, S68-S69.	1.3	0
36	Preoperative Duration of Opioid Use and Return to Work (RTW) Rates after Single-Level Cervical Fusion in Workers' Compensation (WC) Setting. Spine Journal, 2015, 15, S234-S235.	1.3	0

#	ARTICLE	IF	CITATIONS
37	Predictors of Chronic Opioid Therapy after Lumbar Fusion Surgery for Degenerative Disc Disease in a Workers' Compensation Setting. Spine Journal, 2016, 16, S343.	1.3	0
38	Opioid Use Predicts Poor Return-to-Work in Workers' Compensation Patients after Lumbar Discectomy. Spine Journal, 2017, 17, S68-S69.	1.3	0
39	The Impact of Smoking in Workers' Compensation Patients Receiving Spinal Cord Stimulation. Journal of Surgical Orthopaedic Advances, 2021, 30, 185-189.	0.1	0
40	The Role of Antiseptic Irrigation Solutions and Topical Antibiotics in Total Joint Arthroplasty.. Journal of Surgical Orthopaedic Advances, 2021, 30, 226-230.	0.1	0