

Eric Verdon

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4091507/publications.pdf>

Version: 2024-02-01

66
papers

1,553
citations

304743

22
h-index

345221

36
g-index

66
all docs

66
docs citations

66
times ranked

1910
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|--|------|-----------|
| 1 | Tissue distribution, metabolism, and elimination of Victoria Pure Blue BO in rainbow trout: Main metabolite as an appropriate residue marker. <i>Chemosphere</i> , 2021, 262, 127636. | 8.2 | 4 |
| 2 | Composite inclusion complexes containing hyaluronic acid/chitosan nanosystems for dual responsive enrofloxacin release. <i>Carbohydrate Polymers</i> , 2021, 252, 117162. | 10.2 | 33 |
| 3 | Confirmation of five nitrofuran metabolites including nifursol metabolite in meat and aquaculture products by liquid chromatography-tandem mass spectrometry: Validation according to European Union Decision 2002/657/EC. <i>Food Chemistry</i> , 2021, 342, 128389. | 8.2 | 24 |
| 4 | Evaluation of ELISA kits for the screening of four nitrofuran metabolites in aquaculture products according to the European guideline for the validation of screening methods. <i>Food Additives and Contaminants - Part A Chemistry, Analysis, Control, Exposure and Risk Assessment</i> , 2021, 38, 237-254. | 2.3 | 4 |
| 5 | Development and applicability of a multi-residue method for dyes, including new residue markers, to detect drug misuse in aquaculture. <i>Food Additives and Contaminants - Part A Chemistry, Analysis, Control, Exposure and Risk Assessment</i> , 2021, 38, 1332-1349. | 2.3 | 1 |
| 6 | Development of a magnetic MOF-based M-D-1/4SPE methodology combined with LC-MS/MS for the determination of fluorotelomer alcohols and its metabolites in animal derived foods. <i>Food Chemistry</i> , 2021, 363, 130205. | 8.2 | 6 |
| 7 | Preparation and evaluation of valnemulin hydrochloride taste-masking granules. <i>Current Drug Delivery</i> , 2021, 18, . | 1.6 | 1 |
| 8 | Dosage Regimen Formulation and Therapeutic Effect Evaluation of Cyadox Nanosuspension Against Dairy Cow Mastitis Caused by <i>Staphylococcus aureus</i> . <i>Current Drug Delivery</i> , 2021, 18, 965-974. | 1.6 | 1 |
| 9 | Surface plasmon resonance biosensor for the determination of 3-methyl-quinoxaline-2-carboxylic acid, the marker residue of olaquinox, in swine tissues. <i>Food Chemistry</i> , 2020, 302, 124623. | 8.2 | 14 |
| 10 | In vitro investigations of the metabolism of Victoria pure blue BO dye to identify main metabolites for food control in fish. <i>Chemosphere</i> , 2020, 238, 124538. | 8.2 | 7 |
| 11 | Development of a multi-class method to determine nitroimidazoles, nitrofurans, pharmacologically active dyes and chloramphenicol in aquaculture products by liquid chromatography-tandem mass spectrometry. <i>Food Chemistry</i> , 2020, 311, 125924. | 8.2 | 52 |
| 12 | Absorption, distribution, metabolism, and excretion of nanocarriers in vivo and their influences. <i>Advances in Colloid and Interface Science</i> , 2020, 284, 102261. | 14.7 | 83 |
| 13 | Optimal regimens based on PK/PD cutoff evaluation of ceftiofur against <i>Actinobacillus pleuropneumoniae</i> in swine. <i>BMC Veterinary Research</i> , 2020, 16, 366. | 1.9 | 7 |
| 14 | Designing, structural determination and biological effects of rifaximin loaded chitosan-carboxymethyl chitosan nanogel. <i>Carbohydrate Polymers</i> , 2020, 248, 116782. | 10.2 | 65 |
| 15 | The dose regimen formulation of tilmicosin against <i>Lawsonia intracellularis</i> in pigs by pharmacokinetic-pharmacodynamic (PK-PD) model. <i>Microbial Pathogenesis</i> , 2020, 147, 104389. | 2.9 | 12 |
| 16 | Design, Synthesis, and Biological Evaluation of Novel Thiazolidinone-Containing Quinoxaline-1,4-di-N-oxides as Antimycobacterial and Antifungal Agents. <i>Frontiers in Chemistry</i> , 2020, 8, 598. | 3.6 | 18 |
| 17 | Control of Antimicrobials in Feed Using Liquid Chromatography-Tandem Mass Spectrometry: Assessment of Cross-Contamination Rates at the Farm Level. <i>Journal of Agricultural and Food Chemistry</i> , 2020, 68, 9033-9042. | 5.2 | 6 |
| 18 | Intracellular delivery, accumulation, and discrepancy in antibacterial activity of four enrofloxacin-loaded fatty acid solid lipid nanoparticles. <i>Colloids and Surfaces B: Biointerfaces</i> , 2020, 194, 111196. | 5.0 | 18 |

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 19 | Disposition of cyadox in domesticated cats following oral, intramuscular, and intravenous administration. <i>Journal of Veterinary Pharmacology and Therapeutics</i> , 2020, 43, 97-107. | 1.3 | 2 |
| 20 | Tissue distribution and bioaccumulation of 8:2 fluorotelomer alcohol and its metabolites in pigs after oral exposure. <i>Chemosphere</i> , 2020, 249, 126016. | 8.2 | 11 |
| 21 | Development and optimisation of an amperometric immunosensor for the detection of banned antibiotic residues in honey. , 2020, 60, . | | 0 |
| 22 | Enhanced Treatment Effects of Tilmicosin Against <i>Staphylococcus aureus</i> Cow Mastitis by Self-Assembly Sodium Alginate-Chitosan Nanogel. <i>Pharmaceutics</i> , 2019, 11, 524. | 4.5 | 35 |
| 23 | Determination of Tartrazine, Lutein, Capsanthin, Canthaxanthin and β -Carotene in Animal-Derived Foods and Feeds by HPLC Method. <i>Journal of Chromatographic Science</i> , 2019, 57, 462-468. | 1.4 | 10 |
| 24 | A non-targeted LC-HRMS approach for detecting exposure to illegal veterinary treatments: The case of cephalosporins in commercial laying Hens. <i>Journal of Chromatography A</i> , 2019, 1599, 161-171. | 3.7 | 8 |
| 25 | Establishment of pressurized liquid extraction followed by HPLC-MS/MS method for the screening of adrenergic drugs, steroids, sedatives, colorants and antioxidants in swine feed. <i>Journal of Separation Science</i> , 2019, 42, 1915-1929. | 2.5 | 5 |
| 26 | Development of a broad-spectrum monoclonal antibody-based indirect competitive enzyme-linked immunosorbent assay for the multi-residue detection of avermectins in edible animal tissues and milk. <i>Food Chemistry</i> , 2019, 286, 234-240. | 8.2 | 37 |
| 27 | Simultaneous determination of multicomponent of acetylkitasamycin and kitasamycin by LC-MS/MS in swine plasma and its application in a pharmacokinetic study. <i>Biomedical Chromatography</i> , 2018, 32, e4268. | 1.7 | 4 |
| 28 | Construction of Electrochemical Immunosensor Based on Gold-Nanoparticles/Carbon Nanotubes/Chitosan for Sensitive Determination of T-2 Toxin in Feed and Swine Meat. <i>International Journal of Molecular Sciences</i> , 2018, 19, 3895. | 4.1 | 28 |
| 29 | A Convenient and Sensitive LC-MS/MS Method for Simultaneous Determination of Carbadox- and Olaquinox-Related Residues in Swine Muscle and Liver Tissues. <i>Journal of Analytical Methods in Chemistry</i> , 2018, 2018, 1-9. | 1.6 | 2 |
| 30 | The antibacterial activities of aditoprim and its efficacy in the treatment of swine streptococcosis. <i>Scientific Reports</i> , 2017, 7, 41370. | 3.3 | 8 |
| 31 | Preparation, characterization and pharmacokinetics of cyadox nanosuspension. <i>Scientific Reports</i> , 2017, 7, 2289. | 3.3 | 33 |
| 32 | An immunoaffinity column for the selective purification of 3-methyl-quinoxaline-2-carboxylic acid from swine tissues and its determination by high-performance liquid chromatography with ultraviolet detection and a colloidal gold-based immunochromatographic assay. <i>Food Chemistry</i> , 2017, 237, 290-296. | 8.2 | 13 |
| 33 | Solid-phase microextraction set-up for the analysis of liver volatolome to detect livestock exposure to micropollutants. <i>Journal of Chromatography A</i> , 2017, 1497, 9-18. | 3.7 | 12 |
| 34 | Simultaneous Determination of Quinoxalines in Animal Feeds by a Modified QuEChERS Method with MWCNTs as the Sorbent Followed by High-Performance Liquid Chromatography. <i>Food Analytical Methods</i> , 2017, 10, 2085-2091. | 2.6 | 11 |
| 35 | Preparation of a monoclonal antibody against amantadine and rimantadine and development of an indirect competitive enzyme-linked immunosorbent assay for detecting the same in chicken muscle and liver. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2017, 133, 56-63. | 2.8 | 28 |
| 36 | Preparation of a generic monoclonal antibody and development of a highly sensitive indirect competitive ELISA for the detection of phenothiazines in animal feed. <i>Food Chemistry</i> , 2017, 221, 1004-1013. | 8.2 | 36 |

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 37 | Development and Validation of a Monoclonal Antibody-Based Indirect Competitive ELISA for the Detection of Sudan I in Duck Eggs and Crystal Violet in Carp. <i>Food Analytical Methods</i> , 2017, 10, 1442-1451. | 2.6 | 5 |
| 38 | Development and validation of a multiclass method for the determination of antibiotic residues in honey using liquid chromatography-tandem mass spectrometry. <i>Food Additives and Contaminants - Part A Chemistry, Analysis, Control, Exposure and Risk Assessment</i> , 2017, 34, 582-597. | 2.3 | 33 |
| 39 | Pharmacokinetics and Metabolism of Cyadox and Its Main Metabolites in Beagle Dogs Following Oral, Intramuscular, and Intravenous Administration. <i>Frontiers in Pharmacology</i> , 2016, 7, 236. | 3.5 | 4 |
| 40 | Physiologically based pharmacokinetic model for quinocetone in pigs and extrapolation to mequindox. <i>Food Additives and Contaminants - Part A Chemistry, Analysis, Control, Exposure and Risk Assessment</i> , 2016, 34, 1-19. | 2.3 | 2 |
| 41 | Qualitative screening of veterinary anti-microbial agents in tissues, milk, and eggs of food-producing animals using liquid chromatography coupled with tandem mass spectrometry. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2016, 1017-1018, 82-88. | 2.3 | 69 |
| 42 | Development and validation of an indirect competitive enzyme-linked immunosorbent assay for the detection of albendazole 2-aminosulfone residues in animal tissues. <i>Food and Agricultural Immunology</i> , 2016, 27, 273-287. | 1.4 | 5 |
| 43 | Permethrin-induced oxidative stress and toxicity and metabolism. A review. <i>Environmental Research</i> , 2016, 149, 86-104. | 7.5 | 180 |
| 44 | Multiclass method for the quantification of 92 veterinary antimicrobial drugs in livestock excreta, wastewater, and surface water by liquid chromatography with tandem mass spectrometry. <i>Journal of Separation Science</i> , 2016, 39, 4086-4095. | 2.5 | 17 |
| 45 | Development and Validation of a Sensitive Indirect Competitive Enzyme-Linked Immunosorbent Assay for the Screening of Florfenicol and Thiamphenicol in Edible Animal Tissue and Feed. <i>Food Analytical Methods</i> , 2016, 9, 2434-2443. | 2.6 | 17 |
| 46 | Preparation of a Broadly Specific Monoclonal Antibody-Based Indirect Competitive ELISA for the Detection of Benzodiazepines in Edible Animal Tissues and Feed. <i>Food Analytical Methods</i> , 2016, 9, 3407-3419. | 2.6 | 8 |
| 47 | Development and validation of a sensitive monoclonal antibody-based indirect competitive enzyme-linked immunosorbent assay for the determination of the aflatoxin M1 levels in milk. <i>Toxicon</i> , 2016, 113, 18-24. | 1.6 | 17 |
| 48 | Development a monoclonal antibody-based enzyme-linked immunosorbent assay for screening carotenoids in eggs. <i>Food Chemistry</i> , 2016, 202, 141-148. | 8.2 | 8 |
| 49 | Synthesis, 3D-QSAR analysis and biological evaluation of quinoxaline 1,4-di-N-oxide derivatives as antituberculosis agents. <i>Bioorganic and Medicinal Chemistry Letters</i> , 2016, 26, 4146-4153. | 2.2 | 23 |
| 50 | Elimination and Concentration Correlations between Edible Tissues and Biological Fluids and Hair of Ractopamine in Pigs and Goats Fed with Ractopamine-Medicated Feed. <i>Journal of Agricultural and Food Chemistry</i> , 2016, 64, 2012-2020. | 5.2 | 10 |
| 51 | Development of a sensitive monoclonal-based enzyme-linked immunosorbent assay for monitoring T-2 toxin in food and feed. <i>Food Additives and Contaminants - Part A Chemistry, Analysis, Control, Exposure and Risk Assessment</i> , 2016, 33, 1-10. | 2.3 | 8 |
| 52 | Development of a monoclonal antibody-based indirect competitive enzyme-linked immunosorbent assay for nitroimidazoles in edible animal tissues and feeds. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2016, 120, 84-91. | 2.8 | 16 |
| 53 | Development and validation of an indirect competitive enzyme-linked immunosorbent assay for monitoring organoarsenic compounds in edible chicken and pork and feed. <i>Food Chemistry</i> , 2016, 197, 821-828. | 8.2 | 18 |
| 54 | Integration of PK/PD for dose optimization of Cefquinome against <i>Staphylococcus aureus</i> causing septicemia in cattle. <i>Frontiers in Microbiology</i> , 2015, 6, 588. | 3.5 | 32 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 55 | Validation of a liquid chromatography–high-resolution mass spectrometry method for the analysis of ceftiofur in poultry muscle, kidneys and plasma: A unique accuracy profile for each and every matrix. <i>Journal of Chromatography A</i> , 2015, 1407, 119-129. | 3.7 | 11 |
| 56 | Metabolism, Distribution, and Elimination of Mequindox in Pigs, Chickens, and Rats. <i>Journal of Agricultural and Food Chemistry</i> , 2015, 63, 9839-9849. | 5.2 | 31 |
| 57 | A novel hapten and monoclonal-based enzyme-linked immunosorbent assay for 3-methyl-quinoxaline-2-carboxylic acid in edible animal tissues. <i>Analytical Methods</i> , 2015, 7, 6588-6594. | 2.7 | 12 |
| 58 | The Monitoring of Triphenylmethane Dyes in Aquaculture Products Through the European Union Network of Official Control Laboratories. <i>Journal of AOAC INTERNATIONAL</i> , 2015, 98, 649-657. | 1.5 | 10 |
| 59 | Microbiological toxicity of tilmicosin on human colonic microflora in chemostats. <i>Regulatory Toxicology and Pharmacology</i> , 2015, 73, 201-208. | 2.7 | 8 |
| 60 | Metabolic disposition and excretion of quinocetone in rats, pigs, broilers, and carp. <i>Food and Chemical Toxicology</i> , 2014, 69, 109-119. | 3.6 | 29 |
| 61 | Development of a liquid chromatography–tandem mass spectrometry with ultrasound-assisted extraction and auto solid-phase clean-up method for the determination of Fusarium toxins in animal derived foods. <i>Journal of Chromatography A</i> , 2013, 1311, 21-29. | 3.7 | 30 |
| 62 | Development of a liquid chromatography–tandem mass spectrometry with ultrasound-assisted extraction method for the simultaneous determination of sudan dyes and their metabolites in the edible tissues and eggs of food-producing animals. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2013, 939, 45-50. | 2.3 | 42 |
| 63 | Determination of Residues of Three Triphenylmethane Dyes and Their Metabolites (Malachite Green,) Tj ETQq1 1 0.784314 rgBT /Overlock 10 Tf 50 31 Products by LC/MS/MS: First Action 2012.25. <i>Journal of AOAC INTERNATIONAL</i> , 2013, 96, 1152-1157. | 1.5 | 21 |
| 64 | Development of a sensitive and robust liquid chromatography coupled with tandem mass spectrometry and a pressurized liquid extraction for the determination of aflatoxins and ochratoxin A in animal derived foods. <i>Journal of Chromatography A</i> , 2012, 1253, 110-119. | 3.7 | 58 |
| 65 | Simultaneous determination of malachite green, gentian violet and their leuco-metabolites in shrimp and salmon by liquid chromatography–tandem mass spectrometry with accelerated solvent extraction and auto solid-phase clean-up. <i>Food Control</i> , 2011, 22, 1246-1252. | 5.5 | 67 |
| 66 | Multi-residue monitoring for the simultaneous determination of five nitrofurans (furazolidone,) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 31 detection of their five major metabolites (AOZ, AMOZ, SEM, AHD, DNSAH) by liquid chromatography coupled to electrospray tandem mass spectrometry–In-house validation in line with Commission Decision 657/2002/EC. <i>Analytica Chimica Acta</i> , 2007, 586, 336-347. | 5.4 | 123 |