

Herbert Schwarz

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4099396/publications.pdf>

Version: 2024-02-01

90
papers

2,625
citations

172457

29
h-index

214800

47
g-index

92
all docs

92
docs citations

92
times ranked

2361
citing authors

#	ARTICLE	IF	CITATIONS
1	Dendritic cell therapy with CD137L-DC-EBV-VAX in locally recurrent or metastatic nasopharyngeal carcinoma is safe and confers clinical benefit. <i>Cancer Immunology, Immunotherapy</i> , 2022, 71, 1531-1543.	4.2	11
2	Novel Autoantibodies in Idiopathic Small Fiber Neuropathy. <i>Annals of Neurology</i> , 2022, 91, 66-77.	5.3	9
3	Epstein-Barr virus-induced ectopic CD137 expression helps nasopharyngeal carcinoma to escape immune surveillance and enables targeting by chimeric antigen receptors. <i>Cancer Immunology, Immunotherapy</i> , 2022, 71, 2583-2596.	4.2	4
4	Soluble CD137 is a novel serum marker of liver cirrhosis in patients with hepatitis C and alcohol-associated disease etiology. <i>European Journal of Immunology</i> , 2022, 52, 633-645.	2.9	4
5	Ectopic CD137 expression by rhabdomyosarcoma provides selection advantages but allows immunotherapeutic targeting. <i>OncImmunity</i> , 2021, 10, 1877459.	4.6	4
6	Regulatory T Cells Inhibit T Cell Activity by Downregulating CD137 Ligand via CD137 Trogocytosis. <i>Cells</i> , 2021, 10, 353.	4.1	6
7	DUSP16 promotes cancer chemoresistance through regulation of mitochondria-mediated cell death. <i>Nature Communications</i> , 2021, 12, 2284.	12.8	28
8	B7-H7 Is Inducible on T Cells to Regulate Their Immune Response and Serves as a Marker for Exhaustion. <i>Frontiers in Immunology</i> , 2021, 12, 682627.	4.8	7
9	Integration of the Cortical Haemodynamic Response Measured by Functional Near-Infrared Spectroscopy and Amino Acid Analysis to Aid in the Diagnosis of Major Depressive Disorder. <i>Diagnostics</i> , 2021, 11, 1978.	2.6	4
10	No crossreactivity of anti-SARS-CoV-2 spike protein antibodies with Syncytin-1. <i>Cellular and Molecular Immunology</i> , 2021, 18, 2566-2568.	10.5	18
11	Exosomes, a New Star for Targeted Delivery. <i>Frontiers in Cell and Developmental Biology</i> , 2021, 9, 751079.	3.7	104
12	CD137 negatively affects "browning" of white adipose tissue during cold exposure. <i>Journal of Biological Chemistry</i> , 2020, 295, 2034-2042.	3.4	13
13	CD137 Ligand-CD137 Interaction is Required For Inflammasome-Associated Brain Injury Following Ischemic Stroke. <i>NeuroMolecular Medicine</i> , 2020, 22, 474-483.	3.4	9
14	The role of trogocytosis in immune surveillance of Hodgkin lymphoma. <i>OncImmunity</i> , 2020, 9, 1781334.	4.6	9
15	Identification of CD137-Expressing B Cells in Multiple Sclerosis Which Secrete IL-6 Upon Engagement by CD137 Ligand. <i>Frontiers in Immunology</i> , 2020, 11, 571964.	4.8	9
16	Resveratrol attenuates TLR-4 mediated inflammation and elicits therapeutic potential in models of sepsis. <i>Scientific Reports</i> , 2020, 10, 18837.	3.3	14
17	CD137 / CD137 ligand signalling regulates the immune balance: A potential target for novel immunotherapy of autoimmune diseases. <i>Journal of Autoimmunity</i> , 2020, 112, 102499.	6.5	21
18	CD137 ligand interacts with CD32a to trigger reverse CD137 ligand signaling. <i>Cellular and Molecular Immunology</i> , 2020, 17, 1188-1189.	10.5	4

#	ARTICLE	IF	CITATIONS
19	Destroy, what destroys you. <i>Oncolmmunology</i> , 2020, 9, 1685301.	4.6	4
20	The relevance of soluble CD137 in the regulation of immune responses and for immunotherapeutic intervention. <i>Journal of Leukocyte Biology</i> , 2020, 107, 731-738.	3.3	21
21	Deletion of CD137 Ligand Exacerbates Renal and Cutaneous but Alleviates Cerebral Manifestations in Lupus. <i>Frontiers in Immunology</i> , 2019, 10, 1411.	4.8	11
22	Development of a Bispecific Antibody Targeting CD30 and CD137 on Hodgkin and Reed-Sternberg Cells. <i>Frontiers in Oncology</i> , 2019, 9, 945.	2.8	16
23	CD137L-DCs, Potent Immune-Stimulatorsâ€™History, Characteristics, and Perspectives. <i>Frontiers in Immunology</i> , 2019, 10, 2216.	4.8	21
24	The Progress of Investigating the CD137-CD137L Axis as a Potential Target for Systemic Lupus Erythematosus. <i>Cells</i> , 2019, 8, 1044.	4.1	6
25	Increased Akt-Driven Glycolysis Is the Basis for the Higher Potency of CD137L-DCs. <i>Frontiers in Immunology</i> , 2019, 10, 868.	4.8	11
26	Epsteinâ€™Barr virus-encoded LMP1 induces ectopic CD137 expression on Hodgkin and Reedâ€™Sternberg cells via the PI3K-AKT-mTOR pathway. <i>Leukemia and Lymphoma</i> , 2019, 60, 2697-2704.	1.3	20
27	Induction of CD137 expression by viral genes reduces T cell costimulation. <i>Journal of Cellular Physiology</i> , 2019, 234, 21076-21088.	4.1	14
28	Plasma Factors for the Differentiation of Hodgkinâ€™s Lymphoma and Diffused Large B Cell Lymphoma and for Monitoring Remission. <i>Journal of Hematology (Brossard, Quebec)</i> , 2019, 8, 47-54.	1.0	5
29	CD137L dendritic cells induce potent response against cancer-associated viruses and polarize human CD8+ T cells to Tc1 phenotype. <i>Cancer Immunology, Immunotherapy</i> , 2018, 67, 893-905.	4.2	24
30	Anti-CD137 Cancer Immunotherapy Suppresses Tumor Growthâ€™Letter. <i>Cancer Research</i> , 2018, 78, 1571-1571.	0.9	5
31	Hiding in Plain Sight: Soluble Immunomodulatory Receptors. <i>Trends in Immunology</i> , 2018, 39, 771-774.	6.8	9
32	Endothelial dysfunction in systemic lupus erythematosus â€™ a case-control study and an updated meta-analysis and meta-regression. <i>Scientific Reports</i> , 2017, 7, 7320.	3.3	44
33	CD137 and CD137L signals are main drivers of type 1, cell-mediated immune responses. <i>Oncolmmunology</i> , 2016, 5, e1113367.	4.6	60
34	Transcriptional and functional characterization of CD137L-dendritic cells identifies a novel dendritic cell phenotype. <i>Scientific Reports</i> , 2016, 6, 29712.	3.3	10
35	HepaCAM associates with connexin 43 and enhances its localization in cellular junctions. <i>Scientific Reports</i> , 2016, 6, 36218.	3.3	27
36	CD137 signaling in Hodgkin and Reed-Sternberg cell lines induces IL-13 secretion, immune deviation and enhanced growth. <i>Oncolmmunology</i> , 2016, 5, e1160188.	4.6	18

#	ARTICLE	IF	CITATIONS
37	Sourcing of an Alternative Pericyte-Like Cell Type from Peripheral Blood in Clinically Relevant Numbers for Therapeutic Angiogenic Applications. <i>Molecular Therapy</i> , 2015, 23, 510-522.	8.2	28
38	Trogocytic CD137 transfer causes an internalization of CD137 ligand on murine APCs leading to reduced T cell costimulation. <i>Journal of Leukocyte Biology</i> , 2015, 97, 909-919.	3.3	17
39	4-1BB Chimeric Antigen Receptors. <i>Cancer Journal (Sudbury, Mass)</i> , 2014, 20, 134-140.	2.0	48
40	CD137 ligand signalling induces differentiation of primary acute myeloid leukaemia cells. <i>British Journal of Haematology</i> , 2014, 165, 134-144.	2.5	8
41	Role of the CD137 ligand (CD137L) signaling pathway during <i>Mycobacterium tuberculosis</i> infection. <i>Immunobiology</i> , 2014, 219, 78-86.	1.9	4
42	Regulation of Myelopoiesis by CD137L Signaling. <i>International Reviews of Immunology</i> , 2014, 33, 454-469.	3.3	7
43	Tumor necrosis factor receptor 1 associates with CD137 ligand and mediates its reverse signaling. <i>FASEB Journal</i> , 2013, 27, 2957-2966.	0.5	19
44	Ectopic CD137 expression facilitates the escape of Hodgkin and Reed-Sternberg cells from immunosurveillance. <i>Oncolmmunology</i> , 2013, 2, e23441.	4.6	13
45	CD137L-stimulated dendritic cells are more potent than conventional dendritic cells at eliciting cytotoxic T-cell responses. <i>Oncolmmunology</i> , 2013, 2, e26859.	4.6	27
46	Expression of CD137 on Hodgkin and Reed-Sternberg Cells Inhibits T-cell Activation by Eliminating CD137 Ligand Expression. <i>Cancer Research</i> , 2013, 73, 652-661.	0.9	64
47	CD137 ligand signaling enhances myelopoiesis during infections. <i>European Journal of Immunology</i> , 2013, 43, 1555-1567.	2.9	16
48	CD137 Facilitates the Resolution of Acute DSS-Induced Colonic Inflammation in Mice. <i>PLoS ONE</i> , 2013, 8, e73277.	2.5	13
49	CD137 ligand reverse signaling skews hematopoiesis towards myelopoiesis during aging. <i>Aging</i> , 2013, 5, 643-652.	3.1	10
50	Development of Experimental Autoimmune Encephalomyelitis Critically Depends on CD137 Ligand Signaling. <i>Journal of Neuroscience</i> , 2012, 32, 18246-18252.	3.6	32
51	CD137 ligand activated microglia induces oligodendrocyte apoptosis via reactive oxygen species. <i>Journal of Neuroinflammation</i> , 2012, 9, 173.	7.2	56
52	Admission levels of soluble CD137 are increased in patients with acute pancreatitis and are associated with subsequent complications. <i>Experimental and Molecular Pathology</i> , 2012, 92, 1-6.	2.1	10
53	Species Difference of CD137 Ligand Signaling in Human and Murine Monocytes. <i>PLoS ONE</i> , 2011, 6, e16129.	2.5	22
54	Biphasic activity of CD137 ligand-stimulated monocytes on T cell apoptosis and proliferation. <i>Journal of Leukocyte Biology</i> , 2011, 89, 707-720.	3.3	6

#	ARTICLE	IF	CITATIONS
55	Involvement of the Cytokine Receptor CD137 in Murine Hematopoiesis. <i>Advances in Experimental Medicine and Biology</i> , 2011, 691, 375-382.	1.6	2
56	CD137 ligand signaling induces human monocyte to dendritic cell differentiation. <i>European Journal of Immunology</i> , 2010, 40, 1938-1949.	2.9	48
57	CD137 ligand, a member of the tumor necrosis factor family, regulates immune responses via reverse signal transduction. <i>Journal of Leukocyte Biology</i> , 2010, 89, 21-29.	3.3	141
58	Inhibition of Proliferation and Induction of Apoptosis in Multiple Myeloma Cell Lines by CD137 Ligand Signaling. <i>PLoS ONE</i> , 2010, 5, e10845.	2.5	21
59	Regulation of Granulocyte and Macrophage Populations of Murine Bone Marrow Cells by G-CSF and CD137 Protein. <i>PLoS ONE</i> , 2010, 5, e15565.	2.5	29
60	CD137, implications in immunity and potential for therapy. <i>Frontiers in Bioscience - Landmark</i> , 2009, Volume, 4173.	3.0	42
61	CD137 ligand reverse signaling has multiple functions in human dendritic cells during an adaptive immune response. <i>European Journal of Immunology</i> , 2008, 38, 1024-1032.	2.9	54
62	Induction of Proliferation and Monocytic Differentiation of Human CD34+ Cells by CD137 Ligand Signaling. <i>Stem Cells</i> , 2008, 26, 2372-2381.	3.2	35
63	Characterisation of soluble murine CD137 and its association with systemic lupus. <i>Molecular Immunology</i> , 2008, 45, 3990-3999.	2.2	30
64	CD137 Induces Proliferation of Murine Hematopoietic Progenitor Cells and Differentiation to Macrophages. <i>Journal of Immunology</i> , 2008, 181, 3923-3932.	0.8	45
65	CD137 is expressed on blood vessel walls at sites of inflammation and enhances monocyte migratory activity. <i>FASEB Journal</i> , 2007, 21, 456-463.	0.5	101
66	Signal transduction mechanisms of CD137 ligand in human monocytes. <i>Cellular Signalling</i> , 2007, 19, 1899-1908.	3.6	33
67	OX40 interactions in gastrointestinal nematode infection. <i>Immunology</i> , 2006, 117, 108-116.	4.4	5
68	Significance of Reverse Signal Transduction for the Biology of the CD137 Receptor/Ligand System. , 2006, , 29-45.		1
69	Biological activities of reverse signal transduction through CD137 ligand. <i>Journal of Leukocyte Biology</i> , 2005, 77, 281-286.	3.3	67
70	Inhibition of cytokines expression in human microglia infected by virulent and non-virulent mycobacteria. <i>Neurochemistry International</i> , 2004, 44, 381-392.	3.8	40
71	OX40 (CD134) Blockade Inhibits the Co-stimulatory Cascade and Promotes Heart Allograft Survival. <i>Transplantation</i> , 2004, 78, 807-814.	1.0	36
72	Expression of CD137 and its ligand in human neurons, astrocytes, and microglia: Modulation by FGF–2. <i>Journal of Neuroscience Research</i> , 2003, 74, 67-73.	2.9	40

#	ARTICLE	IF	CITATIONS
73	CD137 is expressed by follicular dendritic cells and costimulates B lymphocyte activation in germinal centers. <i>Journal of Leukocyte Biology</i> , 2002, 72, 35-42.	3.3	60
74	In vitro and in vivo activities of OX40 (CD134)-IgG fusion protein isoforms with different levels of immune-effector functions. <i>Journal of Leukocyte Biology</i> , 2002, 72, 522-9.	3.3	18
75	Identification of a soluble OX40 isoform: development of a specific and quantitative immunoassay. <i>Journal of Immunological Methods</i> , 2001, 255, 67-72.	1.4	21
76	CD137 Expression in Tumor Vessel Walls. <i>American Journal of Clinical Pathology</i> , 2001, 115, 543-549.	0.7	87
77	Comparative Analysis of CD137 and LPS Effects on Monocyte Activation, Survival, and Proliferation. <i>Biochemical and Biophysical Research Communications</i> , 2000, 273, 117-122.	2.1	58
78	EXPRESSION OF SOLUBLE CD137 CORRELATES WITH ACTIVATION-INDUCED CELL DEATH OF LYMPHOCYTES. <i>Cytokine</i> , 2000, 12, 742-746.	3.2	48
79	CD137 Induces Proliferation and Endomitosis in Monocytes. <i>Blood</i> , 1999, 94, 3161-3168.	1.4	92
80	Identification of CD137 as a potent monocyte survival factor. <i>Journal of Leukocyte Biology</i> , 1999, 65, 829-833.	3.3	70
81	CD137 Induces Proliferation and Endomitosis in Monocytes. <i>Blood</i> , 1999, 94, 3161-3168.	1.4	5
82	A soluble form of CD137 (ILA/4-1BB), a member of the TNF receptor family, is released by activated lymphocytes and is detectable in sera of patients with rheumatoid arthritis. <i>European Journal of Immunology</i> , 1998, 28, 290-295.	2.9	126
83	A soluble form of CD137 (ILA/4-1BB), a member of the TNF receptor family, is released by activated lymphocytes and is detectable in sera of patients with rheumatoid arthritis. , 1998, 28, 290.		1
84	A soluble form of CD137 (ILA/4-1BB), a member of the TNF receptor family, is released by activated lymphocytes and is detectable in sera of patients with rheumatoid arthritis. , 1998, 28, 290.		2
85	CD137, a Member of the Tumor Necrosis Factor Receptor Family, Is Located on Chromosome 1p36, in a Cluster of Related Genes, and Colocalizes with Several Malignancies. <i>Biochemical and Biophysical Research Communications</i> , 1997, 235, 699-703.	2.1	23
86	Differentiation-dependent and stimulus-specific expression of ILA, the human 4-1BB-homologue, in cells of mesenchymal origin. <i>Osteoarthritis and Cartilage</i> , 1997, 5, 394-406.	1.3	19
87	Recovery of functional, recombinant baculovirus-produced proteins from insoluble precipitates in insect cells. <i>Technical Tips Online</i> , 1997, 2, 30-31.	0.2	0
88	The nerve growth factor/tumor necrosis factor receptor family. <i>Journal of Leukocyte Biology</i> , 1996, 60, 1-7.	3.3	54
89	A mRNA variant encoding a soluble form of 4-1BB, a member of the murine NGF/TNF receptor family. <i>Gene</i> , 1995, 164, 311-315.	2.2	53
90	A receptor induced by lymphocyte activation (ILA): a new member of the human nerve-growth-factor/tumor-necrosis-factor receptor family. <i>Gene</i> , 1993, 134, 295-298.	2.2	114