

Haiping Mao

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4123427/publications.pdf>

Version: 2024-02-01

96
papers

3,021
citations

147801

31
h-index

175258

52
g-index

101
all docs

101
docs citations

101
times ranked

4066
citing authors

#	ARTICLE	IF	CITATIONS
1	A novel STAT3 inhibitor, S3I-201, attenuates renal interstitial fibroblast activation and interstitial fibrosis in obstructive nephropathy. <i>Kidney International</i> , 2010, 78, 257-268.	5.2	219
2	Inhibition of histone deacetylase activity attenuates renal fibroblast activation and interstitial fibrosis in obstructive nephropathy. <i>American Journal of Physiology - Renal Physiology</i> , 2009, 297, F996-F1005.	2.7	188
3	Atg5-mediated autophagy deficiency in proximal tubules promotes cell cycle G ₂ /M arrest and renal fibrosis. <i>Autophagy</i> , 2016, 12, 1472-1486.	9.1	149
4	Clinical Outcome of Hyperuricemia in IgA Nephropathy: A Retrospective Cohort Study and Randomized Controlled Trial. <i>Kidney and Blood Pressure Research</i> , 2012, 35, 153-160.	2.0	127
5	Prevalence and risk factors associated with chronic kidney disease in an adult population from southern China. <i>Nephrology Dialysis Transplantation</i> , 2008, 24, 1205-1212.	0.7	125
6	ATG5-mediated autophagy suppresses NF- κ B signaling to limit epithelial inflammatory response to kidney injury. <i>Cell Death and Disease</i> , 2019, 10, 253.	6.3	105
7	Distinct hsp70 Domains Mediate Apoptosis-inducing Factor Release and Nuclear Accumulation. <i>Journal of Biological Chemistry</i> , 2006, 281, 7873-7880.	3.4	103
8	β -Catenin Promotes Survival of Renal Epithelial Cells by Inhibiting Bax. <i>Journal of the American Society of Nephrology: JASN</i> , 2009, 20, 1919-1928.	6.1	96
9	GSK3 β Promotes Apoptosis after Renal Ischemic Injury. <i>Journal of the American Society of Nephrology: JASN</i> , 2010, 21, 284-294.	6.1	94
10	HSP72 attenuates renal tubular cell apoptosis and interstitial fibrosis in obstructive nephropathy. <i>American Journal of Physiology - Renal Physiology</i> , 2008, 295, F202-F214.	2.7	91
11	The impact of peritoneal dialysis-related peritonitis on mortality in peritoneal dialysis patients. <i>BMC Nephrology</i> , 2017, 18, 186.	1.8	90
12	HSP72 inhibits apoptosis-inducing factor release in ATP-depleted renal epithelial cells. <i>American Journal of Physiology - Cell Physiology</i> , 2003, 285, C1483-C1493.	4.6	78
13	Drp1-mediated mitochondrial fission promotes renal fibroblast activation and fibrogenesis. <i>Cell Death and Disease</i> , 2020, 11, 29.	6.3	73
14	A Crosstalk between the Smad and JNK Signaling in the TGF- β 2-Induced Epithelial-Mesenchymal Transition in Rat Peritoneal Mesothelial Cells. <i>PLoS ONE</i> , 2012, 7, e32009.	2.5	64
15	HSP72 Inhibits Smad3 Activation and Nuclear Translocation in Renal Epithelial-to-Mesenchymal Transition. <i>Journal of the American Society of Nephrology: JASN</i> , 2010, 21, 598-609.	6.1	60
16	The Effect of Fluid Overload on Clinical Outcome in Southern Chinese Patients Undergoing Continuous Ambulatory Peritoneal Dialysis. <i>Peritoneal Dialysis International</i> , 2015, 35, 691-702.	2.3	60
17	Downregulation of Par-3 expression and disruption of Par complex integrity by TGF- β 2 during the process of epithelial to mesenchymal transition in rat proximal epithelial cells. <i>Biochimica Et Biophysica Acta - Molecular Basis of Disease</i> , 2008, 1782, 51-59.	3.8	57
18	Transforming Growth Factor β 21 Induces Epithelial-to-mesenchymal Transition by Activating the Jnk-SMAD3 Pathway in Rat Peritoneal Mesothelial Cells. <i>Peritoneal Dialysis International</i> , 2008, 28, 88-95.	2.3	55

#	ARTICLE	IF	CITATIONS
19	Elevated neutrophil to lymphocyte ratio predicts overall and cardiovascular mortality in maintenance peritoneal dialysis patients. <i>International Urology and Nephrology</i> , 2012, 44, 1521-1528.	1.4	55
20	Heat Shock Protein 72 Enhances Autophagy as a Protective Mechanism in Lipopolysaccharide-Induced Peritonitis in Rats. <i>American Journal of Pathology</i> , 2011, 179, 2822-2834.	3.8	49
21	hsp72 Inhibits Focal Adhesion Kinase Degradation in ATP-depleted Renal Epithelial Cells. <i>Journal of Biological Chemistry</i> , 2003, 278, 18214-18220.	3.4	45
22	Alkaline Phosphatase and Mortality in Patients on Peritoneal Dialysis. <i>Clinical Journal of the American Society of Nephrology: CJASN</i> , 2014, 9, 771-778.	4.5	44
23	Inhibition of EGF Receptor Blocks the Development and Progression of Peritoneal Fibrosis. <i>Journal of the American Society of Nephrology: JASN</i> , 2016, 27, 2631-2644.	6.1	43
24	Clinical outcome and risk factors for mortality in Chinese patients with diabetes on peritoneal dialysis: A 5-year clinical cohort study. <i>Diabetes Research and Clinical Practice</i> , 2013, 100, 354-361.	2.8	41
25	Serum Potassium Levels and Its Variability in Incident Peritoneal Dialysis Patients: Associations with Mortality. <i>PLoS ONE</i> , 2014, 9, e86750.	2.5	41
26	<i>Escherichia Coli</i> Peritonitis in Peritoneal Dialysis: The Prevalence, Antibiotic Resistance and Clinical Outcomes in a South China Dialysis Center. <i>Peritoneal Dialysis International</i> , 2014, 34, 308-316.	2.3	39
27	Prevalence and risk factors of sleep disturbance in continuous ambulatory peritoneal dialysis patients in Guangzhou, southern China. <i>International Urology and Nephrology</i> , 2012, 44, 929-936.	1.4	35
28	Low α -defensin gene copy number increases the risk for IgA nephropathy and renal dysfunction. <i>Science Translational Medicine</i> , 2016, 8, 345ra88.	12.4	35
29	Transforming growth factor β 1 induces epithelial-mesenchymal transition by activating the JNK-Smad3 pathway in rat peritoneal mesothelial cells. <i>Peritoneal Dialysis International</i> , 2008, 28 Suppl 3, S88-95.	2.3	34
30	Decreased expressions of the TNF-alpha signaling adapters in peripheral blood mononuclear cells (PBMCs) are correlated with disease activity in patients with systemic lupus erythematosus. <i>Clinical Rheumatology</i> , 2007, 26, 1481-1489.	2.2	32
31	Association between depression and malnutrition-inflammatory complex syndrome in patients with continuous ambulatory peritoneal dialysis. <i>International Urology and Nephrology</i> , 2011, 43, 875-882.	1.4	32
32	MicroRNA-21 is Overexpressed in Renal Cell Carcinoma. <i>International Journal of Biological Markers</i> , 2013, 28, 201-207.	1.8	30
33	Urgent-start peritoneal dialysis for patients with end stage renal disease: a 10-year retrospective study. <i>BMC Nephrology</i> , 2019, 20, 238.	1.8	27
34	Is cystatin C a better marker than creatinine for evaluating residual renal function in patients on continuous ambulatory peritoneal dialysis?. <i>Nephrology Dialysis Transplantation</i> , 2011, 26, 3358-3365.	0.7	25
35	Malnutrition-inflammation score predicts long-term mortality in Chinese PD patients. <i>Clinical Nephrology</i> , 2013, 79, 477-483.	0.7	25
36	Hsp72 Interacts with Paxillin and Facilitates the Reassembly of Focal Adhesions during Recovery from ATP Depletion. <i>Journal of Biological Chemistry</i> , 2004, 279, 15472-15480.	3.4	24

#	ARTICLE	IF	CITATIONS
37	Src regulates cell cycle protein expression and renal epithelial cell proliferation via PI3K/Akt signaling-dependent and -independent mechanisms. <i>American Journal of Physiology - Renal Physiology</i> , 2008, 295, F145-F152.	2.7	24
38	Association of Pulmonary Hypertension with Mortality in Incident Peritoneal Dialysis Patients. <i>Peritoneal Dialysis International</i> , 2015, 35, 537-544.	2.3	22
39	Impaired TGF- β signalling enhances peritoneal inflammation induced by E. Coli in rats. <i>Nephrology Dialysis Transplantation</i> , 2010, 25, 399-412.	0.7	21
40	Association Analysis of the MHC in Lupus Nephritis. <i>Journal of the American Society of Nephrology: JASN</i> , 2017, 28, 3383-3394.	6.1	21
41	V-ATPase promotes transforming growth factor- β -induced epithelial-mesenchymal transition of rat proximal tubular epithelial cells. <i>American Journal of Physiology - Renal Physiology</i> , 2012, 302, F1121-F1132.	2.7	20
42	High Prevalence and Associated Risk Factors for Impaired Renal Function and Urinary Abnormalities in a Rural Adult Population from Southern China. <i>PLoS ONE</i> , 2012, 7, e47100.	2.5	20
43	Identification of susceptibility locus shared by IgA nephropathy and inflammatory bowel disease in a Chinese Han population. <i>Journal of Human Genetics</i> , 2020, 65, 241-249.	2.3	20
44	Prevalence and risk factors of exit-site infection in incident peritoneal dialysis patients. <i>Peritoneal Dialysis International</i> , 2020, 40, 164-170.	2.3	19
45	Serum magnesium and cardiovascular mortality in peritoneal dialysis patients: a 5-year prospective cohort study. <i>British Journal of Nutrition</i> , 2018, 120, 415-423.	2.3	18
46	Clinicopathologic features and treatment response in nephrotic IgA nephropathy with minimal change disease. <i>Clinical Nephrology</i> , 2013, 79, 37-44.	0.7	18
47	Prevalence and Factors Associated with Hypomagnesemia in Southern Chinese Continuous Ambulatory Peritoneal Dialysis Patients. <i>Peritoneal Dialysis International</i> , 2013, 33, 450-454.	2.3	17
48	The Potential Role of HMGB1 Release in Peritoneal Dialysis-Related Peritonitis. <i>PLoS ONE</i> , 2013, 8, e54647.	2.5	17
49	Association of Body Mass Index and Body Mass Index Change with Mortality in Incident Peritoneal Dialysis Patients. <i>Nutrients</i> , 2015, 7, 8444-8455.	4.1	16
50	Uric acid to high-density lipoprotein cholesterol ratio predicts cardiovascular mortality in patients on peritoneal dialysis. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2021, 31, 561-569.	2.6	15
51	Association of baseline, longitudinal serum high-sensitive C-reactive protein and its change with mortality in peritoneal dialysis patients. <i>BMC Nephrology</i> , 2017, 18, 211.	1.8	13
52	Heat Shock Protein 72 Antagonizes STAT3 Signaling to Inhibit Fibroblast Accumulation in Renal Fibrogenesis. <i>American Journal of Pathology</i> , 2016, 186, 816-828.	3.8	12
53	Autophagy protects against necrotic renal epithelial cell-induced death of renal interstitial fibroblasts. <i>American Journal of Physiology - Renal Physiology</i> , 2012, 303, F83-F91.	2.7	11
54	The predictive study of the relation between elevated low-density lipoprotein cholesterol to high-density lipoprotein cholesterol ratio and mortality in peritoneal dialysis. <i>Lipids in Health and Disease</i> , 2020, 19, 51.	3.0	11

#	ARTICLE	IF	CITATIONS
55	Gender Difference in the Association of Hyperuricemia with Chronic Kidney Disease in Southern China. <i>Kidney and Blood Pressure Research</i> , 2012, 36, 98-106.	2.0	10
56	Association of ITGAX and ITGAM gene polymorphisms with susceptibility to IgA nephropathy. <i>Journal of Human Genetics</i> , 2019, 64, 927-935.	2.3	10
57	Association of Lean Body Mass Index and Peritoneal Protein Clearance in Peritoneal Dialysis Patients. <i>Kidney and Blood Pressure Research</i> , 2019, 44, 94-102.	2.0	10
58	Prevalence and risk factors of chronic kidney disease in first-degree relatives of chronic kidney disease patients in Southern China. <i>Nephrology</i> , 2012, 17, 123-130.	1.6	9
59	Clinical Outcomes of Peritoneal Dialysis Patients Transferred from Hemodialysis: A Matched Case-Control Study. <i>Peritoneal Dialysis International</i> , 2013, 33, 259-266.	2.3	9
60	Elevated Serum Trimethylamine N-Oxide Levels Are Associated with Mortality in Male Patients on Peritoneal Dialysis. <i>Blood Purification</i> , 2021, 50, 837-847.	1.8	9
61	Serum Phosphorus and Albumin in Patients Undergoing Peritoneal Dialysis: Interaction and Association With Mortality. <i>Frontiers in Medicine</i> , 2021, 8, 760394.	2.6	9
62	Heat shock protein 72 suppresses apoptosis by increasing the stability of X-linked inhibitor of apoptosis protein in renal ischemia/reperfusion injury. <i>Molecular Medicine Reports</i> , 2015, 11, 1793-1799.	2.4	8
63	Prevalence of erectile dysfunction and its association with residual renal function in Chinese peritoneal dialysis patients. <i>International Urology and Nephrology</i> , 2015, 47, 383-389.	1.4	8
64	Patient Survival and Technique Failure in Continuous Ambulatory Peritoneal Dialysis Patients with Prior Stroke. <i>Peritoneal Dialysis International</i> , 2016, 36, 308-314.	2.3	8
65	Association of <i>FCRL3</i> Gene Polymorphisms with IgA Nephropathy in a Chinese Han Population. <i>DNA and Cell Biology</i> , 2019, 38, 1155-1165.	1.9	8
66	Plasma fibrinogen and mortality in patients undergoing peritoneal dialysis: a prospective cohort study. <i>BMC Nephrology</i> , 2020, 21, 349.	1.8	8
67	Association between monocyte count to high-density lipoprotein cholesterol ratio and mortality in patients undergoing peritoneal dialysis. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2021, 31, 2081-2088.	2.6	8
68	Abnormal iron status is associated with an increased risk of mortality in patients on peritoneal dialysis. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2021, 31, 1148-1155.	2.6	7
69	High Peritoneal Transport Status Was Not Associated with Mortality in Peritoneal Dialysis Patients with Diabetes. <i>PLoS ONE</i> , 2014, 9, e110445.	2.5	7
70	Ten-year survival of patients treated with peritoneal dialysis: A prospective observational cohort study. <i>Peritoneal Dialysis International</i> , 2020, 40, 573-580.	2.3	6
71	Serum Hepcidin-25 and Risk of Mortality in Patients on Peritoneal Dialysis. <i>Frontiers in Medicine</i> , 2021, 8, 684548.	2.6	6
72	Association of left ventricular systolic dysfunction with mortality in incident peritoneal dialysis patients. <i>Nephrology</i> , 2018, 23, 927-932.	1.6	5

#	ARTICLE	IF	CITATIONS
73	ST6GAL1 polymorphisms influence susceptibility and progression of IgA nephropathy in a Chinese Han population. <i>Immunobiology</i> , 2020, 225, 151973.	1.9	5
74	Age Difference in the Association between Hyponatremia and Infection-Related Mortality in Peritoneal Dialysis Patients. <i>Blood Purification</i> , 2020, 49, 631-640.	1.8	5
75	Serum lipoprotein(a) and risk of mortality in patients on peritoneal dialysis. <i>Journal of Clinical Lipidology</i> , 2020, 14, 252-259.	1.5	5
76	Changes of antibiotic resistance over time among <i>Escherichia coli</i> peritonitis in Southern China. <i>Peritoneal Dialysis International</i> , 2022, 42, 218-222.	2.3	5
77	Acetylation of HMGB1 by JNK1 Signaling Promotes LPS-Induced Peritoneal Mesothelial Cells Apoptosis. <i>BioMed Research International</i> , 2018, 2018, 1-12.	1.9	4
78	The negative impact of depressive symptoms on patient and technique survival in peritoneal dialysis: a prospective cohort study. <i>International Urology and Nephrology</i> , 2020, 52, 2393-2401.	1.4	4
79	Non-high-density lipoprotein cholesterol and mortality among peritoneal dialysis patients. <i>Journal of Clinical Lipidology</i> , 2021, 15, 732-742.	1.5	4
80	Faster Transport Status and Mortality in Anuric Patients Undergoing Continuous Ambulatory Peritoneal Dialysis. <i>Blood Purification</i> , 2015, 40, 160-166.	1.8	3
81	Long-Term Clinical Outcomes of Lupus Nephritis Patients Undergoing Peritoneal Dialysis: A Matched, Case-Control Study. <i>Peritoneal Dialysis International</i> , 2019, 39, 570-573.	2.3	3
82	Association of body mass index and uncontrolled blood pressure with cardiovascular mortality in peritoneal dialysis patients. <i>Journal of Human Hypertension</i> , 2019, 33, 106-114.	2.2	3
83	Peritonitis Affects the Relationship Between Low-Density Lipoprotein Cholesterol and Cardiovascular Events in Peritoneal Dialysis Patients. <i>Canadian Journal of Cardiology</i> , 2020, 36, 92-99.	1.7	3
84	Roles of peritoneal clearance and residual kidney removal in control of uric acid in patients on peritoneal dialysis. <i>BMC Nephrology</i> , 2020, 21, 148.	1.8	3
85	Hepatitis B Virus Infection Rate and Distribution in Chinese Systemic Lupus Erythematosus Patients. <i>Medical Science Monitor</i> , 2015, 21, 1955-1959.	1.1	3
86	Early initiation of PD therapy in elderly patients is associated with increased risk of death. <i>CKJ: Clinical Kidney Journal</i> , 2021, 14, 1649-1656.	2.9	3
87	Association between serum chloride levels with mortality in incident peritoneal dialysis patients. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2022, 32, 624-631.	2.6	3
88	Maintained Folic Acid Supplementation Reduces the Risk of Mortality in Continuous Ambulatory Peritoneal Dialysis Patients. <i>Blood Purification</i> , 2018, 45, 28-35.	1.8	2
89	Gender impact on baseline peritoneal transport properties in incident peritoneal dialysis patients. <i>International Urology and Nephrology</i> , 2019, 51, 2055-2061.	1.4	2
90	Higher serum phosphorus predicts residual renal function loss in male but not female incident peritoneal dialysis patients. <i>Journal of Nephrology</i> , 2020, 33, 829-837.	2.0	2

#	ARTICLE	IF	CITATIONS
91	Higher Eosinophils Predict Death-Censored Technique Failure in Peritoneal Dialysis Patients. <i>International Archives of Allergy and Immunology</i> , 2020, 181, 765-773.	2.1	2
92	Association of Abnormal Iron Status with the Occurrence and Prognosis of Peritoneal Dialysis-Related Peritonitis: A Longitudinal Data-Based 10-Year Retrospective Study. <i>Nutrients</i> , 2022, 14, 1613.	4.1	2
93	Incidence and Risk Factors Associated with Technique Failure in the First Year of Peritoneal Dialysis: A Single Center Retrospective Cohort Study in Southern China. <i>BMC Nephrology</i> , 2022, 23, .	1.8	2
94	Serum Sodium Modifies the Association of Systolic Blood Pressure with Mortality in Peritoneal Dialysis Patients. <i>Kidney and Blood Pressure Research</i> , 2020, 45, 916-925.	2.0	1
95	Rationale and design for Lowering-hyperUricemia treatment on cardiovascular outcomes in peritoneal dialysis patients: a prospective, multicentre, double-blind, randomised controlled trial (LUMINA). <i>BMJ Open</i> , 2020, 10, e037842.	1.9	1
96	Risk factors and clinical outcomes of encapsulating peritoneal sclerosis: A case-control study from China. <i>Peritoneal Dialysis International</i> , 2021, , 089686082110292.	2.3	1