## **Adoracion Cabrera**

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/4125187/publications.pdf

Version: 2024-02-01

20 280 11 17 papers citations h-index g-index

times ranked

citing authors

docs citations

all docs

#	Article	IF	CITATIONS
1	Development and Characterization of Wheat-Agropyron cristatum Introgression Lines Induced by Gametocidal Genes and Wheat ph1b Mutant. Agronomy, 2021, 11, 277.	3.0	7
2	Analysis of Chromosome Associations during Early Meiosis in Wheat Lines Carrying Chromosome Introgressions from Agropyron cristatum. Plants, 2021, 10, 2292.	3 <b>.</b> 5	O
3	Cloning and characterization of a putative orthologue of the wheat vernalization (VRN1) gene in perennial wheatgrass (Agropyron cristatum). Plant Breeding, 2020, 139, 1290-1298.	1.9	4
4	Uncovering homeologous relationships between tetraploid Agropyron cristatum and bread wheat genomes using COS markers. Theoretical and Applied Genetics, 2019, 132, 2881-2898.	3.6	12
5	Development of wheat—Hordeum chilense Chromosome 2Hch Introgression Lines Potentially Useful for Improving Grain Quality Traits. Agronomy, 2019, 9, 493.	3.0	5
6	Chromosomal location and molecular characterization of three grain hardness genes in Agropyron cristatum. Euphytica, 2019, 215, 1.	1.2	4
7	Wx Gene in Hordeum chilense: Chromosomal Location and Characterisation of the Allelic Variation in the Two Main Ecotypes of the Species. Agronomy, 2019, 9, 261.	3.0	10
8	Characterization of a world collection of Agropyron cristatum accessions. Genetic Resources and Crop Evolution, 2018, 65, 1455-1469.	1.6	9
9	Characterization of a set of common wheat– <i>Hordeum chilense</i> chromosome 7H <sup>ch</sup> introgression lines and its potential use in research on grain quality traits. Plant Breeding, 2017, 136, 344-350.	1.9	6
10	Chromosomal location of genes for resistance to powdery mildew in Agropyron cristatum and mapping of conserved orthologous set molecular markers. Euphytica, 2017, 213, 1.	1.2	14
11	Physical mapping of 5S and 45S rDNA genes and ploidy levels of Iranian Asparagus species. Scientia Horticulturae, 2016, 211, 269-276.	3.6	16
12	Molecular and cytogenetic characterization of a common wheat-Agropyron cristatum chromosome translocation conferring resistance to leaf rust. Euphytica, 2015, 201, 89-95.	1.2	35
13	Cytological and molecular characterization of wheat-Hordeum chilense chromosome 7Hch introgression lines. Euphytica, 2015, 203, 165-176.	1.2	12
14	The subtelomeric region is important for chromosome recognition and pairing during meiosis. Scientific Reports, 2014, 4, 6488.	3.3	39
15	Development and characterisation of structural changes in chromosome 3Hch from Hordeum chilense in common wheat and their use in physical mapping. Euphytica, 2012, 188, 429-440.	1.2	16
16	Sub-arm location of prolamin and EST-SSR loci on chromosome 1Hch from Hordeum chilense. Euphytica, 2011, 178, 63-69.	1.2	14
17	A physical map of chromosome 4Hch from H. chilense containing SSR, STS and EST-SSR molecular markers. Euphytica, 2009, 167, 253-259.	1.2	23
18	Genomic Constitution and Expression of Disease Resistance in Agropyron cristatum * Durum Wheat Derivatives. Breeding Science, 2007, 57, 17-21.	1.9	9

## ADORACION CABRERA

#	Article	IF	CITATION
19	A Fertile Amphiploid between Durum Wheat (Triticum Turgidurn) and the ×Agroticum Amphiploid (Agropyron cristatum × T. Tauschii). Hereditas, 2004, 135, 183-186.	1.4	15
20	Physical mapping of ribosomal DNA on several species of the subgenus Rosa. Theoretical and Applied Genetics, 2001, 103, 835-838.	3.6	30