

# Molina Molina Jm

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4146862/publications.pdf>

Version: 2024-02-01

40  
papers

2,098  
citations

257450

24  
h-index

302126

39  
g-index

40  
all docs

40  
docs citations

40  
times ranked

2897  
citing authors

#	ARTICLE	IF	CITATIONS
1	Assessment of chemical mixtures using biomarkers of combined biological activity: A screening study in human placentas. <i>Reproductive Toxicology</i> , 2021, 100, 143-154.	2.9	9
2	Bisphenol A and cognitive function in school-age boys: Is BPA predominantly related to behavior?. <i>NeuroToxicology</i> , 2019, 74, 162-171.	3.0	19
3	Concentrations of bisphenol A and parabens in socks for infants and young children in Spain and their hormone-like activities. <i>Environment International</i> , 2019, 127, 592-600.	10.0	51
4	Presence of Bisphenol A and Parabens in a Neonatal Intensive Care Unit: An Exploratory Study of Potential Sources of Exposure. <i>Environmental Health Perspectives</i> , 2019, 127, 117004.	6.0	32
5	Determination of bisphenol A and bisphenol S concentrations and assessment of estrogen- and anti-androgen-like activities in thermal paper receipts from Brazil, France, and Spain. <i>Environmental Research</i> , 2019, 170, 406-415.	7.5	59
6	Changes in the total effective xenoestrogen burden (TEXB) of breast cancer patients during an 18-month post-surgical follow-up. <i>Reproductive Toxicology</i> , 2017, 69, 212-220.	2.9	4
7	Ecotoxicological assessment of soils polluted with chemical waste from lindane production: Use of bacterial communities and earthworms as bioremediation tools. <i>Ecotoxicology and Environmental Safety</i> , 2017, 145, 539-548.	6.0	24
8	Total Effective Xenoestrogen Burden in Serum Samples and Risk for Breast Cancer in a Population-Based Multicase-Control Study in Spain. <i>Environmental Health Perspectives</i> , 2016, 124, 1575-1582.	6.0	41
9	Bisphenol A and other phenols in human placenta from children with cryptorchidism or hypospadias. <i>Reproductive Toxicology</i> , 2016, 59, 89-95.	2.9	79
10	A novel biomarker for anti-androgenic activity in placenta reveals risks of urogenital malformations. <i>Reproduction</i> , 2015, 149, 605-613.	2.6	13
11	Assessment of hormone-like activities in <i>Ginkgo biloba</i> , <i>Elettaria cardamomum</i> and <i>Plantago ovata</i> extracts using <i>in vitro</i> receptor-specific bioassays. <i>Food Additives and Contaminants - Part A Chemistry, Analysis, Control, Exposure and Risk Assessment</i> , 2015, 32, 1531-1541.	2.3	4
12	Screening of hormone-like activities in bottled waters available in Southern Spain using receptor-specific bioassays. <i>Environment International</i> , 2015, 74, 125-135.	10.0	21
13	Assessment of estrogenic and anti-androgenic activities of the mycotoxin zearalenone and its metabolites using <i>in vitro</i> receptor-specific bioassays. <i>Food and Chemical Toxicology</i> , 2014, 74, 233-239.	3.6	73
14	Adipose tissue concentrations of persistent organic pollutants and total cancer risk in an adult cohort from Southern Spain: Preliminary data from year 9 of the follow-up. <i>Science of the Total Environment</i> , 2014, 500-501, 243-249.	8.0	32
15	Validation of a UHPLC-MS/MS method for quantification of zearalenone, $\hat{1}\pm$ -zearalenol, $\hat{1}^2$ -zearalenol, $\hat{1}\pm$ -zearalanol, $\hat{1}^2$ -zearalanol and zearalanone in human urine. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2014, 962, 68-74.	2.3	37
16	<i>In vitro</i> study on the agonistic and antagonistic activities of bisphenol-S and other bisphenol-A congeners and derivatives via nuclear receptors. <i>Toxicology and Applied Pharmacology</i> , 2013, 272, 127-136.	2.8	305
17	Evaluation of estrogenic, antiestrogenic and genotoxic activity of nemorosone, the major compound found in brown Cuban propolis. <i>BMC Complementary and Alternative Medicine</i> , 2013, 13, 201.	3.7	15
18	Simultaneous determination of the UV-filters benzyl salicylate, phenyl salicylate, octyl salicylate, homosalate, 3-(4-methylbenzylidene) camphor and 3-benzylidene camphor in human placental tissue by LC-MS/MS. Assessment of their <i>in vitro</i> endocrine activity. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2013, 936, 80-87.	2.3	51

#	ARTICLE	IF	CITATIONS
19	Male specific association between xenoestrogen levels in placenta and birthweight. <i>Environment International</i> , 2013, 51, 174-181.	10.0	28
20	Adipose tissue concentrations of persistent organic pollutants and prevalence of type 2 diabetes in adults from Southern Spain. <i>Environmental Research</i> , 2013, 122, 31-37.	7.5	84
21	Predictors of the total effective xenoestrogen burden (TEXB) in human adipose tissue. A pilot study. <i>Reproductive Toxicology</i> , 2012, 33, 45-52.	2.9	16
22	Differential Estrogenic Effects of the Persistent Organochlorine Pesticides Dieldrin, Endosulfan, and Lindane in Primary Neuronal Cultures. <i>Toxicological Sciences</i> , 2011, 120, 413-427.	3.1	83
23	Antitumoral, mutagenic and (anti)estrogenic activities of tingenone and pristimerin. <i>Revista Brasileira De Farmacognosia</i> , 2011, 21, 963-971.	1.4	18
24	Prenatal exposure to organochlorine pesticides and TSH status in newborns from Southern Spain. <i>Science of the Total Environment</i> , 2011, 409, 3281-3287.	8.0	49
25	Relationship between occupational social class and exposure to organochlorine pesticides during pregnancy. <i>Chemosphere</i> , 2011, 83, 831-838.	8.2	11
26	Assessment of PCDD/F, PCB, OCP and BPA dietary exposure of non-breast-fed European infants. <i>Food Additives and Contaminants - Part A Chemistry, Analysis, Control, Exposure and Risk Assessment</i> , 2011, 28, 1110-1122.	2.3	27
27	Assessment of the total effective xenoestrogen burden in extracts of human placentas. <i>Biomarkers</i> , 2009, 14, 271-277.	1.9	27
28	The total effective xenoestrogen burden, a biomarker of exposure to xenoestrogen mixtures, is predicted by the (anti)estrogenicity of its components. <i>Reproductive Toxicology</i> , 2008, 26, 8-12.	2.9	17
29	Profiling of benzophenone derivatives using fish and human estrogen receptor-specific in vitro bioassays. <i>Toxicology and Applied Pharmacology</i> , 2008, 232, 384-395.	2.8	127
30	Polychlorinated biphenyls (PCBs) and hydroxy-PCBs in adipose tissue of women in Southeast Spain. <i>Chemosphere</i> , 2008, 71, 1196-1205.	8.2	66
31	Dioxins in adipose tissue of women in Southern Spain. <i>Chemosphere</i> , 2008, 73, 967-971.	8.2	17
32	Alkylphenols and bisphenol-A and its chlorinated derivatives in adipose tissue of children. , 2008, , .		7
33	Oestrogenicity of paper and cardboard extracts used as food containers. <i>Food Additives and Contaminants</i> , 2007, 24, 95-102.	2.0	69
34	PBDEs and PBBs in the adipose tissue of women from Spain. <i>Chemosphere</i> , 2007, 66, 377-383.	8.2	81
35	Human Exposure to Endocrine-Disrupting Chemicals and Prenatal Risk Factors for Cryptorchidism and Hypospadias: A Nested Caseá€“Control Study. <i>Environmental Health Perspectives</i> , 2007, 115, 8-14.	6.0	215
36	Biomonitoring of environmental estrogens in human tissues. <i>International Journal of Hygiene and Environmental Health</i> , 2007, 210, 429-432.	4.3	13

#	ARTICLE	IF	CITATIONS
37	Lack of activity of cadmium in in vitro estrogenicity assays. <i>Toxicology and Applied Pharmacology</i> , 2006, 216, 20-28.	2.8	66
38	Steroid receptor profiling of vinclozolin and its primary metabolites. <i>Toxicology and Applied Pharmacology</i> , 2006, 216, 44-54.	2.8	106
39	Assessment of total effective xenoestrogen burden in adipose tissue and identification of chemicals responsible for the combined estrogenic effect. <i>Analytical and Bioanalytical Chemistry</i> , 2004, 379, 163-170.	3.7	84
40	<i>Cryptosporidium parvum</i> : oocysts purification using potassium bromide discontinuous gradient. <i>Veterinary Parasitology</i> , 2000, 92, 223-226.	1.8	18