## Anna Enblom

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/4152286/publications.pdf

Version: 2024-02-01

933447 794594 21 373 10 19 citations h-index g-index papers 21 21 21 371 all docs docs citations times ranked citing authors

| #  | Article   | IF  | CITATIONS |
|----|---|-----|-----------|
| 1  | Identifying female pelvic cancer survivors withÂlow levels ofÂphysical activity after radiotherapy:<br>women withÂfecal andÂurinary leakage need additional support. Supportive Care in Cancer, 2020, 28,<br>2669-2681.   | 2.2 | 7         |
| 2  | Treatment expectations seem to affect bowel health when using acupuncture during radiotherapy for cancer: Secondary outcomes from a clinical randomized sham-controlled trial. Complementary Therapies in Medicine, 2020, 52, 102404.                                   | 2.7 | 3         |
| 3  | Frequent Stools Were Related to Reduced Quality of Life and Capacity in Daily Activities. Cancer Nursing, 2020, 43, 478-488.  | 1.5 | 2         |
| 4  | Participation in water-exercising long-term after breast cancer surgery: Experiences of significant factors for continuing exercising as a part of cancer rehabilitation. European Journal of Cancer Care, 2018, 27, e12736.  | 1.5 | 11        |
| 5  | Understanding rationales for acupuncture treated individuals' beliefs in acupuncture effects, to be able to maximize therapeutic results: A qualitative analysis. Complementary Therapies in Medicine, 2018, 39, 101-108.   | 2.7 | 1         |
| 6  | Patients— and Physiotherapists— Belief in and Use of Acupuncture for Cancer-Related Symptoms.<br>Acupuncture in Medicine, 2017, 35, 251-258.  | 1.0 | 19        |
| 7  | Complementary and alternative medicine self-care strategies for nausea in patients undergoing abdominal or pelvic irradiation for cancer: A longitudinal observational study of implementation in routine care. Complementary Therapies in Medicine, 2017, 34, 141-148. | 2.7 | 5         |
| 8  | Emesis in patients receiving acupuncture, sham acupuncture or standard care during chemo-radiation: A randomized controlled study. Complementary Therapies in Medicine, 2017, 34, 16-25.  | 2.7 | 13        |
| 9  | Type and Frequency of Side Effects during Pc6 Acupuncture: Observations from Therapists and Patients Participating in Clinical Efficacy Trials of Acupuncture. Acupuncture in Medicine, 2017, 35, 421-429.  | 1.0 | 8         |
| 10 | Experiences of incontinence and pelvic floor muscle training after gynaecologic cancer treatment. Supportive Care in Cancer, 2017, 25, 157-166.   | 2.2 | 22        |
| 11 | Reduced Need for Rescue Antiemetics and Improved Capacity to Eat in Patients Receiving Acupuncture<br>Compared to Patients Receiving Sham Acupuncture or Standard Care during Radiotherapy.<br>Evidence-based Complementary and Alternative Medicine, 2017, 2017, 1-11. | 1.2 | 8         |
| 12 | The Effects of Positive or Neutral Communication during Acupuncture for Relaxing Effects: A Sham-Controlled Randomized Trial. Evidence-based Complementary and Alternative Medicine, 2016, 2016, 1-11.  | 1.2 | 2         |
| 13 | Health-related quality of life and musculoskeletal function in patients with musculoskeletal disorders: after compared to before short-term group-based aqua-exercises. European Journal of Physiotherapy, 2016, 18, 218-225.   | 1.3 | 7         |
| 14 | WATER EXERCISE COMPARED TO LAND EXERCISE OR STANDARD CARE IN FEMALE CANCER SURVIVORS WITH SECONDARY LYMPHEDEMA. Lymphology, 2015, 48, 64-79.  | 0.2 | 11        |
| 15 | Acupuncture compared with placebo acupuncture in radiotherapy-induced nausea—a randomized controlled study. Annals of Oncology, 2012, 23, 1353-1361.  | 1.2 | 74        |
| 16 | Getting the Grip on Nonspecific Treatment Effects: Emesis in Patients Randomized to Acupuncture or Sham Compared to Patients Receiving Standard Care. PLoS ONE, 2011, 6, e14766.  | 2.5 | 39        |
| 17 | The Nonpenetrating Telescopic Sham Needle May Blind Patients with Different Characteristics and Experiences When Treated by Several Therapists. Evidence-based Complementary and Alternative Medicine, 2011, 2011, 1-12.  | 1.2 | 12        |
| 18 | Pilot Testing of Methods for Evaluation of Acupuncture for Emesis during Radiotherapy: A Randomised Single Subject Experimental Design. Acupuncture in Medicine, 2011, 29, 94-102.  | 1.0 | 17        |

## Anna Enblom

| #  | Article  | lF  | CITATION |
|----|--|-----|----------|
| 19 | Emesis and gastrointestinal problems during radiotherapy: A comparison of performance of daily activities between patients experiencing nausea and patients free from nausea. European Journal of Oncology Nursing, 2010, 14, 359-366. | 2.1 | 8        |
| 20 | One third of patients with radiotherapy-induced nausea consider their antiemetic treatment insufficient. Supportive Care in Cancer, 2009, 17, 23-32.   | 2.2 | 71       |
| 21 | Can individuals identify if needling was performed with an acupuncture needle or a non-penetrating sham needle?. Complementary Therapies in Medicine, 2008, 16, 288-294.   | 2.7 | 33       |