

# Morten T Kristensen

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/415593/publications.pdf>

Version: 2024-02-01

95  
papers

3,142  
citations

147801

31  
h-index

175258

52  
g-index

103  
all docs

103  
docs citations

103  
times ranked

2626  
citing authors

| #  | ARTICLE  | IF  | CITATIONS |
|----|--|-----|-----------|
| 1  | Anaemia impedes functional mobility after hip fracture surgery. <i>Age and Ageing</i> , 2008, 37, 173-178.   | 1.6 | 193       |
| 2  | Prediction of postoperative morbidity, mortality and rehabilitation in hip fracture patients: the cumulated ambulation score. <i>Clinical Rehabilitation</i> , 2006, 20, 701-708.                                | 2.2 | 156       |
| 3  | The effects of liberal versus restrictive transfusion thresholds on ambulation after hip fracture surgery. <i>Transfusion</i> , 2009, 49, 227-234.   | 1.6 | 156       |
| 4  | Loss of Knee-Extension Strength Is Related to Knee Swelling After Total Knee Arthroplasty. <i>Archives of Physical Medicine and Rehabilitation</i> , 2010, 91, 1770-1776.  | 0.9 | 139       |
| 5  | Timed "Up & Go" Test as a Predictor of Falls Within 6 Months After Hip Fracture Surgery. <i>Physical Therapy</i> , 2007, 87, 24-30.  | 2.4 | 137       |
| 6  | Effect of Postoperative Epidural Analgesia on Rehabilitation and Pain after Hip Fracture Surgery. <i>Anesthesiology</i> , 2005, 102, 1197-1204.  | 2.5 | 128       |
| 7  | Prefracture functional level evaluated by the New Mobility Score predicts in-hospital outcome after hip fracture surgery. <i>Monthly Notices of the Royal Astronomical Society: Letters</i> , 2010, 81, 296-302. | 3.3 | 128       |
| 8  | Factors affecting functional prognosis of patients with hip fracture. <i>European Journal of Physical and Rehabilitation Medicine</i> , 2011, 47, 257-64.  | 2.2 | 107       |
| 9  | High inter-tester reliability of the new mobility score in patients with hip fracture. <i>Journal of Rehabilitation Medicine</i> , 2008, 40, 589-591.  | 1.1 | 86        |
| 10 | Very low survival rates after non-traumatic lower limb amputation in a consecutive series: what to do?. <i>Interactive Cardiovascular and Thoracic Surgery</i> , 2012, 14, 543-547.                              | 1.1 | 84        |
| 11 | Postoperative pain after hip fracture is procedure specific. <i>British Journal of Anaesthesia</i> , 2009, 102, 111-116.   | 3.4 | 81        |
| 12 | The role of pain for early rehabilitation in fast track total knee arthroplasty. <i>Disability and Rehabilitation</i> , 2010, 32, 300-306.   | 1.8 | 70        |
| 13 | Physical Therapy Management of Older Adults With Hip Fracture. <i>Journal of Orthopaedic and Sports Physical Therapy</i> , 2021, 51, CPG1-CPG81.   | 3.5 | 70        |
| 14 | High intertester reliability of the Cumulated Ambulation Score for the evaluation of basic mobility in patients with hip fracture. <i>Clinical Rehabilitation</i> , 2009, 23, 1116-1123.                         | 2.2 | 56        |
| 15 | Hip fractures in the non-elderly "Who, why and whither?. <i>Injury</i> , 2018, 49, 1445-1450.  | 1.7 | 52        |
| 16 | Knee-extension strength, postural control and function are related to fracture type and thigh edema in patients with hip fracture. <i>Clinical Biomechanics</i> , 2009, 24, 218-224.                             | 1.2 | 50        |
| 17 | Change of residence and functional status within three months and one year following hip fracture surgery. <i>Disability and Rehabilitation</i> , 2014, 36, 685-690.   | 1.8 | 49        |
| 18 | Predictors of Long-Term Mortality in Older People With Hip Fracture. <i>Archives of Physical Medicine and Rehabilitation</i> , 2015, 96, 1215-1221.  | 0.9 | 48        |

| #  | ARTICLE  | IF  | CITATIONS |
|----|--|-----|-----------|
| 19 | Non-Weight-Bearing Status Compromises the Functional Level Up to 1 yr After Hip Fracture Surgery. <i>American Journal of Physical Medicine and Rehabilitation</i> , 2014, 93, 641-648.   | 1.4 | 46        |
| 20 | Physiotherapy following fragility fractures. <i>Injury</i> , 2018, 49, 1413-1417.  | 1.7 | 45        |
| 21 | The basic mobility status upon acute hospital discharge is an independent risk factor for mortality up to 5 years after hip fracture surgery. <i>Monthly Notices of the Royal Astronomical Society: Letters</i> , 2018, 89, 47-52.   | 3.3 | 44        |
| 22 | High perceived caregiver burden for relatives of patients following hip fracture surgery. <i>Disability and Rehabilitation</i> , 2019, 41, 311-318.  | 1.8 | 44        |
| 23 | Fatigue and pain limit independent mobility and physiotherapy after hip fracture surgery. <i>Disability and Rehabilitation</i> , 2018, 40, 1808-1816.  | 1.8 | 43        |
| 24 | Factors with independent influence on the "timed up and go"™ test in patients with hip fracture. <i>Physiotherapy Research International</i> , 2009, 14, 30-41.  | 1.5 | 41        |
| 25 | Effect of a single preoperative 125 mg dose of methylprednisolone on postoperative delirium in hip fracture patients; a randomised, double-blind, placebo-controlled trial. <i>Anaesthesia</i> , 2018, 73, 1353-1360.  | 3.8 | 41        |
| 26 | Most patients regain prefracture basic mobility after hip fracture surgery in a fast-track programme. <i>Danish Medical Journal</i> , 2012, 59, A4447.   | 0.5 | 40        |
| 27 | Efficacy of early controlled motion of the ankle compared with immobilisation in non-operative treatment of patients with an acute Achilles tendon rupture: an assessor-blinded, randomised controlled trial. <i>British Journal of Sports Medicine</i> , 2020, 54, 719-724. | 6.7 | 37        |
| 28 | Timed Up & Go Test Score in Patients With Hip Fracture Is Related to the Type of Walking Aid. <i>Archives of Physical Medicine and Rehabilitation</i> , 2009, 90, 1760-1765.   | 0.9 | 36        |
| 29 | How Many Trials Are Needed to Achieve Performance Stability of the Timed Up & Go Test in Patients With Hip Fracture?. <i>Archives of Physical Medicine and Rehabilitation</i> , 2010, 91, 885-889.   | 0.9 | 36        |
| 30 | Hip Fracture-Related Pain Strongly Influences Functional Performance of Patients With an Intertrochanteric Fracture Upon Discharge From the Hospital. <i>PM and R</i> , 2013, 5, 135-141.  | 1.6 | 36        |
| 31 | Regaining pre-fracture basic mobility status after hip fracture and association with post-discharge mortality and readmission—a nationwide register study in Denmark. <i>Age and Ageing</i> , 2019, 48, 278-284.   | 1.6 | 36        |
| 32 | Feasibility of Progressive Strength Training Implemented in the Acute Ward after Hip Fracture Surgery. <i>PLoS ONE</i> , 2014, 9, e93332.  | 2.5 | 33        |
| 33 | Physical Activity in the Acute Ward Following Hip Fracture Surgery is Associated with Less Fear of Falling. <i>Journal of Aging and Physical Activity</i> , 2016, 24, 525-532.   | 1.0 | 33        |
| 34 | Effectiveness of acute in-hospital physiotherapy with knee-extension strength training in reducing strength deficits in patients with a hip fracture: A randomised controlled trial. <i>PLoS ONE</i> , 2017, 12, e0179867.   | 2.5 | 31        |
| 35 | Cumulated Ambulation Score to evaluate mobility is feasible in geriatric patients and in patients with hip fracture. <i>Danish Medical Journal</i> , 2012, 59, A4464.  | 0.5 | 30        |
| 36 | Standardised test protocol (Constant Score) for evaluation of functionality in patients with shoulder disorders. <i>Danish Medical Journal</i> , 2013, 60, A4608.  | 0.5 | 30        |

| #  | ARTICLE  | IF  | CITATIONS |
|----|--|-----|-----------|
| 37 | RELATIVE AND ABSOLUTE INTERTESTER RELIABILITY OF THE TIMED UP AND GO TEST TO QUANTIFY FUNCTIONAL MOBILITY IN PATIENTS WITH HIP FRACTURE. <i>Journal of the American Geriatrics Society</i> , 2011, 59, 565-567.  | 2.6 | 29        |
| 38 | Introducing a Third Timed Up & Go Test Trial Improves Performances of Hospitalized and Community-Dwelling Older Individuals. <i>Journal of Geriatric Physical Therapy</i> , 2017, 40, 121-126.   | 1.1 | 29        |
| 39 | Feasibility of progressive strength training shortly after hip fracture surgery. <i>World Journal of Orthopedics</i> , 2013, 4, 248.   | 1.8 | 29        |
| 40 | Interrater reliability of the standardized Timed Up and Go Test when used in hospitalized and community-dwelling older individuals. <i>Physiotherapy Research International</i> , 2019, 24, e1769.   | 1.5 | 28        |
| 41 | The Barthel Index and the Cumulated Ambulation Score are superior to the de Morton Mobility Index for the early assessment of outcome in patients with a hip fracture admitted to an acute geriatric ward. <i>Disability and Rehabilitation</i> , 2019, 41, 1351-1359. | 1.8 | 26        |
| 42 | Thigh and Knee Circumference, Knee Extension Strength, and Functional Performance After Fast Track Total Hip Arthroplasty. <i>PM and R</i> , 2011, 3, 117-124.   | 1.6 | 25        |
| 43 | Physical performance following acute high-risk abdominal surgery: a prospective cohort study. <i>Canadian Journal of Surgery</i> , 2018, 61, 42-49.  | 1.2 | 23        |
| 44 | Development and reliability of the Achilles Tendon Length Measure and comparison with the Achilles Tendon Resting Angle on patients with an Achilles tendon rupture. <i>Foot and Ankle Surgery</i> , 2017, 23, 275-280.  | 1.7 | 22        |
| 45 | Number of test trials needed for performance stability and interrater reliability of the one leg stand test in patients with a major non-traumatic lower limb amputation. <i>Gait and Posture</i> , 2014, 39, 424-429.   | 1.4 | 21        |
| 46 | Simultaneous Disulfide and Boronic Acid Ester Exchange in Dynamic Combinatorial Libraries. <i>International Journal of Molecular Sciences</i> , 2015, 16, 21858-21872.   | 4.1 | 21        |
| 47 | Surgical apgar score predicts early complication in transfemoral amputees: Retrospective study of 170 major amputations. <i>World Journal of Orthopedics</i> , 2016, 7, 832.   | 1.8 | 19        |
| 48 | Intra- and Interrater Reliability and Agreement of the Danish Version of the Dynamic Gait Index in Older People With Balance Impairments. <i>Archives of Physical Medicine and Rehabilitation</i> , 2011, 92, 1630-1635.   | 0.9 | 18        |
| 49 | High inter-rater reliability, agreement, and convergent validity of Constant score in patients with clavicle fractures. <i>Journal of Shoulder and Elbow Surgery</i> , 2016, 25, 1577-1582.  | 2.6 | 14        |
| 50 | Psychometric properties of the Cumulated Ambulation Score: a systematic review. <i>European Journal of Physical and Rehabilitation Medicine</i> , 2018, 54, 766-771.   | 2.2 | 14        |
| 51 | An exercise program for people with severe peripheral neuropathy and diabetic foot ulcers – a case series on feasibility and safety. <i>Disability and Rehabilitation</i> , 2020, 42, 183-189.   | 1.8 | 14        |
| 52 | COVID-19 and acute respiratory failure treated with CPAP. <i>European Clinical Respiratory Journal</i> , 2021, 8, 1910191.   | 1.5 | 14        |
| 53 | Cross-cultural validation of the Italian version of the Cumulated Ambulation Score. <i>International Journal of Rehabilitation Research</i> , 2016, 39, 160-164.   | 1.3 | 13        |
| 54 | An enhanced treatment program with markedly reduced mortality after a transtibial or higher non-traumatic lower extremity amputation. <i>Monthly Notices of the Royal Astronomical Society: Letters</i> , 2016, 87, 306-311.   | 3.3 | 13        |

| #  | ARTICLE  | IF  | CITATIONS |
|----|--|-----|-----------|
| 55 | Interrater Reliability of the 6-Minute Walk Test in Women With Hip Fracture. <i>Journal of Geriatric Physical Therapy</i> , 2017, 40, 158-166.   | 1.1 | 13        |
| 56 | Neither operative nor nonoperative approach is superior for treating displaced midshaft clavicle fractures: a partially blinded randomized controlled clinical trial. <i>Bone and Joint Journal</i> , 2021, 103-B, 762-768.  | 4.4 | 13        |
| 57 | Translation, inter-rater reliability, agreement, and internal consistency of the Spanish version of the cumulated ambulation score in patients after hip fracture. <i>Disability and Rehabilitation</i> , 2020, 42, 2766-2771.   | 1.8 | 12        |
| 58 | Tourniquets do not increase the total blood loss or re-amputation risk in transtibial amputations. <i>World Journal of Orthopedics</i> , 2017, 8, 62.  | 1.8 | 12        |
| 59 | Reliability of the Copenhagen Achilles length measure (CALM) on patients with an Achilles tendon rupture. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> , 2020, 28, 281-290.   | 4.2 | 11        |
| 60 | Knee Extension Strength Measures Indicating Probable Sarcopenia Is Associated with Health-Related Outcomes and a Strong Predictor of 1-Year Mortality in Patients Following Hip Fracture Surgery. <i>Geriatrics (Switzerland)</i> , 2021, 6, 8.  | 1.7 | 11        |
| 61 | Early recovery of basic mobility under femoral nerve block after hip fracture surgery – A propensity score matched pilot study. <i>Injury</i> , 2021, 52, 3382-3387.   | 1.7 | 11        |
| 62 | Exercise Therapy Is Effective at Improving Short- and Long-Term Mobility, Activities of Daily Living, and Balance in Older Patients Following Hip Fracture: A Systematic Review and Meta-Analysis. <i>Journals of Gerontology - Series A Biological Sciences and Medical Sciences</i> , 2022, 77, 861-871. | 3.6 | 11        |
| 63 | Efficacy of early controlled motion of the ankle compared with no motion after non-operative treatment of an acute Achilles tendon rupture: study protocol for a randomized controlled trial. <i>Trials</i> , 2016, 17, 564.   | 1.6 | 10        |
| 64 | Performance Stability and Interrater Reliability of Culturally Adapted 10-Meter Walking Test for Danes with Neurological Disorders. <i>Journal of Stroke and Cerebrovascular Diseases</i> , 2019, 28, 2459-2467.   | 1.6 | 10        |
| 65 | Orthopaedic physiotherapists'™ perceptions of mechanisms for observed variation in the implementation of physiotherapy practices in the early postoperative phase after hip fracture: a UK qualitative study. <i>Age and Ageing</i> , 2021, 50, 1961-1970.   | 1.6 | 10        |
| 66 | Quality indicators in the treatment of geriatric hip fractures: literature review and expert consensus. <i>Archives of Osteoporosis</i> , 2021, 16, 152.   | 2.4 | 10        |
| 67 | Frailty and osteoporosis in patients with hip fractures under the age of 60 – a prospective cohort of 218 individuals. <i>Osteoporosis International</i> , 2022, 33, 1037-1055.  | 3.1 | 10        |
| 68 | Municipality-based physical rehabilitation after acute hip fracture surgery in Denmark. <i>Danish Medical Journal</i> , 2015, 62, A5023.   | 0.5 | 10        |
| 69 | Effect of Automated Oxygen Titration during Walking on Dyspnea and Endurance in Chronic Hypoxemic Patients with COPD: A Randomized Crossover Trial. <i>Journal of Clinical Medicine</i> , 2021, 10, 4820.  | 2.4 | 9         |
| 70 | Resistance Training With Ankle Weight Cuffs Is Feasible in Patients With Acute Exacerbation of COPD. <i>Journal of Cardiopulmonary Rehabilitation and Prevention</i> , 2017, 37, 49-56.  | 2.1 | 8         |
| 71 | Individualized treatment for acute Achilles tendon rupture based on the Copenhagen Achilles Rupture Treatment Algorithm (CARTA): a study protocol for a multicenter randomized controlled trial. <i>Trials</i> , 2020, 21, 399.  | 1.6 | 8         |
| 72 | Validation of Two Activity Monitors in Slow and Fast Walking Hospitalized Patients. <i>Rehabilitation Research and Practice</i> , 2022, 2022, 1-14.  | 0.6 | 8         |

| #  | ARTICLE   | IF  | CITATIONS |
|----|---|-----|-----------|
| 73 | Reliability and agreement between 2 strength devices used in the newly modified and standardized Constant score. <i>Journal of Shoulder and Elbow Surgery</i> , 2014, 23, 1806-1812.  | 2.6 | 7         |
| 74 | Development and psychometric properties of the Basic Amputee Mobility Score for use in patients with a major lower extremity amputation. <i>Geriatrics and Gerontology International</i> , 2018, 18, 138-145.   | 1.5 | 7         |
| 75 | Early mobilisation of patients with community-acquired pneumonia reduce length of hospitalisation—a pilot study. <i>Journal of Physical Therapy Science</i> , 2018, 30, 926-932.  | 0.6 | 7         |
| 76 | Factors influencing performances and indicating risk of falls using the true Timed Up and Go test time of patients with hip fracture upon acute hospital discharge. <i>Physiotherapy Research International</i> , 2020, 25, e1841.  | 1.5 | 7         |
| 77 | Total Blood Loss After Transfemoral Amputations Is Twice the Intraoperative Loss: An Observational Cohort Study of 81 Nontraumatic Amputations. <i>Geriatric Orthopaedic Surgery and Rehabilitation</i> , 2017, 8, 123-127.   | 1.4 | 6         |
| 78 | The Achilles Tendon Length Measure and the Achilles Tendon Resting Angle show acceptable construct validity using the Copenhagen Achilles Length Measure as gold standard. <i>Foot and Ankle Surgery</i> , 2021, 27, 655-659.   | 1.7 | 6         |
| 79 | Translation, Inter-rater Reliability, Agreement, and Internal Consistency of the Japanese Version of the Cumulated Ambulation Score in Patients after Hip Fracture Surgery. <i>Progress in Rehabilitation Medicine</i> , 2020, 5, n/a.  | 0.9 | 6         |
| 80 | Translation, reliability, agreement and validity of the Turkish version of Cumulated Ambulation Score in patients with hip fracture. <i>Joint Diseases and Related Surgery</i> , 2020, 31, 346-352.   | 1.0 | 6         |
| 81 | The cumulated ambulation score is superior to the new mobility score and the de Morton Mobility Index in predicting discharge destination of patients admitted to an acute geriatric ward; a 1-year cohort study of 491 patients. <i>Disability and Rehabilitation</i> , 2022, 44, 1481-1488.                                       | 1.8 | 5         |
| 82 | Feasibility and preliminary effect of anabolic steroids in addition to strength training and nutritional supplement in rehabilitation of patients with hip fracture: a randomized controlled pilot trial (HIP-SAP1 trial). <i>BMC Geriatrics</i> , 2021, 21, 323.   | 2.7 | 5         |
| 83 | Loss of pre-fracture basic mobility status at hospital discharge for hip fracture is associated with 30-day post-discharge risk of infections - a four-year nationwide cohort study of 23,309 Danish patients. <i>Injury</i> , 2021, 52, 1833-1840.   | 1.7 | 5         |
| 84 | Avoidable 30-day mortality analysis and failure to rescue in dysvascular lower extremity amputees. <i>Monthly Notices of the Royal Astronomical Society: Letters</i> , 2018, 89, 246-250.   | 3.3 | 4         |
| 85 | Preliminary effect and feasibility of physiotherapy with strength training and protein-rich nutritional supplement in combination with anabolic steroids in cross-continuum rehabilitation of patients with hip fracture: protocol for a blinded randomized controlled pilot trial (HIP-SAP1 trial). <i>Trials</i> , 2019, 20, 763. | 1.6 | 4         |
| 86 | Six Versus 12 Weeks of Outpatient Physical Therapy Including Progressive Resistance Training in Cognitively Intact Older Adults After Hip Fracture: A Multicenter Randomized Controlled Trial. <i>Journals of Gerontology - Series A Biological Sciences and Medical Sciences</i> , 2022, 77, 1455-1462.                            | 3.6 | 3         |
| 87 | Translation, Reliability and Validity of the Spanish Version of the Modified New Mobility Score (NMS-ES). <i>International Journal of Environmental Research and Public Health</i> , 2021, 18, 723.   | 2.6 | 3         |
| 88 | The Copenhagen Achilles Length Measurement performed at time of acute Achilles tendon rupture is correlated to tendon elongation after one year.. <i>Journal of Foot and Ankle Surgery</i> , 2022, , .  | 1.0 | 3         |
| 89 | Validity, Responsiveness, and Predictive Ability of the Japanese Version of the Cumulated Ambulation Score in Patients with Hip Fracture. <i>Progress in Rehabilitation Medicine</i> , 2022, 7, n/a.  | 0.9 | 3         |
| 90 | Difficult to predict early failure after major lower-extremity amputations. <i>Danish Medical Journal</i> , 2015, 62, A5172.  | 0.5 | 3         |

| #  | ARTICLE   | IF  | CITATIONS |
|----|---|-----|-----------|
| 91 | 12â€¦Efficacy of early controlled motion of the ankle in non-operative treatment of acute achilles tendon rupture. An assessor-blinded RCT. , 2019, , .   |     | 2         |
| 92 | Mobilisation and physiotherapy intervention following hip fracture: snapshot survey across six countries from the Fragility Fracture Network Physiotherapy Group. Disability and Rehabilitation, 2022, 44, 6788-6795.         | 1.8 | 2         |
| 93 | 21â€¦Validation of achilles tendon length measure and achilles tendon resting angle in relation to copenhagen achilles length measure. , 2019, , .  |     | 0         |
| 94 | Older patientsâ€™ perspectives on participating in multimodal rehabilitation including anabolic steroids following hip fracture: a qualitative study embedded within a pilot RCT. Disability and Rehabilitation, 2022, , 1-9. | 1.8 | 0         |
| 95 | Motives and Barriers to Exercise Training during Hospitalization in Patients with Type 2 Diabetes: A Cross-Sectional Study. International Journal of Environmental Research and Public Health, 2022, 19, 1035.                | 2.6 | 0         |