Ricardo Reyes-Chilpa

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/4187284/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Antimycobacterial Activity of Alkaloids and Extracts from Tabernaemontana alba and T. arborea. Planta Medica, 2022, 88, 53-61.	1.3	7
2	Tabernaemontana arborea and ibogaine induce paroxysmal EEG activity in freely moving mice: Involvement of serotonin 5-HT1A receptors. NeuroToxicology, 2022, 89, 79-91.	3.0	7
3	Microsatellites, morphological, and alkaloids characterization of Zephyranthes fosteri and Z. alba (Amaryllidaceae): Allopatric populations. Biochemical Systematics and Ecology, 2022, 101, 104398.	1.3	3
4	On the first book of medicinal plants written in the American Continent:The LibellusMedicinalibus Indorum Herbisfrom Mexico, 1552. A review. Boletin Latinoamericano Y Del Caribe De Plantas Medicinales Y Aromaticas, 2021, 20, 1-27.	0.5	6
5	Plants of the †Libellus de Medicinalibus Indorum Herbis' from Mexico, 1552. Zephyranthes fosteri (Amaryllidaceae) Alkaloids. Chemistry and Biodiversity, 2021, 18, e2000834.	2.1	5
6	Phytotoxic activity of aqueous extracts of ruderal plants and its potential application to tomato crop. Botanical Sciences, 2021, 99, 487-498.	0.8	3
7	The New World Bays (Litsea, Lauraceae). A Botanical, Chemical, Pharmacological and Ecological Review in Relation to their Traditional and Potential Applications as Phytomedicines. Botanical Review, The, 2021, 87, 392-420.	3.9	3
8	Mammea type coumarins isolated from Calophyllum brasiliense induced apoptotic cell death of Trypanosoma cruzi through mitochondrial dysfunction, ROS production and cell cycle alterations. Bioorganic Chemistry, 2020, 100, 103894.	4.1	7
9	Quantitative Evaluation of a Mexican and a Chanaian <i>Tabernaemontana</i> Species as Alternatives to <i>Voacanga africana</i> for the Production of Antiaddictive Ibogan Type Alkaloids. Chemistry and Biodiversity, 2020, 17, e2000002.	2.1	4
10	Network Pharmacology Uncovers Anticancer Activity of Mammea-Type Coumarins from Calophyllum brasiliense. Planta Medica, 2019, 85, 14-23.	1.3	9
11	Extraction and Conversion Studies of the Antiaddictive Alkaloids Coronaridine, Ibogamine, Voacangine, and Ibogaine from Two Mexican <i>Tabernaemontana</i> Species (Apocynaceae). Chemistry and Biodiversity, 2019, 16, e1900175.	2.1	12
12	Strategies for the in vitro production of antiaddictive ibogan type alkaloids from Apocynaceae species. Plant Cell, Tissue and Organ Culture, 2019, 138, 215-227.	2.3	3
13	Trypanocidal and toxicological assessment in vitro and in silico of three sesquiterpene lactones from Asteraceae plant species. Food and Chemical Toxicology, 2019, 125, 55-61.	3.6	18
14	Coumarins isolated from Calophyllum brasiliense produce ultrastructural alterations and affect in vitro infectivity of Trypanosoma cruzi. Phytomedicine, 2019, 61, 152827.	5.3	15
15	Chemoinformatic Analysis of Selected Cacalolides from Psacalium decompositum (A. Gray) H. Rob. & Brettell and Psacalium peltatum (Kunth) Cass. and Their Effects on FcεRI-Dependent Degranulation in Mast Cells. Molecules, 2018, 23, 3367.	3.8	6
16	Antinociceptive, antiâ€inflammatory, and central nervous system (CNS) effects of the natural coumarin soulattrolide. Drug Development Research, 2018, 79, 332-338.	2.9	6
17	Broccoli sprouts produce abdominal antinociception but not spasmolytic effects like its bioactive metabolite sulforaphane. Biomedicine and Pharmacotherapy, 2018, 107, 1770-1778.	5.6	10
18	Jacareubin inhibits FcεRI-induced extracellular calcium entry and production of reactive oxygen species required for anaphylactic degranulation of mast cells. Biochemical Pharmacology, 2018, 154, 344-356.	4.4	7

#	Article	IF	CITATIONS
19	Bioactive Coumarins and Xanthones From Calophyllum Genus and Analysis of Their Druglikeness and Toxicological Properties. Studies in Natural Products Chemistry, 2017, 53, 277-307.	1.8	7
20	Cytogenetic effects of Jacareubin from Calophyllum brasiliense on human peripheral blood mononucleated cells in vitro and on mouse polychromatic erythrocytes in vivo. Toxicology and Applied Pharmacology, 2017, 335, 6-15.	2.8	9
21	Risk assessment of Soulatrolide and Mammea (A/BA+A/BB) coumarins from Calophyllum brasiliense by a toxicogenomic and toxicological approach. Food and Chemical Toxicology, 2016, 91, 117-129.	3.6	9
22	Antimycobacterial and HIV-1 Reverse Transcriptase Activity of Julianaceae and Clusiaceae Plant Species from Mexico. Evidence-based Complementary and Alternative Medicine, 2015, 2015, 1-8.	1.2	15
23	Toxicogenomic analysis of pharmacological active coumarins isolated from Calophyllum brasiliense. Genomics Data, 2015, 6, 258-259.	1.3	4
24	Linalool and β-pinene exert their antidepressant-like activity through the monoaminergic pathway. Life Sciences, 2015, 128, 24-29.	4.3	120
25	Trends in the chemical and pharmacological research on the tropical trees Calophyllum brasiliense and Calophyllum inophyllum, a global context. Scientometrics, 2015, 105, 1019-1030.	3.0	13
26	Anthraquinones from Vismia mexicana. Zeitschrift Fur Naturforschung - Section C Journal of Biosciences, 2014, 69, 29-34.	1.4	6
27	Molecular mechanisms involved in the cytotoxicity induced by coumarins from Calophyllum brasiliense in K562 leukaemia cells. Journal of Pharmacy and Pharmacology, 2014, 66, 1189-1195.	2.4	7
28	Rotenone isolated from Pachyrhizus erosus displays cytotoxicity and genotoxicity in K562 cells. Natural Product Research, 2014, 28, 1780-1785.	1.8	15
29	Neuropharmacological in vivo effects and phytochemical profile of the extract from the aerial parts of Heteropterys brachiata (L.) DC. (Malpighiaceae). Journal of Ethnopharmacology, 2013, 146, 311-317.	4.1	19
30	Antioxidant properties of xanthones from Calophyllum brasiliense: prevention of oxidative damage induced by FeSO4. BMC Complementary and Alternative Medicine, 2013, 13, 262.	3.7	21
31	Antidepressant activity of Litsea glaucescens essential oil: Identification of β-pinene and linalool as active principles. Journal of Ethnopharmacology, 2012, 143, 673-679.	4.1	124
32	Acetylcholinesterase-inhibiting Alkaloids from Zephyranthes concolor. Molecules, 2011, 16, 9520-9533.	3.8	23
33	Antidepressant effect and pharmacological evaluation of standardized extract of flavonoids from Byrsonima crassifolia. Phytomedicine, 2011, 18, 1255-1261.	5.3	50
34	Essential Oils in Mexican Bays (Litsea spp., Lauraceae): Taxonomic Assortment and Ethnobotanical Implications1. Economic Botany, 2011, 65, 178-189.	1.7	28
35	Leaf Structure of Two Chemotypes of Calophyllum brasiliense from Mexico. Microscopy and Microanalysis, 2011, 17, 340-341.	0.4	4
36	Production of anti-HIV-1 calanolides in a callus culture of Calophyllum brasiliense (Cambes). Plant Cell, Tissue and Organ Culture, 2010, 103, 33-40.	2.3	30

RICARDO REYES-CHILPA

#	Article	IF	CITATIONS
37	Coumarin A/AA induces apoptosisâ€like cell death in HeLa cells mediated by the release of apoptosisâ€inducing factor. Journal of Biochemical and Molecular Toxicology, 2009, 23, 263-272.	3.0	25
38	Sesquiterpenoids from antidiabetic Psacalium decompositum block ATP sensitive potassium channels. Journal of Ethnopharmacology, 2009, 123, 489-493.	4.1	7
39	X-ray and high-resolution ¹ H and ¹³ C NMR of smooth muscle relaxant sesquiterpene lactones. Canadian Journal of Chemistry, 2008, 86, 1077-1084.	1.1	7
40	Reactivity of Several Reactive Oxygen Species (ROS) with the Sesquiterpene Cacalol. Natural Product Communications, 2008, 3, 1934578X0800300.	0.5	1
41	Trypanocidal constituents in plants: 7. Mammea-type coumarins. Memorias Do Instituto Oswaldo Cruz, 2008, 103, 431-436.	1.6	41
42	Productos forestales no maderables en México: Aspectos económicos para el desarrollo sustentable. Madera Bosques, 2008, 14, 95-112.	0.2	21
43	Revised NMR data for Incartine: an Alkaloid from Galanthus elwesii. Molecules, 2007, 12, 1430-1435.	3.8	33
44	Inhibition of gastric H+,K+-ATPase activity by flavonoids, coumarins and xanthones isolated from Mexican medicinal plants. Journal of Ethnopharmacology, 2006, 105, 167-172.	4.1	73
45	Trypanocidal Constituents in Plants 5. Evaluation of Some Mexican Plants for Their Trypanocidal Activity and Active Constituents in the Seeds of Persea americana. Biological and Pharmaceutical Bulletin, 2005, 28, 1314-1317.	1.4	47
46	Antibacterial activity of crude extracts from Mexican medicinal plants and purified coumarins and xanthones. Journal of Ethnopharmacology, 2005, 97, 293-299.	4.1	152
47	Cytotoxic effects of mammea type coumarins from Calophyllum brasiliense. Life Sciences, 2004, 75, 1635-1647.	4.3	79
48	Trypanocidal Constituents in Plants 3. Leaves of Garcinia intermedia and Heartwood of Calophyllum brasiliense. Biological and Pharmaceutical Bulletin, 2004, 27, 141-143.	1.4	72
49	Methanol Extracts of Hamelia patens Containing Oxindole Alkaloids Relax KCl-Induced Contraction in Rat Myometrium. Biological and Pharmaceutical Bulletin, 2004, 27, 1617-1620.	1.4	20
50	HIV-1 Inhibition by Extracts of Clusiaceae Species from Mexico. Biological and Pharmaceutical Bulletin, 2004, 27, 916-920.	1.4	21
51	HIV-1 Inhibitory Compounds from Calophyllum brasiliense Leaves. Biological and Pharmaceutical Bulletin, 2004, 27, 1471-1475.	1.4	75
52	Seasonal evaluation of the postharvest fungicidal activity of powders and extracts of huamuchil (Pithecellobium dulce): action against Botrytris cinerea, Penicillium digitatum and Rhizopus stolonifer of strawberry fruit. Postharvest Biology and Technology, 2003, 29, 81-92.	6.0	26
53	Relaxation of Uterine and Aortic Smooth Muscle by Claucolides D and E from Vernonia liatroides Biological and Pharmaceutical Bulletin, 2003, 26, 112-115.	1.4	13
54	A Novel Cacalolide from Psacalium Decompositum. Natural Product Research, 2002, 16, 239-242.	0.4	3

#	Article	IF	CITATIONS
55	Trypanocidal Constituents in Plants. 1. Evaluation of Some Mexican Plants for Their Trypanocidal Activity and Active Constituents in Guaco, Roots of Aristolochia taliscana Biological and Pharmaceutical Bulletin, 2002, 25, 1188-1191.	1.4	59
56	Natural products isolated from Mexican medicinal plants: Novel inhibitors of sulfotransferases, SULT1A1 and SULT2A1. Phytomedicine, 2001, 8, 481-488.	5.3	27
57	Antifungal monoterpene production in elicited cell suspension cultures of Piqueria trinervia. Phytochemistry, 2000, 55, 51-57.	2.9	16
58	Flavanones and 3-hydroxyflavanones from Lonchocarpus oaxacensis. Phytochemistry, 2000, 55, 953-957.	2.9	21
59	Hypoglycemic effect of extracts and fractions from Psacalium decompositum in healthy and alloxan-diabetic mice. Journal of Ethnopharmacology, 2000, 72, 21-27.	4.1	53
60	Hypoglycemic activity of root water decoction, sesquiterpenoids, and one polysaccharide fraction from Psacalium decompositum in mice. Journal of Ethnopharmacology, 2000, 69, 207-215.	4.1	29
61	Bioactive Compounds from Selected Plants used in the xvi Century Mexican Traditional Medicine. Studies in Natural Products Chemistry, 2000, , 799-844.	1.8	18
62	Flavonoids and Isoflavonoids with Antifungal Properties from <i>Platymiscium yucatanum</i> Heartwood. Holzforschung, 1998, 52, 459-462.	1.9	38
63	Effects of three Mexican medicinal plants (Asteraceae) on blood glucose levels in healthy mice and rabbits. Journal of Ethnopharmacology, 1997, 55, 171-177.	4.1	62
64	Antifungal Xanthones from Calophyllum brasiliensis Heartwood. Journal of Chemical Ecology, 1997, 23, 1901-1911.	1.8	46
65	Transformation of Terpene Piquerol A to Hydroquinone and Phenolic Derivatives. Effect of These Compounds on Weeds. Journal of Agricultural and Food Chemistry, 1996, 44, 2839-2841.	5.2	8
66	A Survey of Alkaloids in the Genera Harpalyce and Brongniartia (FabaceaeBrongniartieae). Biochemical Systematics and Ecology, 1996, 24, 749-755.	1.3	5
67	Antitermitic activity ofLonchocarpus castilloi flavonoids and heartwood extracts. Journal of Chemical Ecology, 1995, 21, 455-463.	1.8	36
68	NMR and X-ray studies of apetalic acid isolated from Calophyllum brasiliense and of its chiral amides. Canadian Journal of Chemistry, 0, , .	1.1	0