Markku Peltonen

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/4199682/publications.pdf

Version: 2024-02-01

292 papers 29,897 citations

76 h-index 165 g-index

295 all docs

295 docs citations

times ranked

295

29885 citing authors

| # | Article | IF | CITATIONS |
|----|---|-----|-----------|
| 1 | Dementia prevention: The potential longâ€term costâ€effectiveness of the FINGER prevention program. Alzheimer's and Dementia, 2023, 19, 999-1008. | 0.4 | 9 |
| 2 | The effect of adherence on cognition in a multidomain lifestyle intervention (FINGER). Alzheimer's and Dementia, 2022, 18, 1325-1334. | 0.4 | 24 |
| 3 | Association of Bariatric Surgery With Cancer Incidence in Patients With Obesity and Diabetes: Long-term Results From the Swedish Obese Subjects Study. Diabetes Care, 2022, 45, 444-450. | 4.3 | 31 |
| 4 | Effect of a multi-domain lifestyle intervention on cardiovascular risk in older people: the FINGER trial. European Heart Journal, 2022, 43, 2054-2061. | 1.0 | 26 |
| 5 | Occupational complexity and cognition in the FINGER multidomain intervention trial. Alzheimer's and Dementia, 2022, 18, 2438-2447. | 0.4 | 4 |
| 6 | Risk factors for major gastrointestinal bleeding in the general population in Finland. World Journal of Gastroenterology, 2022, 28, 2008-2020. | 1.4 | 3 |
| 7 | Prediction of Suicide and Nonfatal Self-harm After Bariatric Surgery: A Risk Score Based on Sociodemographic Factors, Lifestyle Behavior, and Mental Health. Annals of Surgery, 2021, 274, 339-345. | 2.1 | 17 |
| 8 | High need for recovery from work and sleep problems are associated with workers' unhealthy dietary habits. Public Health Nutrition, 2021, 24, 1-10. | 1.1 | 5 |
| 9 | Remission and progression of pre-existing micro- and macroalbuminuria over 15 years after bariatric surgery in Swedish Obese Subjects study. International Journal of Obesity, 2021, 45, 535-546. | 1.6 | 9 |
| 10 | Telomere Length Change in a Multidomain Lifestyle Intervention to Prevent Cognitive Decline: A Randomized Clinical Trial. Journals of Gerontology - Series A Biological Sciences and Medical Sciences, 2021, 76, 491-498. | 1.7 | 11 |
| 11 | Heart failure development in obesity: underlying risk factors and mechanistic pathways. ESC Heart Failure, 2021, 8, 356-367. | 1.4 | 12 |
| 12 | Depression, anxiety, and suicidal ideation in young adults 5Âyears after undergoing bariatric surgery as adolescents. Eating and Weight Disorders, 2021, 26, 1211-1221. | 1.2 | 6 |
| 13 | Diabetes and Cardiovascular Disease Risk Perception and Risk Indicators: a 5-Year Follow-up. International Journal of Behavioral Medicine, 2021, 28, 337-348. | 0.8 | 6 |
| 14 | Earlier life leisure-time physical activity in relation to age-related frailty syndrome. Age and Ageing, 2021, 50, 161-168. | 0.7 | 7 |
| 15 | Impact of the COVIDâ€19 pandemic on statistical design and analysis plans for multidomain intervention clinical trials: Experience from Worldâ€Wide FINGERS. Alzheimer's and Dementia: Translational Research and Clinical Interventions, 2021, 7, e12143. | 1.8 | 13 |
| 16 | Long-term incidence of colorectal cancer after bariatric surgery or usual care in the Swedish Obese Subjects study. PLoS ONE, 2021, 16, e0248550. | 1.1 | 27 |
| 17 | Longâ€term incidence of hypoglycaemiaâ€related events after bariatric surgery or usual care in the Swedish Obese Subjects study: A registerâ€based analysis. Diabetes, Obesity and Metabolism, 2021, 23, 1917-1925. | 2.2 | 2 |
| 18 | Change in CAIDE Dementia Risk Score and Neuroimaging Biomarkers During a 2-Year Multidomain Lifestyle Randomized Controlled Trial: Results of a Post-Hoc Subgroup Analysis. Journals of Gerontology - Series A Biological Sciences and Medical Sciences, 2021, 76, 1407-1414. | 1.7 | 17 |

| # | Article | IF | CITATIONS |
|----|---|------|-----------|
| 19 | Adiponectin Associates with Rheumatoid Arthritis Risk in Overweight and Obesity Independently of Other Adipokines. Journal of Clinical Medicine, 2021, 10, 2791. | 1.0 | 9 |
| 20 | Long-term outcomes of lifestyle intervention to prevent type 2 diabetes in people at high risk in primary health care. Primary Care Diabetes, 2021, 15, 444-450. | 0.9 | 10 |
| 21 | Effect of a Multidomain Lifestyle Intervention on Estimated Dementia Risk. Journal of Alzheimer's Disease, 2021, 82, 1461-1466. | 1.2 | 16 |
| 22 | Long-term risk of anaemia after bariatric surgery: results from the Swedish Obese Subjects study. Lancet Diabetes and Endocrinology,the, 2021, 9, 515-524. | 5.5 | 20 |
| 23 | Pharmacy-based screening to detect persons at elevated risk of type 2 diabetes: a cost-utility analysis. BMC Health Services Research, 2021, 21, 916. | 0.9 | 2 |
| 24 | Koronapandemia - kenen kriisi?. Sosiaalilaaketieteellinen Aikakauslehti, 2021, 58, . | 0.0 | 0 |
| 25 | Type 2 Diabetes-Related Health Economic Impact Associated with Increased Whole Grains Consumption among Adults in Finland. Nutrients, 2021, 13, 3583. | 1.7 | 11 |
| 26 | Bariatric surgery and the incidence of rheumatoid arthritis – a Swedish Obese Subjects study. Rheumatology, 2020, 59, 303-309. | 0.9 | 26 |
| 27 | Association of Bariatric Surgery With Skin Cancer Incidence in Adults With Obesity. JAMA Dermatology, 2020, 156, 38. | 2.0 | 13 |
| 28 | Life Expectancy after Bariatric Surgery in the Swedish Obese Subjects Study. New England Journal of Medicine, 2020, 383, 1535-1543. | 13.9 | 272 |
| 29 | Thirty-Year Incidence and Mortality Trends in Upper and Lower Gastrointestinal Bleeding in Finland. JAMA Network Open, 2020, 3, e2020172. | 2.8 | 18 |
| 30 | Estimating expected life-years and risk factor associations with mortality in Finland: cohort study. BMJ Open, 2020, 10, e033741. | 0.8 | 15 |
| 31 | Elevated adiponectin predicts the development of rheumatoid arthritis in subjects with obesity. Scandinavian Journal of Rheumatology, 2020, 49, 452-460. | 0.6 | 17 |
| 32 | Bariatric surgery versus standard obesity treatment and the risk of severe liver disease: Data from the Swedish Obese Subjects study. Clinical Gastroenterology and Hepatology, 2020, 19, 2675-2676.e2. | 2.4 | 3 |
| 33 | A randomized controlled trial comparing intensive non-surgical treatment with bariatric surgery in adolescents aged 13–16 years (AMOS2): Rationale, study design, and patient recruitment. Contemporary Clinical Trials Communications, 2020, 19, 100592. | 0.5 | 11 |
| 34 | 9p21.3 Coronary Artery Disease Locus Identifies Patients With Treatment Benefit From Bariatric Surgery in the Nonrandomized Prospective Controlled Swedish Obese Subjects Study. Circulation Genomic and Precision Medicine, 2020, 13, 460-465. | 1.6 | 1 |
| 35 | Comparison of Preoperative Remission Scores and Diabetes Duration Alone as Predictors of Durable Type 2 Diabetes Remission and Risk of Diabetes Complications After Bariatric Surgery: A Post Hoc Analysis of Participants From the Swedish Obese Subjects Study. Diabetes Care, 2020, 43, 2804-2811. | 4.3 | 18 |
| 36 | Non-Alcoholic Fatty Liver Disease Markers Associated with Fasting Serum Insulin and Urinary Albumin Excretion Independent of Fasting Plasma Glucose. Journal of Clinical Medicine, 2020, 9, 3161. | 1.0 | 0 |

3

| # | Article | IF | CITATIONS |
|----|--|-----|-----------|
| 37 | Evaluation of Prediction Models for Type 2 Diabetes Relapse After Post-bariatric Surgery Remission: a Post hoc Analysis of 15-Year Follow-up Data from the Swedish Obese Subjects (SOS) Study. Obesity Surgery, 2020, 30, 3955-3960. | 1.1 | 10 |
| 38 | Long-term incidence of gallstone disease after bariatric surgery. Surgery for Obesity and Related Diseases, 2020, 16, 1474-1482. | 1.0 | 24 |
| 39 | Effect and safety of 4% albumin in the treatment of cardiac surgery patients: study protocol for the randomized, double-blind, clinical ALBICS (ALBumin In Cardiac Surgery) trial. Trials, 2020, 21, 235. | 0.7 | 14 |
| 40 | Fracture risk after three bariatric surgery procedures in Swedish obese subjects: up to 26 years followâ€up of a controlled intervention study. Journal of Internal Medicine, 2020, 287, 546-557. | 2.7 | 50 |
| 41 | Worldâ€Wide FINGERS Network: A global approach to risk reduction and prevention of dementia. Alzheimer's and Dementia, 2020, 16, 1078-1094. | 0.4 | 257 |
| 42 | 5-year mental health and eating pattern outcomes following bariatric surgery in adolescents: a prospective cohort study. The Lancet Child and Adolescent Health, 2020, 4, 210-219. | 2.7 | 37 |
| 43 | Letter to the Editor: Effects of Bariatric Surgery on Cancer Risk. Obesity Surgery, 2020, 30, 2036-2036. | 1.1 | 1 |
| 44 | Effects of Bariatric Surgery in Early- and Adult-Onset Obesity in the Prospective Controlled Swedish Obese Subjects Study. Diabetes Care, 2020, 43, 860-866. | 4.3 | 12 |
| 45 | The association of work stress and night work with nutrient intake $\hat{a} \in \hat{a}$ a prospective cohort study. Scandinavian Journal of Work, Environment and Health, 2020, 46, 533-541. | 1.7 | 10 |
| 46 | Long-term incidence of serious fall-related injuries after bariatric surgery in Swedish obese subjects. International Journal of Obesity, 2019, 43, 933-937. | 1.6 | 17 |
| 47 | Trajectories in hypnotic use and approaching death: a register linked case–control study. Sleep Medicine, 2019, 57, 153-161. | 0.8 | 2 |
| 48 | Long-term and recent trends in hypertension awareness, treatment, and control in 12 high-income countries: an analysis of 123 nationally representative surveys. Lancet, The, 2019, 394, 639-651. | 6.3 | 325 |
| 49 | BEST: Bypass equipoise sleeve trial; rationale and design of a randomized, registry-based, multicenter trial comparing Roux-en-Y gastric bypass with sleeve gastrectomy. Contemporary Clinical Trials, 2019, 84, 105809. | 0.8 | 14 |
| 50 | Contribution of rare and common variants to intellectual disability in a sub-isolate of Northern Finland. Nature Communications, 2019, 10, 410. | 5.8 | 32 |
| 51 | Surgical obesity treatment and the risk of heart failure. European Heart Journal, 2019, 40, 2131-2138. | 1.0 | 51 |
| 52 | The Effect of Multidomain Lifestyle Intervention on Daily Functioning in Older People. Journal of the American Geriatrics Society, 2019, 67, 1138-1144. | 1.3 | 35 |
| 53 | Finnish Parkinson's disease study integrating protein-protein interaction network data with exome sequencing analysis. Scientific Reports, 2019, 9, 18865. | 1.6 | 7 |
| 54 | Reoperations After Bariatric Surgery in 26 Years of Follow-up of the Swedish Obese Subjects Study. JAMA Surgery, 2019, 154, 319. | 2,2 | 60 |

| # | Article | IF | CITATIONS |
|----|--|--------------|-----------|
| 55 | Dietary changes and cognition over 2 years within a multidomain intervention trial—The Finnish Geriatric Intervention Study to Prevent Cognitive Impairment and Disability (FINGER). Alzheimer's and Dementia, 2019, 15, 410-417. | 0.4 | 63 |
| 56 | Human PNPLA3-I148M variant increases hepatic retention of polyunsaturated fatty acids. JCI Insight, 2019, 4, . | 2.3 | 93 |
| 57 | Microvascular Outcomes in Patients With Diabetes After Bariatric Surgery. Annals of Internal Medicine, 2019, 170, 506. | 2.0 | 0 |
| 58 | Incidence of end-stage renal disease following bariatric surgery in the Swedish Obese Subjects Study. International Journal of Obesity, 2018, 42, 964-973. | 1.6 | 62 |
| 59 | Differences in participation rates between urban and rural areas are diminishing in Finland. Scandinavian Journal of Public Health, 2018, 46, 755-757. | 1.2 | 5 |
| 60 | Effect of the Apolipoprotein E Genotype on Cognitive Change During a Multidomain Lifestyle Intervention. JAMA Neurology, 2018, 75, 462. | 4.5 | 136 |
| 61 | The effect of age and gender on the genetic regulation of serum 25-hydroxyvitamin D - the FIN-D2D population-based study. Journal of Steroid Biochemistry and Molecular Biology, 2018, 178, 229-233. | 1.2 | 10 |
| 62 | Risk of suicide and non-fatal self-harm after bariatric surgery: results from two matched cohort studies. Lancet Diabetes and Endocrinology,the, 2018, 6, 197-207. | 5 . 5 | 124 |
| 63 | Associations of Bariatric Surgery With Changes in Interpersonal Relationship Status. JAMA Surgery, 2018, 153, 654. | 2.2 | 44 |
| 64 | Cohort Profile: The National FINRISK Study. International Journal of Epidemiology, 2018, 47, 696-696i. | 0.9 | 214 |
| 65 | Multidomain lifestyle intervention benefits a large elderly population at risk for cognitive decline and dementia regardless of baseline characteristics: The FINGER trial. Alzheimer's and Dementia, 2018, 14, 263-270. | 0.4 | 236 |
| 66 | The Effect of a 2-Year Intervention Consisting of Diet, Physical Exercise, Cognitive Training, and Monitoring of Vascular Risk on Chronic Morbidity—the FINGER Randomized Controlled Trial. Journal of the American Medical Directors Association, 2018, 19, 355-360.e1. | 1.2 | 48 |
| 67 | Associations of serum indolepropionic acid, a gut microbiota metabolite, with type 2 diabetes and low-grade inflammation in high-risk individuals. Nutrition and Diabetes, 2018, 8, 35. | 1.5 | 147 |
| 68 | Language as a determinant of participation rates in Finnish health examination surveys. Scandinavian Journal of Public Health, 2018, 46, 240-243. | 1.2 | 7 |
| 69 | Reply: Bariatric surgery and chronic kidney disease: much hope, but proof is still awaited. International Journal of Obesity, 2018, 42, 1534-1534. | 1.6 | 0 |
| 70 | Binge eating and other eating-related problems in adolescents undergoing gastric bypass: results from a Swedish nationwide study (AMOS). Appetite, 2018, 127, 349-355. | 1.8 | 19 |
| 71 | Laparoscopic Roux-en-Y gastric bypass in adolescents with severe obesity (AMOS): a prospective, 5-year, Swedish nationwide study. Lancet Diabetes and Endocrinology,the, 2017, 5, 174-183. | 5. 5 | 226 |
| 72 | Health-related quality of life in a multidomain intervention trial to prevent cognitive decline (FINGER). European Geriatric Medicine, 2017, 8, 164-167. | 1.2 | 24 |

| # | Article | IF | CITATIONS |
|----|---|--------------|-----------|
| 73 | Long-term incidence of microvascular disease after bariatric surgery or usual care in patients with obesity, stratified by baseline glycaemic status: a post-hoc analysis of participants from the Swedish Obese Subjects study. Lancet Diabetes and Endocrinology,the, 2017, 5, 271-279. | 5 . 5 | 111 |
| 74 | Sustained diabetes risk reduction after real life and primary health care setting implementation of the diabetes in Europe prevention using lifestyle, physical activity and nutritional intervention (DE-PLAN) project. BMC Public Health, 2017, 17, 198. | 1.2 | 44 |
| 75 | Effects of bariatric surgery on gout incidence in the Swedish Obese Subjects study: a non-randomised, prospective, controlled intervention trial. Annals of the Rheumatic Diseases, 2017, 76, 688-693. | 0.5 | 55 |
| 76 | Long-term incidence of female-specific cancer after bariatric surgery or usual care in the Swedish Obese Subjects Study. Gynecologic Oncology, 2017, 145, 224-229. | 0.6 | 98 |
| 77 | Diabetes and impaired glucose metabolism is associated with more cold-related cardiorespiratory symptoms. Diabetes Research and Clinical Practice, 2017, 129, 116-125. | 1.1 | 8 |
| 78 | Long-term effects of bariatric surgery in patients with obesity and chromosome 16 p11.2 microdeletion. Surgery for Obesity and Related Diseases, 2017, 13, 1321-1325. | 1.0 | 3 |
| 79 | Changes in total energy intake and macronutrient composition after bariatric surgery predict long-term weight outcome: findings from the Swedish Obese Subjects (SOS) study. American Journal of Clinical Nutrition, 2017, 106, 136-145. | 2.2 | 59 |
| 80 | Sociodemographic and lifestyle factors as determinants of energy intake and macronutrient composition: a 10-year follow-up after bariatric surgery. Surgery for Obesity and Related Diseases, 2017, 13, 1572-1583. | 1.0 | 9 |
| 81 | Bariatric surgery, glycaemic status, and microvascular complications – Authors' reply. Lancet Diabetes and Endocrinology,the, 2017, 5, 416-417. | 5 . 5 | 0 |
| 82 | Pregnancy outcomes after maternal varenicline use; analysis of surveillance data collected by the European Network of Teratology Information Services. Reproductive Toxicology, 2017, 67, 26-34. | 1.3 | 25 |
| 83 | Bariatric Surgery and the Incidence of Psoriasis and Psoriatic Arthritis in the Swedish Obese Subjects Study. Obesity, 2017, 25, 2068-2073. | 1.5 | 41 |
| 84 | Use of HOMA-IR to diagnose non-alcoholic fatty liver disease: a population-based and inter-laboratory study. Diabetologia, 2017, 60, 1873-1882. | 2.9 | 85 |
| 85 | Eveningness has the increased odds for spinal diseases but the decreased odds for articular diseases with prospective hospital treatments. Biological Rhythm Research, 2017, 48, 263-274. | 0.4 | 8 |
| 86 | Baseline Telomere Length and Effects of a Multidomain Lifestyle Intervention on Cognition: The FINGER Randomized Controlled Trial. Journal of Alzheimer's Disease, 2017, 59, 1459-1470. | 1.2 | 20 |
| 87 | Branched-Chain Amino Acid Levels Are Related with Surrogates of Disturbed Lipid Metabolism among Older Men. Frontiers in Medicine, 2016, 3, 57. | 1.2 | 32 |
| 88 | Bariatric Surgery and the Risk of New-Onset Atrial Fibrillation in SwedishÂObese Subjects. Journal of the American College of Cardiology, 2016, 68, 2497-2504. | 1.2 | 159 |
| 89 | Health 2000 score – development and validation of a novel cardiovascular risk score. Annals of Medicine, 2016, 48, 403-409. | 1.5 | 3 |
| 90 | Former male elite athletes have better metabolic health in late life than their controls. Scandinavian Journal of Medicine and Science in Sports, 2016, 26, 284-290. | 1.3 | 16 |

| # | Article | IF | CITATIONS |
|-----|--|-----|-----------|
| 91 | Leisure-Time and Occupational Physical Activity in Early and Late Adulthood in Relation to Later Life Physical Functioning. Journal of Physical Activity and Health, 2016, 13, 1079-1087. | 1.0 | 9 |
| 92 | Determination of a Normal value for HOMA-IR and cut-off Value for Non-Alcoholic Fatty Liver Disease. Journal of Hepatology, 2016, 64, S481. | 1.8 | 1 |
| 93 | Primary prevention and risk factor reduction in coronary heart disease mortality among working aged men and women in eastern Finland over 40 years: population based observational study. BMJ, The, 2016, 352, i721. | 3.0 | 93 |
| 94 | Response to Comment on Sjöholm et al. Weight Change–Adjusted Effects of Gastric Bypass Surgery on Glucose Metabolism: 2- and 10-Year Results From the Swedish Obese Subjects (SOS) Study. Diabetes Care 2016;39:625–631. Diabetes Care, 2016, 39, e85-e85. | 4.3 | 0 |
| 95 | Prevalence of insomniaâ€related symptoms continues to increase in the Finnish workingâ€age population. Journal of Sleep Research, 2016, 25, 454-457. | 1.7 | 66 |
| 96 | Determinants of Diabetes Remission and Glycemic Control After Bariatric Surgery. Diabetes Care, 2016, 39, 166-174. | 4.3 | 152 |
| 97 | Characteristics of adolescents with poor mental health after bariatric surgery. Surgery for Obesity and Related Diseases, 2016, 12, 882-890. | 1.0 | 27 |
| 98 | Weight Change–Adjusted Effects of Gastric Bypass Surgery on Glucose Metabolism: 2- and 10-Year Results From the Swedish Obese Subjects (SOS) Study. Diabetes Care, 2016, 39, 625-631. | 4.3 | 61 |
| 99 | Editorial commentary: Obesity, surgically induced weight loss, and cardiovascular disease. Trends in Cardiovascular Medicine, 2016, 26, 290-291. | 2.3 | 1 |
| 100 | Analysis of the Japanese Diabetes Risk Score and fatty liver markers for incident diabetes in a Japanese cohort. Primary Care Diabetes, 2016, 10, 19-26. | 0.9 | 4 |
| 101 | Changes in CVD Incidence and Mortality Rates, and Life Expectancy: North Karelia and National. Global Heart, 2016, 11, 201. | 0.9 | 10 |
| 102 | Following in the Footsteps of the North Karelia Project: Prevention of Type 2 Diabetes. Global Heart, 2016, 11, 223. | 0.9 | 8 |
| 103 | Predicting Coronary Heart Disease and Stroke: The FINRISK Calculator. Global Heart, 2016, 11, 213. | 0.9 | 19 |
| 104 | Daily Sedentary Time and Risk of Cardiovascular Disease: The National FINRISK 2002 Study. Journal of Physical Activity and Health, 2015, 12, 904-908. | 1.0 | 32 |
| 105 | Adapting existing diabetes risk scores for an Asian population: a risk score for detecting undiagnosed diabetes in the Mongolian population. BMC Public Health, 2015, 15, 938. | 1.2 | 19 |
| 106 | Changes in glucose metabolism in people with different glucose metabolism disorders at baseline: followâ€up results of a Finnish national diabetes prevention programme. Diabetic Medicine, 2015, 32, 1611-1616. | 1.2 | 3 |
| 107 | Twoâ€year trends in psychological outcomes after gastric bypass in adolescents with severe obesity. Obesity, 2015, 23, 1966-1972. | 1.5 | 48 |
| 108 | Former male elite athletes and risk of hypertension in later life. Journal of Hypertension, 2015, 33, 1549-1554. | 0.3 | 10 |

| # | Article | IF | CITATIONS |
|-----|--|-------------|-----------|
| 109 | Forty-year trends in cardiovascular risk factors in Finland. European Journal of Public Health, 2015, 25, 539-546. | 0.1 | 208 |
| 110 | Food and nutrient intake among workers with different shift systems. Occupational and Environmental Medicine, 2015, 72, 513-520. | 1.3 | 66 |
| 111 | Balancing risks and benefits of bariatric surgery for type 2 diabetes. Lancet Diabetes and Endocrinology,the, 2015, 3, 394-395. | 5. 5 | O |
| 112 | The incidence of albuminuria after bariatric surgery and usual care in swedish obese subjects (SOS): a prospective controlled intervention trial. International Journal of Obesity, 2015, 39, 169-175. | 1.6 | 60 |
| 113 | A 2 year multidomain intervention of diet, exercise, cognitive training, and vascular risk monitoring versus control to prevent cognitive decline in at-risk elderly people (FINGER): a randomised controlled trial. Lancet, The, 2015, 385, 2255-2263. | 6.3 | 2,307 |
| 114 | National diabetes prevention program (DEHKO): awareness and self-reported lifestyle changes in Finnish middle-aged population. Public Health, 2015, 129, 210-217. | 1.4 | 15 |
| 115 | Clinical and lifestyle-related risk factors for incident multimorbidity: 10-year follow-up of Finnish population-based cohorts 1982–2012. European Journal of Internal Medicine, 2015, 26, 211-216. | 1.0 | 91 |
| 116 | Secular trends and educational differences in the incidence of type 2 diabetes in Finland, 1972–2007. European Journal of Epidemiology, 2015, 30, 649-659. | 2.5 | 16 |
| 117 | Interrelationships of Physical Activity and Sleep with Cardiovascular Risk Factors: a Person-Oriented Approach. International Journal of Behavioral Medicine, 2015, 22, 735-747. | 0.8 | 10 |
| 118 | Incidence and remission of type 2 diabetes in relation to degree of obesity at baseline and 2Âyear weight change: the Swedish Obese Subjects (SOS) study. Diabetologia, 2015, 58, 1448-1453. | 2.9 | 77 |
| 119 | Changes in lifestyle modestly reduce the estimated cardiovascular disease risk in one-year follow-up of the Finnish diabetes prevention program (FIN-D2D). European Journal of Cardiovascular Nursing, 2015, 14, 145-152. | 0.4 | 18 |
| 120 | Health-care costs over 15 years after bariatric surgery for patients with different baseline glucose status: results from the Swedish Obese Subjects study. Lancet Diabetes and Endocrinology,the, 2015, 3, 855-865. | 5.5 | 66 |
| 121 | Selfâ€Reported Walking Difficulty Predicts Lateâ€Life Mortality in Finnish War Veterans: Results from the Veteran 1992 Project Survey. Journal of the American Geriatrics Society, 2015, 63, 118-123. | 1.3 | 7 |
| 122 | Evaluation of AUSDRISK as a screening tool for lifestyle modification programs: international implications for policy and cost-effectiveness. BMJ Open Diabetes Research and Care, 2015, 3, e000125. | 1.2 | 19 |
| 123 | Circadian preference links to depression in general adult population. Journal of Affective Disorders, 2015, 188, 143-148. | 2.0 | 135 |
| 124 | Evening typology and morning tiredness associates with low leisure time physical activity and high sitting. Chronobiology International, 2015, 32, 1090-1100. | 0.9 | 40 |
| 125 | Psychological aspects of eating behavior as predictors of 10-y weight changes after surgical and conventional treatment of severe obesity: results from the Swedish Obese Subjects intervention study. American Journal of Clinical Nutrition, 2015, 101, 16-24. | 2.2 | 68 |
| 126 | Prevention of diabetes and cardiovascular diseases in occupational health care: Feasibility and effectiveness. Primary Care Diabetes, 2015, 9, 96-104. | 0.9 | 16 |

| # | Article | IF | Citations |
|-----|---|-----|-----------|
| 127 | All-cause and disease-specific mortality among male, former elite athletes: an average 50-year follow-up. British Journal of Sports Medicine, 2015, 49, 893-897. | 3.1 | 86 |
| 128 | Effect of intensive exercise in early adult life on telomere length in later life in men. Journal of Sports Science and Medicine, 2015, 14, 239-45. | 0.7 | 13 |
| 129 | Impact of Health Counselling on Cardiovascular Disease Risk in Middle Aged Men: Influence of Socioeconomic Status. PLoS ONE, 2014, 9, e88959. | 1.1 | 13 |
| 130 | Association of Serum 25-Hydroxyvitamin D with Lifestyle Factors and Metabolic and Cardiovascular Disease Markers: Population-Based Cross-Sectional Study (FIN-D2D). PLoS ONE, 2014, 9, e100235. | 1.1 | 29 |
| 131 | Recruitment and Baseline Characteristics of Participants in the Finnish Geriatric Intervention Study to Prevent Cognitive Impairment and Disability (FINGER)—A Randomized Controlled Lifestyle Trial. International Journal of Environmental Research and Public Health, 2014, 11, 9345-9360. | 1.2 | 69 |
| 132 | Lifestyle Intervention in Prevention of Type 2 Diabetes in Women With a History of Gestational Diabetes Mellitus: One-Year Results of the FIN-D2D Project. Journal of Women's Health, 2014, 23, 506-512. | 1.5 | 25 |
| 133 | Fatty liver and serum cholinesterase are independently correlated with HbA1c levels: Cross-sectional analysis of 5384 people. Journal of International Medical Research, 2014, 42, 542-553. | 0.4 | 11 |
| 134 | Adiponectin and Bariatric Surgery: Associations With Diabetes and Cardiovascular Disease in the Swedish Obese Subjects Study. Diabetes Care, 2014, 37, 1401-1409. | 4.3 | 41 |
| 135 | The impact of weight reduction in the prevention of the progression of obstructive sleep apnea: an explanatory analysis of a 5-year observational follow-up trial. Sleep Medicine, 2014, 15, 329-335. | 0.8 | 38 |
| 136 | Perceived Sufficiency of Physical Activity Levels Among Adults at High Risk of Type 2 Diabetes: The FIN-D2D Study. International Journal of Behavioral Medicine, 2014, 21, 99-108. | 0.8 | 5 |
| 137 | Endothelial function is well preserved in obese patients with mild obstructive sleep apnea. Sleep and Breathing, 2014, 18, 177-186. | 0.9 | 11 |
| 138 | A former career as a male elite athleteâ€"does it protect against type 2 diabetes in later life?. Diabetologia, 2014, 57, 270-274. | 2.9 | 27 |
| 139 | Physical activity and sleep profiles in Finnish men and women. BMC Public Health, 2014, 14, 82. | 1.2 | 32 |
| 140 | Do depressive symptoms have an impact on the effectiveness of lifestyle counseling in prevention of type 2 diabetes? One-year follow-up of FIN-D2D. Primary Care Diabetes, 2014, 8, 43-47. | 0.9 | 4 |
| 141 | Long-term consistency of diurnal-type preferences among men. Chronobiology International, 2014, 31, 182-188. | 0.9 | 79 |
| 142 | Evening chronotypes have the increased odds for bronchial asthma and nocturnal asthma. Chronobiology International, 2014, 31, 95-101. | 0.9 | 50 |
| 143 | Association of Bariatric Surgery With Long-term Remission of Type 2 Diabetes and With Microvascular and Macrovascular Complications. JAMA - Journal of the American Medical Association, 2014, 311, 2297. | 3.8 | 849 |
| 144 | Telomere length in circulating leukocytes is associated with lung function and disease. European Respiratory Journal, 2014, 43, 983-992. | 3.1 | 103 |

| # | Article | IF | CITATIONS |
|-----|---|-----|-----------|
| 145 | A population-based study on the prevalence of NASH using scores validated against liver histology. Journal of Hepatology, 2014, 60, 839-846. | 1.8 | 107 |
| 146 | Gaby Mahlberg and Dirk Wiemann, eds. European Contexts for English Republicanism. Politics and Culture in Europe, 1650–1750. Farnham: Ashgate Publishing Limited, 2013. xiii + 274 pp. \$124.95. ISBN: 978-1-49094-5556-1 Renaissance Quarterly, 2014, 67, 634-635. | 0.0 | 0 |
| 147 | The Association between HbA1c, Fasting Glucose, 1-Hour Glucose and 2-Hour Glucose during an Oral Glucose Tolerance Test and Cardiovascular Disease in Individuals with Elevated Risk for Diabetes. PLoS ONE, 2014, 9, e109506. | 1.1 | 38 |
| 148 | Behavioral Trait of Morningness-Eveningness in Association with Articular and Spinal Diseases in a Population. PLoS ONE, 2014, 9, e114635. | 1.1 | 35 |
| 149 | Evening types are prone to depression. Chronobiology International, 2013, 30, 719-725. | 0.9 | 192 |
| 150 | Fatty liver score and 15-year incidence of type 2 diabetes. Hepatology International, 2013, 7, 610-621. | 1.9 | 11 |
| 151 | Mortality in middle-aged men with obstructive sleep apnea in Finland. Sleep and Breathing, 2013, 17, 1047-1053. | 0.9 | 21 |
| 152 | Computationally estimated apolipoproteins B and A1 in predicting cardiovascular risk. Atherosclerosis, 2013, 226, 245-251. | 0.4 | 23 |
| 153 | Associations of Chronotype and Sleep With Cardiovascular Diseases and Type 2 Diabetes. Chronobiology International, 2013, 30, 470-477. | 0.9 | 270 |
| 154 | The Finnish Geriatric Intervention Study to Prevent Cognitive Impairment and Disability (FINGER): Study design and progress. Alzheimer's and Dementia, 2013, 9, 657-665. | 0.4 | 385 |
| 155 | Changes in all-cause and cardiovascular disease mortality in three different Finnish population cohorts with and without diabetes. International Journal of Cardiology, 2013, 168, 4734-4738. | 0.8 | 4 |
| 156 | Alcohol consumption and alcohol problems after bariatric surgery in the swedish obese subjects study. Obesity, 2013, 21, 2444-2451. | 1.5 | 136 |
| 157 | Financial satisfaction and its relationship to depressive symptoms in middle-aged and older adults: Results from the FIN-D2D survey. International Journal of Social Psychiatry, 2013, 59, 239-246. | 1.6 | 17 |
| 158 | Predictors of Success of a Lifestyle Intervention in Relation to Weight Loss and Improvement in Glucose Tolerance Among Individuals at High Risk for Type 2 Diabetes. Journal of Primary Care and Community Health, 2013, 4, 59-66. | 1.0 | 15 |
| 159 | Improved lifestyle and decreased diabetes risk over 13Âyears: long-term follow-up of the randomised Finnish Diabetes Prevention Study (DPS). Diabetologia, 2013, 56, 284-293. | 2.9 | 416 |
| 160 | Obesity surgery and incidence of heart failure. European Heart Journal, 2013, 34, 3555-3555. | 1.0 | 1 |
| 161 | Obesity surgery and incidence of atrial fibrillation. European Heart Journal, 2013, 34, P4091-P4091. | 1.0 | 0 |
| 162 | Evaluation of Current Eligibility Criteria for Bariatric Surgery. Diabetes Care, 2013, 36, 1335-1340. | 4.3 | 68 |

| # | Article | IF | Citations |
|-----|---|------------|-----------|
| 163 | Prevalence, predictors and covariates of functional status impairment among Finnish Second World War veterans during 1992-2004. Age and Ageing, 2013, 42, 508-514. | 0.7 | 9 |
| 164 | Long-Term Effect of Bariatric Surgery on Liver Enzymes in the Swedish Obese Subjects (SOS) Study. PLoS ONE, 2013, 8, e60495. | 1.1 | 69 |
| 165 | Relation of Chronotype to Sleep Complaints in the General Finnish Population < /b>. Chronobiology International, 2012, 29, 311-317. | 0.9 | 205 |
| 166 | Bariatric Surgery and Long-term Cardiovascular Events. JAMA - Journal of the American Medical Association, 2012, 307, 56. | 3.8 | 1,341 |
| 167 | Relative validity of a FFQ in measuring carbohydrate fractions, dietary glycaemic index and load: exploring the effects of subject characteristics. British Journal of Nutrition, 2012, 107, 1367-1375. | 1.2 | 35 |
| 168 | Bariatric Surgery and Prevention of Type 2 Diabetes in Swedish Obese Subjects. New England Journal of Medicine, 2012, 367, 695-704. | 13.9 | 698 |
| 169 | Two-year outcome of laparoscopic Roux-en-Y gastric bypass in adolescents with severe obesity: results from a Swedish Nationwide Study (AMOS). International Journal of Obesity, 2012, 36, 1388-1395. | 1.6 | 119 |
| 170 | Virtues in Elizabethan and Early Stuart Grammar Schools. Journal of Medieval and Early Modern Studies, 2012, 42, 157-179. | 0.2 | 3 |
| 171 | Effect of short-term carbohydrate overfeeding and long-term weight loss on liver fat in overweight humans. American Journal of Clinical Nutrition, 2012, 96, 727-734. | 2.2 | 171 |
| 172 | Do statins interfere with lifestyle intervention in the prevention of diabetes in primary healthcare? One-year follow-up of the FIN-D2D project. BMJ Open, 2012, 2, e001472. | 0.8 | 14 |
| 173 | Intra-individual changes in body weight in population-based cohorts during four decades: the Finnish FINRISK study. European Journal of Public Health, 2012, 22, 107-112. | 0.1 | 17 |
| 174 | Comparison of the MDRD Study and the CKD-EPI Study equations in evaluating trends of estimated kidney function at population level: findings from the National FINRISK Study. Nephrology Dialysis Transplantation, 2012, 27, 3210-3217. | 0.4 | 14 |
| 175 | Health Care Use During 20 Years Following Bariatric Surgery. JAMA - Journal of the American Medical Association, 2012, 308, 1132. | 3.8 | 131 |
| 176 | James Kuzner. Open Subjects: English Renaissance Republicans, Modern Selfhoods, and the Virtue of Vulnerability. Edinburgh Critical Studies in Renaissance Culture. Edinburgh: Edinburgh University Press, 2011. x + 222 pp. £65. ISBN: 978–0–7486–4253–3 Renaissance Quarterly, 2012, 65, 1349-135 | 0.0 51. | 0 |
| 177 | Effect of weight loss on inflammation in patients with mild obstructive sleep apnea. Nutrition, Metabolism and Cardiovascular Diseases, 2012, 22, 583-590. | 1.1 | 19 |
| 178 | Determinants of changes in self-rated health among Finnish war veterans: Results from the Veteran Project 1992 and 2004 surveys. Archives of Gerontology and Geriatrics, 2012, 55, 343-348. | 1.4 | 6 |
| 179 | Perceived need to increase physical activity levels among adults at high risk of type 2 diabetes. A cross-sectional analysis within a community-based diabetes prevention project FIN-D2D. BMC Public Health, 2012, 12, 514. | 1.2 | 14 |
| 180 | Relationships between depressive symptoms and self-reported unintentional injuries: the cross-sectional populationâ€"based FIN-D2D survey. BMC Public Health, 2012, 12, 516. | 1.2 | 9 |

| # | Article | IF | CITATIONS |
|-----|---|-----|-----------|
| 181 | Occupational health care identifies risk for type 2 diabetes and cardiovascular disease. Primary Care Diabetes, 2012, 6, 95-102. | 0.9 | 24 |
| 182 | Participation, socioeconomic status and group or individual counselling intervention in individuals at high risk for type 2 diabetes: One-year follow-up study of the FIN-D2D-project. Primary Care Diabetes, 2012, 6, 277-283. | 0.9 | 11 |
| 183 | Cardiovascular Events After Bariatric Surgery in Obese Subjects With Type 2 Diabetes. Diabetes Care, 2012, 35, 2613-2617. | 4.3 | 152 |
| 184 | Paradoxical Lower Serum Triglyceride Levels and Higher Type 2 Diabetes Mellitus Susceptibility in Obese Individuals with the PNPLA3 148M Variant. PLoS ONE, 2012, 7, e39362. | 1.1 | 78 |
| 185 | Trends in estimated kidney function: the FINRISK surveys. European Journal of Epidemiology, 2012, 27, 305-313. | 2.5 | 23 |
| 186 | Family history of diabetes and effectiveness of lifestyle counselling on the cardioâ€metabolic risk profile in individuals at high risk of Type 2 diabetes: 1â€year followâ€up of the FINâ€D2D project. Diabetic Medicine, 2012, 29, 207-211. | 1,2 | 17 |
| 187 | Effects of bariatric surgery on disability pension in Swedish obese subjects. International Journal of Obesity, 2012, 36, 356-362. | 1.6 | 17 |
| 188 | Changes in body mass index and measures of abdominal obesity in Finnish adults between 1992 and 2007, the National FINRISK Study. Clinical Obesity, 2012, 2, 57-63. | 1.1 | 19 |
| 189 | Lifetime leisure-time physical activity and the risk of depressive symptoms at the ages of 65–74years: The FIN-D2D survey. Preventive Medicine, 2012, 54, 313-315. | 1.6 | 16 |
| 190 | Leukocyte Telomere Length in the Finnish Diabetes Prevention Study. PLoS ONE, 2012, 7, e34948. | 1.1 | 65 |
| 191 | Diurnal Evening Type is Associated with Current Smoking, Nicotine Dependence and Nicotine Intake in the Population Based National FINRISK 2007 Study. Journal of Addiction Research & Therapy, 2012, 01, . | 0.2 | 20 |
| 192 | Ageing and associations of fasting plasma glucose and 2h plasma glucose with HbA1C in apparently healthy population. â€∞FIN-D2D―study. Diabetes Research and Clinical Practice, 2011, 93, 344-349. | 1.1 | 12 |
| 193 | Self-reported sleep duration, all-cause mortality, cardiovascular mortality and morbidity in Finland. Sleep Medicine, 2011, 12, 215-221. | 0.8 | 159 |
| 194 | HbA $<$ sub $>$ 1c $<$ /sub $>$ in diagnosing and predicting Typeâ \in f2 diabetes in impaired glucose tolerance: the Finnish Diabetes Prevention Study. Diabetic Medicine, 2011, 28, 36-42. | 1,2 | 36 |
| 195 | The use of fasting vs. non-fasting triglyceride concentration for estimating the prevalence of high LDL-cholesterol and metabolic syndrome in population surveys. BMC Medical Research Methodology, 2011, 11, 63. | 1.4 | 13 |
| 196 | Population-level effects of the national diabetes prevention programme (FIN-D2D) on the body weight, the waist circumference, and the prevalence of obesity. BMC Public Health, 2011, 11, 350. | 1.2 | 18 |
| 197 | Metabolically healthy and unhealthy obesity phenotypes in the general population: the FIN-D2D Survey. BMC Public Health, 2011, 11, 754. | 1.2 | 133 |
| 198 | Increase in physical activity and cardiometabolic risk profile change during lifestyle intervention in primary healthcare: 1-year follow-up study among individuals at high risk for type 2 diabetes. BMJ Open, 2011, 1, e000292-e000292. | 0.8 | 31 |

| # | Article | IF | Citations |
|-----|---|-----|-----------|
| 199 | Socioeconomic position and effectiveness of lifestyle intervention in prevention of type 2 diabetes: One-year follow-up of the FIN-D2D project. Scandinavian Journal of Public Health, 2011, 39, 561-570. | 1.2 | 28 |
| 200 | Serum calcium level is associated with metabolic syndrome in the general population: FIN-D2D study. European Journal of Endocrinology, 2011, 165, 429-434. | 1.9 | 40 |
| 201 | Impact of Positive Family History and Genetic Risk Variants on the Incidence of Diabetes: The Finnish Diabetes Prevention Study. Diabetes Care, 2011, 34, 418-423. | 4.3 | 44 |
| 202 | Prevention of type 2 diabetes by lifestyle intervention in primary health care setting in Poland: Diabetes in Europe Prevention using Lifestyle, physical Activity and Nutritional intervention (DE-PLAN) project. British Journal of Diabetes and Vascular Disease, 2011, 11, 198-203. | 0.6 | 23 |
| 203 | Socio-economic differences in dysglycemia and lifestyle-related risk factors in the Finnish middle-aged population. European Journal of Public Health, 2011, 21, 768-774. | 0.1 | 23 |
| 204 | Reply to E Hemmingsson et al. American Journal of Clinical Nutrition, 2011, 93, 474-475. | 2.2 | 0 |
| 205 | Self-reported sleep apnoea and mortality in patients from the Swedish Obese Subjects study. European Respiratory Journal, 2011, 38, 1349-1354. | 3.1 | 16 |
| 206 | Impaired nasal breathing may prevent the beneficial effect of weight loss in the treatment of OSA. Rhinology, 2011, 49, 587-592. | 0.7 | 2 |
| 207 | Impaired nasal breathing may prevent the beneficial effect of weight loss in the treatment of OSA. Rhinology, 2011, 49, 587-592. | 0.7 | 4 |
| 208 | The activation of the inflammatory cytokines in overweight patients with mild obstructive sleep apnoea. Journal of Sleep Research, 2010, 19, 341-348. | 1.7 | 68 |
| 209 | Non-alcoholic and alcoholic Fatty Liver Disease - two Diseases of Affluence associated with the Metabolic Syndrome and Type 2 Diabetes: the FIN-D2D Survey. BMC Public Health, 2010, 10, 237. | 1.2 | 66 |
| 210 | Changes in sleep duration and changes in weight in obese patients: The Swedish Obese Subjects Study. Sleep and Biological Rhythms, 2010, 8, 63-71. | 0.5 | 15 |
| 211 | Thirty-five-year trends in cardiovascular risk factors in Finland. International Journal of Epidemiology, 2010, 39, 504-518. | 0.9 | 429 |
| 212 | Trends in health status and the use of services among the 80-84-year-old Finnish war veterans during 1992 and 2004. Scandinavian Journal of Public Health, 2010, 38, 434-441. | 1.2 | 5 |
| 213 | Take Action to Prevent Diabetes – The IMAGE Toolkit for the Prevention of Type 2 Diabetes in Europe. Hormone and Metabolic Research, 2010, 42, S37-S55. | 0.7 | 197 |
| 214 | Reducing the risk of type 2 diabetes with nutrition and physical activity – efficacy and implementation of lifestyle interventions in Finland. Public Health Nutrition, 2010, 13, 993-999. | 1.1 | 50 |
| 215 | A European Evidence-Based Guideline for the Prevention of Type 2 Diabetes. Hormone and Metabolic Research, 2010, 42, S3-S36. | 0.7 | 385 |
| 216 | Despite evidence-based guidelines, systolic blood pressure remains inadequately controlled in older hypertensive adults. Journal of Human Hypertension, 2010, 24, 439-446. | 1.0 | 16 |

| # | Article | IF | Citations |
|-----|---|-------------|---------------|
| 217 | Quality Indicators for the Prevention of Type 2 Diabetes in Europe – IMAGE. Hormone and Metabolic Research, 2010, 42, S56-S63. | 0.7 | 40 |
| 218 | Sustained improvement in mild obstructive sleep apnea after a diet- and physical activity–based lifestyle intervention: postinterventional follow-up. American Journal of Clinical Nutrition, 2010, 92, 688-696. | 2.2 | 87 |
| 219 | Cardiometabolic profile of people screened for high risk of type 2 diabetes in a national diabetes prevention programme (FIN-D2D). Primary Care Diabetes, 2010, 4, 231-239. | 0.9 | 24 |
| 220 | Lifestyle Intervention for Prevention of Type 2 Diabetes in Primary Health Care. Diabetes Care, 2010, 33, 2146-2151. | 4.3 | 265 |
| 221 | Development and validation of a risk-score model for subjects with impaired glucose tolerance for the assessment of the risk of type 2 diabetes mellitus—The STOP-NIDDM risk-score. Diabetes Research and Clinical Practice, 2010, 87, 267-274. | 1.1 | 35 |
| 222 | Leisure-time physical activity and metabolic syndrome plus depressive symptoms in the FIN-D2D survey. Preventive Medicine, 2010, 51, 466-470. | 1.6 | 16 |
| 223 | Leisure-Time Physical Activity and the Metabolic Syndrome in the Finnish Diabetes Prevention Study. Diabetes Care, 2010, 33, 1610-1617. | 4.3 | 74 |
| 224 | Diabetes, Alzheimer disease, and vascular dementia. Neurology, 2010, 75, 1195-1202. | 1.5 | 422 |
| 225 | Ten-Year Mortality and Cardiovascular Morbidity in the Finnish Diabetes Prevention Study—Secondary Analysis of the Randomized Trial. PLoS ONE, 2009, 4, e5656. | 1.1 | 158 |
| 226 | Lifestyle Intervention with Weight Reduction. American Journal of Respiratory and Critical Care Medicine, 2009, 179, 320-327. | 2.5 | 361 |
| 227 | Sleep Duration, Lifestyle Intervention, and Incidence of Type 2 Diabetes in Impaired Glucose Tolerance. Diabetes Care, 2009, 32, 1965-1971. | 4.3 | 102 |
| 228 | Surgical intervention represents a feasible option for patients with mild obstructive sleep apnoea. Acta Oto-Laryngologica, 2009, 129, 1266-1273. | 0.3 | 7 |
| 229 | Anti-inflammatory effect of lifestyle changes in the Finnish Diabetes Prevention Study. Diabetologia, 2009, 52, 433-442. | 2.9 | 133 |
| 230 | Educational attainment and effectiveness of lifestyle intervention in the Finnish Diabetes Prevention Study. Diabetes Research and Clinical Practice, 2009, 86, e1-e5. | 1.1 | 24 |
| 231 | Prediction of Non-Alcoholic Fatty Liver Disease and Liver Fat Using Metabolic and Genetic Factors. Gastroenterology, 2009, 137, 865-872. | 0.6 | 646 |
| 232 | Effects of bariatric surgery on cancer incidence in obese patients in Sweden (Swedish Obese Subjects) Tj ETQq0 | 0 g.rgBT /0 | Overlock 10 1 |
| 233 | Prevalence, awareness and treatment of hypertension in Finland during 1982–2007. Journal of Hypertension, 2009, 27, 1552-1559. | 0.3 | 65 |
| 234 | Effectiveness of Superficial Venous Surgery in Terms of Quality-Adjusted Life Years and Costs. Scandinavian Journal of Surgery, 2009, 98, 229-233. | 1.3 | 7 |

| # | Article | IF | Citations |
|-----|--|-----|-----------|
| 235 | Sleep-disordered breathing is related to an increased risk for type 2 diabetes in middle-aged men, but not in women – the FIN-D2D survey. Diabetes, Obesity and Metabolism, 2008, 10, 468-475. | 2.2 | 35 |
| 236 | Trends in selfâ€reported sleep duration and insomniaâ€related symptoms in Finland from 1972 to 2005: a comparative review and reâ€analysis of Finnish population samples. Journal of Sleep Research, 2008, 17, 54-62. | 1.7 | 216 |
| 237 | High prevalence of obesity, central obesity and abnormal glucose tolerance in the middle-aged Finnish population. BMC Public Health, 2008, 8, 423. | 1.2 | 52 |
| 238 | Sleep duration is associated with an increased risk for the prevalence of type 2 diabetes in middle-aged women – The FIN-D2D survey. Sleep Medicine, 2008, 9, 221-227. | 0.8 | 88 |
| 239 | Leisure time physical activity in individuals with screen-detected type 2 diabetes compared to those with known type 2 diabetes. Diabetes Research and Clinical Practice, 2008, 81, 110-116. | 1.1 | 13 |
| 240 | Determinants for the Effectiveness of Lifestyle Intervention in the Finnish Diabetes Prevention Study. Diabetes Care, 2008, 31, 857-862. | 4.3 | 134 |
| 241 | Effect of Lifestyle Intervention on the Occurrence of Metabolic Syndrome and its Components in the Finnish Diabetes Prevention Study. Diabetes Care, 2008, 31, 805-807. | 4.3 | 178 |
| 242 | Determinants for the Effectiveness of Lifestyle Intervention in the Finnish Diabetes Prevention Study. Diabetes Care, 2008, 31, e88-e88. | 4.3 | 0 |
| 243 | The European Perspective of Type 2 Diabetes Prevention: Diabetes in Europe - Prevention Using Lifestyle, Physical Activity and Nutritional Intervention (DE-PLAN) Project. Experimental and Clinical Endocrinology and Diabetes, 2008, 116, 167-172. | 0.6 | 144 |
| 244 | The Increasing Prevalence of Metabolic Syndrome among Finnish Men and Women over a Decade. Journal of Clinical Endocrinology and Metabolism, 2008, 93, 832-836. | 1.8 | 58 |
| 245 | Effect of Weight Reduction on Rhinometric Measurements in Overweight Patients with Obstructive Sleep Apnea. American Journal of Rhinology & Allergy, 2008, 22, 410-415. | 2.3 | 32 |
| 246 | Costs of a Self-Selected, Health-Promoting Diet Among the Participants of the Finnish Diabetes Prevention Study. Diabetes Care, 2007, 30, 1275-1277. | 4.3 | 18 |
| 247 | Prevention of Type 2 Diabetes - Lessons we have Learnt for Implementation. Hormone and Metabolic Research, 2007, 39, 636-641. | 0.7 | 32 |
| 248 | National type 2 diabetes prevention programme in Finland: FIN-D2D. International Journal of Circumpolar Health, 2007, 66, 101-112. | 0.5 | 162 |
| 249 | Two Year Reduction In Sleep Apnea Symptoms and Associated Diabetes Incidence After Weight Loss In Severe Obesity. Sleep, 2007, 30, 703-710. | 0.6 | 128 |
| 250 | Lifestyle intervention to prevent diabetes in men and women with impaired glucose tolerance is cost-effective. International Journal of Technology Assessment in Health Care, 2007, 23, 177-183. | 0.2 | 77 |
| 251 | Effect of smoking on lifestyle interventions to prevent diabetes – Authors' reply. Lancet, The, 2007, 369, 365-366. | 6.3 | 0 |
| 252 | Sustained reduction in the incidence of type 2 diabetes by lifestyle intervention: follow-up of the Finnish Diabetes Prevention Study. Lancet, The, 2006, 368, 1673-1679. | 6.3 | 1,530 |

| # | Article | IF | Citations |
|-----|--|------|-----------|
| 253 | Association of leisure time physical activity and abdominal obesity with fasting serum insulin and 2-h postchallenge plasma glucose levels. Diabetic Medicine, 2006, 23, 1025-1028. | 1.2 | 23 |
| 254 | High-fibre, low-fat diet predicts long-term weight loss and decreased type 2 diabetes risk: the Finnish Diabetes Prevention Study. Diabetologia, 2006, 49, 912-920. | 2.9 | 249 |
| 255 | Joint association of coffee consumption and other factors to the risk of type 2 diabetes: a prospective study in Finland. International Journal of Obesity, 2006, 30, 1742-1749. | 1.6 | 56 |
| 256 | Systemic Immune Mediators and Lifestyle Changes in the Prevention of Type 2 Diabetes: Results From the Finnish Diabetes Prevention Study. Diabetes, 2006, 55, 2340-2346. | 0.3 | 110 |
| 257 | Effortâ€Related Calf Pain in the Obese and Longâ€Term Changes after Surgical Obesity Treatment ^{**} . Obesity, 2005, 13, 137-145. | 4.0 | 11 |
| 258 | The effects of physical activity and body mass index on cardiovascular, cancer and all-cause mortality among 47 212 middle-aged Finnish men and women. International Journal of Obesity, 2005, 29, 894-902. | 1.6 | 237 |
| 259 | Urinary sodium and potassium excretion and the risk of type 2 diabetes: a prospective study in Finland. Diabetologia, 2005, 48, 1477-1483. | 2.9 | 106 |
| 260 | The Impact of History of Hypertension and Type 2 Diabetes at Baseline on the Incidence of Stroke and Stroke Mortality. Stroke, 2005, 36, 2538-2543. | 1.0 | 142 |
| 261 | The Gender-Specific Impact of Diabetes and Myocardial Infarction at Baseline and During Follow-Up on Mortality From All Causes and Coronary Heart Disease. Journal of the American College of Cardiology, 2005, 45, 1413-1418. | 1.2 | 68 |
| 262 | Cross-sectional evaluation of the Finnish Diabetes Risk Score: a tool to identify undetected type 2 diabetes, abnormal glucose tolerance and metabolic syndrome. Diabetes and Vascular Disease Research, 2005, 2, 67-72. | 0.9 | 273 |
| 263 | Lifestyle strategies for weight control: experience from the Finnish Diabetes Prevention Study. Proceedings of the Nutrition Society, 2005, 64, 81-88. | 0.4 | 42 |
| 264 | Low and high circulating cortisol levels predict mortality and cognitive dysfunction early after stroke. Journal of Internal Medicine, 2004, 256, 15-21. | 2.7 | 86 |
| 265 | Body composition in the SOS (Swedish Obese Subjects) reference study. International Journal of Obesity, 2004, 28, 1317-1324. | 1.6 | 44 |
| 266 | Lifestyle, Diabetes, and Cardiovascular Risk Factors 10 Years after Bariatric Surgery. New England Journal of Medicine, 2004, 351, 2683-2693. | 13.9 | 4,023 |
| 267 | Intermittent versus on-demand use of a very low calorie diet: a randomized 2-year clinical trial. Journal of Internal Medicine, 2003, 253, 463-471. | 2.7 | 49 |
| 268 | A dietary and behavioural programme for the treatment of obesity. A 4-year clinical trial and a long-term posttreatment follow-up. Journal of Internal Medicine, 2003, 254, 272-279. | 2.7 | 65 |
| 269 | High Expression of Complement Components in Omental Adipose Tissue in Obese Men. Obesity, 2003, 11, 699-708. | 4.0 | 195 |
| 270 | Gallstones, gallbladder disease, and pancreatitis Cross-Sectional and 2-year data from the Swedish obese subjects (SOS) and SOS reference studies. American Journal of Gastroenterology, 2003, 98, 1032-1041. | 0.2 | 1 |

| # | Article | IF | CITATIONS |
|-----|--|-----|-----------|
| 271 | Musculoskeletal pain in the obese: a comparison with a general population and long-term changes after conventional and surgical obesity treatment. Pain, 2003, 104, 549-557. | 2.0 | 232 |
| 272 | Impact of blood pressure and insulin on the relationship between body fat and left ventricular structure. European Heart Journal, 2003, 24, 1500-1505. | 1.0 | 38 |
| 273 | Gallstones, Gallbladder Disease, and Pancreatitis: Cross-Sectional and 2-Year Data From The Swedish Obese Subjects (Sos) and Sos Reference Studies. American Journal of Gastroenterology, 2003, 98, 1032-1041. | 0.2 | 87 |
| 274 | Potassium per kilogram fat-free mass and total body potassium: predictions from sex, age, and anthropometry. American Journal of Physiology - Endocrinology and Metabolism, 2003, 284, E416-E423. | 1.8 | 10 |
| 275 | Pharmaceutical Costs in Obese Individuals. Archives of Internal Medicine, 2002, 162, 2061. | 4.3 | 128 |
| 276 | Long-term effects of weight loss on pharmaceutical costs in obese subjects. A report from the SOS intervention study. International Journal of Obesity, 2002, 26, 184-192. | 1.6 | 55 |
| 277 | Cost of Inâ€Patient Care over 7 Years among Surgically and Conventionally Treated Obese Patients. Obesity, 2002, 10, 1276-1283. | 4.0 | 43 |
| 278 | Depotâ€Specific Expression of Fibroblast Growth Factors in Human Adipose Tissue. Obesity, 2002, 10, 608-616. | 4.0 | 74 |
| 279 | Blood Pressure and Pulse Pressure during Longâ€Term Weight Loss in the Obese: The Swedish Obese Subjects (SOS) Intervention Study. Obesity, 2001, 9, 188-195. | 4.0 | 117 |
| 280 | Widening Gap of Stroke Between East and West. Stroke, 2000, 31, 2-8. | 1.0 | 65 |
| 281 | Marked improvement in survival after acute myocardial infarction in middle-aged men but not in women. The Northern Sweden MONICA study 1985-94. Journal of Internal Medicine, 2000, 247, 579-587. | 2.7 | 42 |
| 282 | Social Patterning of Myocardial Infarction and Stroke in Sweden: Incidence and Survival. American Journal of Epidemiology, 2000, 151, 283-292. | 1.6 | 76 |
| 283 | Differentiated Long-Term Effects of Intentional Weight Loss on Diabetes and Hypertension. Hypertension, 2000, 36, 20-25. | 1.3 | 340 |
| 284 | Factors Associated With Delayed Admission to Hospital and In-Hospital Delays> in Acute Stroke and TIA. Stroke, 1999, 30, 40-48. | 1.0 | 243 |
| 285 | High prevalence of undiagnosed coeliac disease in adults: a Swedish population-based study. Journal of Internal Medicine, 1999, 245, 63-68. | 2.7 | 169 |
| 286 | Stroke Units in Their Natural Habitat. Stroke, 1999, 30, 709-714. | 1.0 | 107 |
| 287 | Secular trends in social patterning of cardiovascular risk factor levels in Sweden. The Northern Sweden MONICA Study 1986–1994. Journal of Internal Medicine, 1998, 243, 1-9. | 2.7 | 3 |
| 288 | Secular trends in social patterning of cardiovascular risk factor levels in Sweden. The Northern Sweden MONICA Study 1986–1994. Journal of Internal Medicine, 1998, 244, 1-9. | 2.7 | 52 |

| # | Article | IF | CITATIONS |
|-----|---|-----|-----------|
| 289 | Time Trends in Long-term Survival After Stroke. Stroke, 1998, 29, 1358-1365. | 1.0 | 49 |
| 290 | Age-period-cohort effects on ischaemic heart disease mortality in Sweden from 1969 to 1993, and forecasts up to 2003. European Heart Journal, 1997, 18, 1307-1312. | 1.0 | 27 |
| 291 | Two variants of extracellular-superoxide dismutase: relationship to cardiovascular risk factors in an unselected middle-aged population. Journal of Internal Medicine, 1997, 242, 5-14. | 2.7 | 81 |
| 292 | Age-Period-Cohort Effects on Stroke Mortality in Sweden 1969-1993 and Forecasts Up to the Year 2003. Stroke, 1996, 27, 1981-1985. | 1.0 | 30 |