
Tomomasa Takamiya

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/4217353/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	The Sequential Organ Failure Assessment (SOFA) score predicts mortality and neurological outcome in patients with post-cardiac arrest syndrome. Journal of Cardiology, 2020, 76, 295-302.	1.9	20
2	Propensity-matched comparison of cryoballoon and radiofrequency ablation for atrial fibrillation in elderly patients. Heart Rhythm, 2019, 16, 838-845.	0.7	18
3	Characteristics of the nonpulmonary vein foci induced after secondâ€generation cryoballoon ablation for paroxysmal atrial fibrillation. Journal of Cardiovascular Electrophysiology, 2020, 31, 174-184.	1.7	14
4	The impact of left atrium size on selection of the pulmonary vein isolation method for atrial fibrillation: Cryoballoon or radiofrequency catheter ablation. American Heart Journal, 2021, 231, 82-92.	2.7	14
5	Presence of non-pulmonary vein foci in patients with atrial fibrillation undergoing standard ablation of pulmonary vein isolation: Clinical characteristics and long-term ablation outcome. IJC Heart and Vasculature, 2021, 32, 100717.	1.1	14
6	Differences in the electrophysiological findings of repeat ablation between patients who first underwent cryoballoon ablation and radiofrequency catheter ablation for paroxysmal atrial fibrillation. Journal of Cardiovascular Electrophysiology, 2019, 30, 1792-1800.	1.7	13
7	Pulmonary vein isolation plus left atrial posterior wall isolation and additional nonpulmonary vein trigger ablation using highâ€dose isoproterenol for longâ€standing persistent atrial fibrillation. Journal of Arrhythmia, 2019, 35, 215-222.	1.2	11
8	Cryoballoon versus radiofrequency ablation for paroxysmal atrial fibrillation in hemodialysis patients. Heart and Vessels, 2020, 35, 1709-1716.	1.2	8
9	Association between the locations of non-pulmonary vein triggers and ablation outcomes in repeat procedures after cryoballoon ablation of paroxysmal atrial fibrillation. Journal of Interventional Cardiac Electrophysiology, 2021, 62, 381-389.	1.3	6
10	One-year outcomes after pulmonary vein isolation plus posterior wall isolation and additional non-pulmonary vein trigger ablation for persistent atrial fibrillation with or without contact force sensing: a propensity score-matched comparison. Journal of Interventional Cardiac Electrophysiology, 2020, 59, 585-593.	1.3	5
11	Impact of diagnosisâ€ŧoâ€ablation time on nonâ€pulmonary vein triggers and ablation outcomes in persistent atrial fibrillation. Journal of Cardiovascular Electrophysiology, 2021, 32, 1251-1258.	1.7	4
12	Effect of isolation feasibility of non-pulmonary vein foci on efficacy of ablation for atrial fibrillation: comparison of the isolation and focal ablation methods. Journal of Interventional Cardiac Electrophysiology, 2022, 65, 441-451.	1.3	4
13	Successful Percutaneous Coronary Intervention for Atherosclerotic Coronary Lesion with Anomalous Origin of the Right Coronary Artery. Case Reports in Medicine, 2018, 2018, 1-7.	0.7	3
14	Identification of an epicardial slow conduction channel using ripple mapping in ablation of postinfarct ventricular tachycardia. Journal of Cardiovascular Electrophysiology, 2019, 30, 1156-1158.	1.7	2
15	Catheter ablation for atrial fibrillation in patients with congestive heart failure. International Journal of Cardiology, 2021, 333, 98-104.	1.7	1
16	Optimal single procedure strategy of pulmonary vein isolation with cryoballoon or radiofrequency and non-pulmonary vein triggers ablation for non-paroxysmal atrial fibrillation. IJC Heart and Vasculature, 2022, 40, 101021.	1.1	1