

Robin P F Dullaart

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4236553/publications.pdf>

Version: 2024-02-01

124
papers

3,685
citations

101543

36
h-index

161849

54
g-index

124
all docs

124
docs citations

124
times ranked

5971
citing authors

#	ARTICLE	IF	CITATIONS
1	Application of non-HDL cholesterol for population-based cardiovascular risk stratification: results from the Multinational Cardiovascular Risk Consortium. <i>Lancet, The</i> , 2019, 394, 2173-2183.	13.7	177
2	Disturbed Vitamin A Metabolism in Non-Alcoholic Fatty Liver Disease (NAFLD). <i>Nutrients</i> , 2018, 10, 29.	4.1	138
3	Genome Wide Association Identifies Common Variants at the SERPINA6/SERPINA1 Locus Influencing Plasma Cortisol and Corticosteroid Binding Globulin. <i>PLoS Genetics</i> , 2014, 10, e1004474.	3.5	105
4	Role of HDL Cholesterol and Estimates of HDL Particle Composition in Future Development of Type 2 Diabetes in the General Population: The PREVEND Study. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2013, 98, E1352-E1359.	3.6	98
5	Circulating Total Bilirubin and Risk of Incident Cardiovascular Disease in the General Population. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2015, 35, 716-724.	2.4	96
6	TMAO is Associated with Mortality: Impact of Modestly Impaired Renal Function. <i>Scientific Reports</i> , 2017, 7, 13781.	3.3	96
7	Genetic Variation at the <i>Phospholipid Transfer Protein</i> Locus Affects Its Activity and High-Density Lipoprotein Size and Is a Novel Marker of Cardiovascular Disease Susceptibility. <i>Circulation</i> , 2010, 122, 470-477.	1.6	86
8	Efficacy of β -Blockers on Hemodynamic Control during Pheochromocytoma Resection: A Randomized Controlled Trial. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2020, 105, 2381-2391.	3.6	85
9	GlycA, a marker of acute phase glycoproteins, and the risk of incident type 2 diabetes mellitus: PREVEND study. <i>Clinica Chimica Acta</i> , 2016, 452, 10-17.	1.1	80
10	Skin Autofluorescence Is Associated With 5-Year Mortality and Cardiovascular Events in Patients With Peripheral Artery Disease. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2014, 34, 933-938.	2.4	78
11	HDL (High-Density Lipoprotein) Cholesterol Efflux Capacity Is Associated With Incident Cardiovascular Disease in the General Population. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2019, 39, 1874-1883.	2.4	78
12	Prevalence and determinants of non-alcoholic fatty liver disease in lifelines: A large Dutch population cohort. <i>PLoS ONE</i> , 2017, 12, e0171502.	2.5	74
13	Combined Effects of Smoking and Alcohol on Metabolic Syndrome: The LifeLines Cohort Study. <i>PLoS ONE</i> , 2014, 9, e96406.	2.5	73
14	HDL Cholesterol Efflux Predicts Graft Failure in Renal Transplant Recipients. <i>Journal of the American Society of Nephrology: JASN</i> , 2016, 27, 595-603.	6.1	71
15	The anti-inflammatory function of HDL is impaired in type 2 diabetes: role of hyperglycemia, paraoxonase-1 and low grade inflammation. <i>Cardiovascular Diabetology</i> , 2017, 16, 132.	6.8	71
16	A novel NMR-based assay to measure circulating concentrations of branched-chain amino acids: Elevation in subjects with type 2 diabetes mellitus and association with carotid intima media thickness. <i>Clinical Biochemistry</i> , 2018, 54, 92-99.	1.9	71
17	High plasma cholesteryl ester transfer protein levels may favour reduced incidence of cardiovascular events in men with low triglycerides. <i>European Heart Journal</i> , 2007, 28, 1012-1018.	2.2	68
18	The Effect of Cholesteryl Ester Transfer Protein β 29Câ†’A Promoter Polymorphism on High-Density Lipoprotein Cholesterol Is Dependent on Serum Triglycerides. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2005, 90, 4198-4204.	3.6	67

#	ARTICLE	IF	CITATIONS
19	Carotid artery intima media thickness is inversely related to serum free thyroxine in euthyroid subjects. <i>Clinical Endocrinology</i> , 2007, 67, 668-673.	2.4	67
20	Non-Alcoholic Fatty Liver Disease and Risk of Incident Type 2 Diabetes: Role of Circulating Branched-Chain Amino Acids. <i>Nutrients</i> , 2019, 11, 705.	4.1	67
21	High-Density Lipoprotein Anti-Inflammatory Capacity and Incident Cardiovascular Events. <i>Circulation</i> , 2021, 143, 1935-1945.	1.6	67
22	Assessing the functional properties of high-density lipoproteins: an emerging concept in cardiovascular research. <i>Biomarkers in Medicine</i> , 2013, 7, 457-472.	1.4	62
23	Pleiotropic Effects of Lipid Genes on Plasma Glucose, HbA1c, and HOMA-IR Levels. <i>Diabetes</i> , 2014, 63, 3149-3158.	0.6	61
24	Plasma Branched-Chain Amino Acids and Risk of Incident Type 2 Diabetes: Results from the PREVENT Prospective Cohort Study. <i>Journal of Clinical Medicine</i> , 2018, 7, 513.	2.4	60
25	Lower plasma adiponectin is a marker of increased intima-media thickness associated with type 2 diabetes mellitus and with male gender. <i>European Journal of Endocrinology</i> , 2007, 156, 387-394.	3.7	54
26	Plasma Lecithin: Cholesterol Acyltransferase Activity Is Elevated in Metabolic Syndrome and Is an Independent Marker of Increased Carotid Artery Intima Media Thickness. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2008, 93, 4860-4866.	3.6	54
27	Impaired HDL cholesterol efflux in metabolic syndrome is unrelated to glucose tolerance status: the CODAM study. <i>Scientific Reports</i> , 2016, 6, 27367.	3.3	53
28	High plasma C-reactive protein (CRP) is related to low paraoxonase (PON) activity independently of high leptin and low adiponectin in type 2 diabetes mellitus. <i>Clinical Endocrinology</i> , 2009, 70, 221-226.	2.4	52
29	Association of Thyroid Dysfunction With Cognitive Function. <i>JAMA Internal Medicine</i> , 2021, 181, 1440.	5.1	51
30	HDL function is impaired in acute myocardial infarction independent of plasma HDL cholesterol levels. <i>Journal of Clinical Lipidology</i> , 2016, 10, 1318-1328.	1.5	50
31	Branched Chain Amino Acids Are Associated with Insulin Resistance Independent of Leptin and Adiponectin in Subjects with Varying Degrees of Glucose Tolerance. <i>Metabolic Syndrome and Related Disorders</i> , 2017, 15, 183-186.	1.3	49
32	Associations Between High-Density Lipoprotein Particles and Ischemic Events by Vascular Domain, Sex, and Ethnicity. <i>Circulation</i> , 2020, 142, 657-669.	1.6	49
33	Common variation in the CETP gene and the implications for cardiovascular disease and its treatment: an updated analysis. <i>Pharmacogenomics</i> , 2008, 9, 747-763.	1.3	48
34	Association of Plasma Concentration of Vitamin B ₁₂ With All-Cause Mortality in the General Population in the Netherlands. <i>JAMA Network Open</i> , 2020, 3, e1919274.	5.9	45
35	Unenhanced CT imaging is highly sensitive to exclude pheochromocytoma: a multicenter study. <i>European Journal of Endocrinology</i> , 2018, 178, 431-437.	3.7	44
36	Ketone Bodies Are Mildly Elevated in Subjects with Type 2 Diabetes Mellitus and Are Inversely Associated with Insulin Resistance as Measured by the Lipoprotein Insulin Resistance Index. <i>Journal of Clinical Medicine</i> , 2020, 9, 321.	2.4	40

#	ARTICLE	IF	CITATIONS
37	Increased Risk of Atrial Fibrillation After Treatment for Differentiated Thyroid Carcinoma. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2015, 100, 4563-4569.	3.6	39
38	The Relation Between Thyroid Function and Anemia: A Pooled Analysis of Individual Participant Data. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2018, 103, 3658-3667.	3.6	39
39	Effects of Hydrocortisone on the Regulation of Blood Pressure: Results From a Randomized Controlled Trial. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2016, 101, 3691-3699.	3.6	34
40	The HDL anti-inflammatory function is impaired in myocardial infarction and may predict new cardiac events independent of HDL cholesterol. <i>Clinica Chimica Acta</i> , 2014, 433, 34-38.	1.1	32
41	Cigarette smoking is associated with higher thyroid hormone and lower TSH levels: the PREVENT study. <i>Endocrine</i> , 2020, 67, 613-622.	2.3	32
42	Lipoprotein insulin resistance index, a high-throughput measure of insulin resistance, is associated with incident type II diabetes mellitus in the Prevention of Renal and Vascular End-Stage Disease study. <i>Journal of Clinical Lipidology</i> , 2019, 13, 129-137.e1.	1.5	31
43	Low thyroid function is not associated with an accelerated deterioration in renal function. <i>Nephrology Dialysis Transplantation</i> , 2019, 34, 650-659.	0.7	31
44	Circulating trimethylamine N-oxide is associated with all-cause mortality in subjects with nonalcoholic fatty liver disease. <i>Liver International</i> , 2021, 41, 2371-2382.	3.9	31
45	Calculating the optimal surveillance for head and neck paraganglioma in SDHB-mutation carriers. <i>Familial Cancer</i> , 2017, 16, 123-130.	1.9	30
46	Genome-Wide Association Study and Identification of a Protective Missense Variant on Lipoprotein(a) Concentration. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2021, 41, 1792-1800.	2.4	29
47	Oxidative stress is associated with suspected nonalcoholic fatty liver disease and all-cause mortality in the general population. <i>Liver International</i> , 2020, 40, 2148-2159.	3.9	28
48	High Serum PCSK9 Is Associated With Increased Risk of New-Onset Diabetes After Transplantation in Renal Transplant Recipients. <i>Diabetes Care</i> , 2017, 40, 894-901.	8.6	27
49	High Betaine, a Trimethylamine N-Oxide Related Metabolite, Is Prospectively Associated with Low Future Risk of Type 2 Diabetes Mellitus in the PREVENT Study. <i>Journal of Clinical Medicine</i> , 2019, 8, 1813.	2.4	27
50	HDL Particle Subspecies and Their Association With Incident Type 2 Diabetes: The PREVENT Study. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2021, 106, 1761-1772.	3.6	27
51	Association of beta-hydroxybutyrate with development of heart failure: Sex differences in a Dutch population cohort. <i>European Journal of Clinical Investigation</i> , 2021, 51, e13468.	3.4	25
52	Plasma Levels of Retinol Binding Protein 4 Relate to Large VLDL and Small LDL Particles in Subjects with and without Type 2 Diabetes. <i>Journal of Clinical Medicine</i> , 2019, 8, 1792.	2.4	24
53	The effects of two different doses of hydrocortisone on cognition in patients with secondary adrenal insufficiency – Results from a randomized controlled trial. <i>Psychoneuroendocrinology</i> , 2015, 55, 36-47.	2.7	23
54	Triglyceride-rich lipoprotein and LDL particle subfractions and their association with incident type 2 diabetes: the PREVENT study. <i>Cardiovascular Diabetology</i> , 2021, 20, 156.	6.8	23

#	ARTICLE	IF	CITATIONS
55	The Interaction of Age and Type 2 Diabetes on Executive Function and Memory in Persons Aged 35 Years or Older. PLoS ONE, 2013, 8, e82991.	2.5	22
56	The Relationship of the Anti-Oxidant Bilirubin with Free Thyroxine Is Modified by Insulin Resistance in Euthyroid Subjects. PLoS ONE, 2014, 9, e90886.	2.5	21
57	Soluble Vascular Cell Adhesion Molecules May be Protective of Future Cardiovascular Disease Risk: Findings from the PREVEND Prospective Cohort Study. Journal of Atherosclerosis and Thrombosis, 2017, 24, 804-818.	2.0	21
58	A Newly Developed Diabetes Risk Index, Based on Lipoprotein Subfractions and Branched Chain Amino Acids, is Associated with Incident Type 2 Diabetes Mellitus in the PREVEND Cohort. Journal of Clinical Medicine, 2020, 9, 2781.	2.4	21
59	Nonalcoholic fatty liver disease, circulating ketone bodies and all-cause mortality in a general population-based cohort. European Journal of Clinical Investigation, 2021, 51, e13627.	3.4	20
60	An individual participant data analysis of prospective cohort studies on the association between subclinical thyroid dysfunction and depressive symptoms. Scientific Reports, 2020, 10, 19111.	3.3	19
61	Prescription of statins in suspected non-alcoholic fatty liver disease and high cardiovascular risk, a population-based study. Liver International, 2019, 39, 1343-1354.	3.9	18
62	Characterization of LP-Z Lipoprotein Particles and Quantification in Subjects with Liver Disease Using a Newly Developed NMR-Based Assay. Journal of Clinical Medicine, 2020, 9, 2915.	2.4	18
63	Plasma lecithin:cholesterol acyltransferase and phospholipid transfer protein activity independently associate with nonalcoholic fatty liver disease. European Journal of Clinical Investigation, 2018, 48, e12988.	3.4	17
64	Lower Plasma Magnesium, Measured by Nuclear Magnetic Resonance Spectroscopy, is Associated with Increased Risk of Developing Type 2 Diabetes Mellitus in Women: Results from a Dutch Prospective Cohort Study. Journal of Clinical Medicine, 2019, 8, 169.	2.4	16
65	Higher Sodium Intake Assessed by 24 Hour Urinary Sodium Excretion Is Associated with Non-Alcoholic Fatty Liver Disease: The PREVEND Cohort Study. Journal of Clinical Medicine, 2019, 8, 2157.	2.4	16
66	Hypomagnesaemia and its determinants in a contemporary primary care cohort of persons with type 2 diabetes. Endocrine, 2020, 67, 80-86.	2.3	16
67	Downregulation of cholesteryl ester transfer protein by glucocorticoids: a randomised study on HDL. European Journal of Clinical Investigation, 2017, 47, 494-503.	3.4	15
68	Glucocorticoid replacement is associated with hypertriglyceridaemia, elevated glucose and higher non-HDL cholesterol and may diminish the association of HDL cholesterol with the CETP promoter polymorphism in GH-receiving hypopituitary patients. Clinical Endocrinology, 2008, 69, 359-366.	2.4	14
69	Plasma C-Peptide and Risk of Developing Type 2 Diabetes in the General Population. Journal of Clinical Medicine, 2020, 9, 3001.	2.4	14
70	Primary aldosteronism is associated with decreased low-density and high-density lipoprotein particle concentrations and increased GlycA, a pro-inflammatory glycoprotein biomarker. Clinical Endocrinology, 2019, 90, 79-87.	2.4	13
71	HDL Cholesterol Efflux Capacity is Impaired in Severe Short-Term Hypothyroidism Despite Increased HDL Cholesterol. Journal of Clinical Endocrinology and Metabolism, 2020, 105, e3355-e3362.	3.6	13
72	Effect of Metformin Treatment on Lipoprotein Subfractions in Non-Diabetic Patients with Acute Myocardial Infarction: A Glycometabolic Intervention as Adjunct to Primary Coronary Intervention in ST Elevation Myocardial Infarction (GIPS-III) Trial. PLoS ONE, 2016, 11, e0145719.	2.5	13

#	ARTICLE	IF	CITATIONS
73	The predictive value of the antioxidative function of HDL for cardiovascular disease and graft failure in renal transplant recipients. <i>Atherosclerosis</i> , 2016, 249, 181-185.	0.8	12
74	Emerging role of dopamine in neovascularization of pheochromocytoma and paraganglioma. <i>FASEB Journal</i> , 2017, 31, 2226-2240.	0.5	12
75	Plasma ApoE elevations are associated with NAFLD: The PREVEND Study. <i>PLoS ONE</i> , 2019, 14, e0220659.	2.5	11
76	Antibodies Against the C-Terminus of ApoA-1 Are Inversely Associated with Cholesterol Efflux Capacity and HDL Metabolism in Subjects with and without Type 2 Diabetes Mellitus. <i>International Journal of Molecular Sciences</i> , 2019, 20, 732.	4.1	11
77	Obesity, adipokines and COVID-19. <i>European Journal of Clinical Investigation</i> , 2020, 50, e13313.	3.4	11
78	Plasma creatine and incident type 2 diabetes in a general population-based cohort: The PREVEND study. <i>Clinical Endocrinology</i> , 2021, 94, 563-574.	2.4	11
79	Remnant lipoprotein cholesterol is associated with incident new onset diabetes after transplantation (NODAT) in renal transplant recipients: results of the TransplantLines Biobank and cohort Studies. <i>Cardiovascular Diabetology</i> , 2022, 21, 41.	6.8	11
80	No influence of antihypertensive agents on plasma free metanephrines. <i>Clinical Biochemistry</i> , 2016, 49, 1368-1371.	1.9	10
81	Mass spectrometric quantification of salivary metanephrines—A study in healthy subjects. <i>Clinical Biochemistry</i> , 2016, 49, 983-988.	1.9	10
82	Autoantibodies to Apolipoprotein A-1 as Independent Predictors of Cardiovascular Mortality in Renal Transplant Recipients. <i>Journal of Clinical Medicine</i> , 2019, 8, 948.	2.4	10
83	Thyroid function and risk of all-cause and cardiovascular mortality: a prospective population-based cohort study. <i>Endocrine</i> , 2021, 71, 385-396.	2.3	10
84	Circulating Trimethylamine N-Oxide Is Associated with Increased Risk of Cardiovascular Mortality in Type-2 Diabetes: Results from a Dutch Diabetes Cohort (ZODIAC-59). <i>Journal of Clinical Medicine</i> , 2021, 10, 2269.	2.4	10
85	Dopamine concentration in blood platelets is elevated in patients with head and neck paragangliomas. <i>Clinical Chemistry and Laboratory Medicine</i> , 2016, 54, 1395-401.	2.3	9
86	High-Density Lipoprotein Particles and Their Relationship to Posttransplantation Diabetes Mellitus in Renal Transplant Recipients. <i>Biomolecules</i> , 2020, 10, 481.	4.0	9
87	The Systemic Redox Status Is Maintained in Non-Smoking Type 2 Diabetic Subjects Without Cardiovascular Disease: Association with Elevated Triglycerides and Large VLDL. <i>Journal of Clinical Medicine</i> , 2020, 9, 49.	2.4	8
88	Association of Circulating Trimethylamine N-Oxide and Its Dietary Determinants with the Risk of Kidney Graft Failure: Results of the TransplantLines Cohort Study. <i>Nutrients</i> , 2021, 13, 262.	4.1	8
89	Adiposity and the development of dyslipidemia in APOE $\epsilon\epsilon$ homozygous subjects: A longitudinal analysis in two population-based cohorts. <i>Atherosclerosis</i> , 2021, 325, 57-62.	0.8	8
90	Higher hydrocortisone dose increases bilirubin in hypopituitary patients—results from an RCT. <i>European Journal of Clinical Investigation</i> , 2016, 46, 475-480.	3.4	7

#	ARTICLE	IF	CITATIONS
91	Vitamin B6, Inflammation, and Cardiovascular Outcome in a Population-Based Cohort: The Prevention of Renal and Vascular End-Stage Disease (PREVEND) Study. <i>Nutrients</i> , 2020, 12, 2711.	4.1	7
92	Endogenous urinary glucocorticoid metabolites and mortality in prednisolone-treated renal transplant recipients. <i>Clinical Transplantation</i> , 2020, 34, e13824.	1.6	7
93	Association of hepcidin with survival after kidney transplantation. <i>European Journal of Clinical Investigation</i> , 2016, 46, 994-1001.	3.4	6
94	High Plasma Branched-Chain Amino Acids Are Associated with Higher Risk of Post-Transplant Diabetes Mellitus in Renal Transplant Recipients. <i>Journal of Clinical Medicine</i> , 2020, 9, 511.	2.4	6
95	Blood Mg ²⁺ is more closely associated with hyperglycaemia than with hypertriglycerolaemia: the PREVEND study. <i>Diabetologia</i> , 2019, 62, 1732-1734.	6.3	5
96	Plasma branched chain amino acids are lower in short-term profound hypothyroidism and increase in response to thyroid hormone supplementation. <i>Scandinavian Journal of Clinical and Laboratory Investigation</i> , 2020, 80, 562-566.	1.2	5
97	Renin-angiotensin-aldosterone responsiveness to low sodium and blood pressure reactivity to angiotensin-II are unrelated to cholesteryl ester transfer protein mass in healthy subjects. <i>Expert Opinion on Therapeutic Targets</i> , 2008, 12, 1321-1328.	3.4	4
98	The HDL anti-inflammatory function is impaired in the context of low-normal free thyroxine in diabetic and nondiabetic individuals. <i>Clinical Endocrinology</i> , 2018, 88, 752-754.	2.4	4
99	Sex Differences in Lipid Profile across the Life Span in Patients with Type 2 Diabetes: A Primary Care-Based Study. <i>Journal of Clinical Medicine</i> , 2021, 10, 1775.	2.4	4
100	Lipoprotein particle sizes and incident type 2 diabetes: the PREVEND cohort study. <i>Diabetologia</i> , 2022, 65, 402-405.	6.3	4
101	Temporal Course of Plasma Trimethylamine N-Oxide (TMAO) Levels in ST-Elevation Myocardial Infarction. <i>Journal of Clinical Medicine</i> , 2021, 10, 5677.	2.4	4
102	Subclinical thyroid dysfunction and depressive symptoms: protocol for a systematic review and individual participant data meta-analysis of prospective cohort studies. <i>BMJ Open</i> , 2019, 9, e029716.	1.9	3
103	Lipoprotein(a): The Renaissance of an Enigmatic Lipoprotein. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2020, 105, e896-e898.	3.6	3
104	Association of diuretic use with increased risk for long-term post-transplantation diabetes mellitus in kidney transplant recipients. <i>Nephrology Dialysis Transplantation</i> , 2022, , .	0.7	3
105	Higher free triiodothyronine is associated with higher HDL particle concentration and smaller HDL particle size. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2022, , .	3.6	3
106	Profoundly Disturbed Lipoproteins in Cirrhotic Patients: Role of Lipoprotein-Z, a Hepatotoxic LDL-like Lipoprotein. <i>Journal of Clinical Medicine</i> , 2022, 11, 1223.	2.4	3
107	Research update for articles published in <i>EJCI</i> in 2014. <i>European Journal of Clinical Investigation</i> , 2016, 46, 880-894.	3.4	2
108	Urinary prednisolone excretion is a determinant of serum hepcidin levels in renal transplant recipients. <i>American Journal of Hematology</i> , 2017, 92, E173-E175.	4.1	2

#	ARTICLE	IF	CITATIONS
109	Nuclear Magnetic Resonance-Measured Ionized Magnesium Is Inversely Associated with Type 2 Diabetes in the Insulin Resistance Atherosclerosis Study. <i>Nutrients</i> , 2022, 14, 1792.	4.1	2
110	High plasma levels of betaine, a trimethylamine N-oxide related metabolite, are associated with the severity of cirrhosis. <i>Liver International</i> , 2022, , .	3.9	2
111	Lipoprotein insulin resistance score and branched-chain amino acids increase after adrenalectomy for unilateral aldosterone-producing adenoma: a preliminary study. <i>Endocrine</i> , 2020, 68, 420-426.	2.3	1
112	Low Detection Rates of Genetic FH in Cohort of Patients With Severe Hypercholesterolemia in the United Arab Emirates. <i>Frontiers in Genetics</i> , 2021, 12, 809256.	2.3	1
113	Incorporating Baseline Outcome Data in Individual Participant Data Meta-Analysis of Non-randomized Studies. <i>Frontiers in Psychiatry</i> , 2022, 13, 774251.	2.6	1
114	Fasting Proinsulin Independently Predicts Incident Type 2 Diabetes in the General Population. <i>Journal of Personalized Medicine</i> , 2022, 12, 1131.	2.5	1
115	Research update for articles published in EJCI in 2009. <i>European Journal of Clinical Investigation</i> , 2011, 41, 1149-1163.	3.4	0
116	PS4 - 22. HDL-cholesterol, Apolipoprotein A-I/A-II, and HDL-cholesterol particle composition for the risk of developing type 2 diabetes in the community: the PREVEND Study. <i>Nederlands Tijdschrift Voor Diabetologie</i> , 2012, 10, 113-113.	0.0	0
117	Research update for articles published in EJCI in 2016. <i>European Journal of Clinical Investigation</i> , 2018, 48, e13016.	3.4	0
118	Research update for articles published in EJCI in 2017. <i>European Journal of Clinical Investigation</i> , 2019, 49, e13163.	3.4	0
119	P1650HIGH-DENSITY LIPOPROTEIN PARTICLES AND THEIR RELATIONSHIP TO POSTTRANSPLANTATION DIABETES IN RENAL TRANSPLANT RECIPIENTS. <i>Nephrology Dialysis Transplantation</i> , 2020, 35, .	0.7	0
120	Plasma Vitamin C and Risk of Late Graft Failure in Kidney Transplant Recipients: Results of the TransplantLines Biobank and Cohort Study. <i>Antioxidants</i> , 2021, 10, 631.	5.1	0
121	MO589PLASMA CONCENTRATIONS OF TRIMETHYLAMINE N-OXIDE, AND ITS DIETARY DETERMINANTS ARE ASSOCIATED WITH INCREASED RISK OF GRAFT FAILURE. <i>Nephrology Dialysis Transplantation</i> , 2021, 36, .	0.7	0
122	Hdl Cholesterol Efflux Predicts Graft Failure But Not Cardiovascular And Overall Mortality In Renal Transplant Recipients. , 2014, , .		0
123	Somatosensory function in patients with secondary adrenal insufficiency treated with two different doses of hydrocortisone—Results from a randomized controlled trial. <i>PLoS ONE</i> , 2017, 12, e0180326.	2.5	0
124	Lipoprotein Z, an abnormal LDL-like lipoprotein, independently predicts mortality in cirrhosis. <i>European Journal of Internal Medicine</i> , 2022, , .	2.2	0