

Robin P F Dullaart

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4236553/publications.pdf>

Version: 2024-02-01

124
papers

3,685
citations

101543

36
h-index

161849

54
g-index

124
all docs

124
docs citations

124
times ranked

5971
citing authors

#	ARTICLE	IF	CITATIONS
1	Lipoprotein particle sizes and incident type 2 diabetes: the PREVEND cohort study. <i>Diabetologia</i> , 2022, 65, 402-405.	6.3	4
2	Association of diuretic use with increased risk for long-term post-transplantation diabetes mellitus in kidney transplant recipients. <i>Nephrology Dialysis Transplantation</i> , 2022, , .	0.7	3
3	Higher free triiodothyronine is associated with higher HDL particle concentration and smaller HDL particle size. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2022, , .	3.6	3
4	Incorporating Baseline Outcome Data in Individual Participant Data Meta-Analysis of Non-randomized Studies. <i>Frontiers in Psychiatry</i> , 2022, 13, 774251.	2.6	1
5	Profoundly Disturbed Lipoproteins in Cirrhotic Patients: Role of Lipoprotein-Z, a Hepatotoxic LDL-like Lipoprotein. <i>Journal of Clinical Medicine</i> , 2022, 11, 1223.	2.4	3
6	Remnant lipoprotein cholesterol is associated with incident new onset diabetes after transplantation (NODAT) in renal transplant recipients: results of the TransplantLines Biobank and cohort Studies. <i>Cardiovascular Diabetology</i> , 2022, 21, 41.	6.8	11
7	Lipoprotein Z, an abnormal LDL-like lipoprotein, independently predicts mortality in cirrhosis. <i>European Journal of Internal Medicine</i> , 2022, , .	2.2	0
8	Nuclear Magnetic Resonance-Measured Ionized Magnesium Is Inversely Associated with Type 2 Diabetes in the Insulin Resistance Atherosclerosis Study. <i>Nutrients</i> , 2022, 14, 1792.	4.1	2
9	High plasma levels of betaine, a trimethylamine <sc>Nâ€Oxide</sc> â€related metabolite, are associated with the severity of cirrhosis. <i>Liver International</i> , 2022, , .	3.9	2
10	Fasting Proinsulin Independently Predicts Incident Type 2 Diabetes in the General Population. <i>Journal of Personalized Medicine</i> , 2022, 12, 1131.	2.5	1
11	Thyroid function and risk of all-cause and cardiovascular mortality: a prospective population-based cohort study. <i>Endocrine</i> , 2021, 71, 385-396.	2.3	10
12	Plasma creatine and incident type 2 diabetes in a general populationâ€based cohort: The PREVEND study. <i>Clinical Endocrinology</i> , 2021, 94, 563-574.	2.4	11
13	Association of betaâ€hydroxybutyrate with development of heart failure: Sex differences in a Dutch population cohort. <i>European Journal of Clinical Investigation</i> , 2021, 51, e13468.	3.4	25
14	Association of Circulating Trimethylamine N-Oxide and Its Dietary Determinants with the Risk of Kidney Graft Failure: Results of the TransplantLines Cohort Study. <i>Nutrients</i> , 2021, 13, 262.	4.1	8
15	HDL Particle Subspecies and Their Association With Incident Type 2 Diabetes: The PREVEND Study. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2021, 106, 1761-1772.	3.6	27
16	Plasma Vitamin C and Risk of Late Graft Failure in Kidney Transplant Recipients: Results of the TransplantLines Biobank and Cohort Study. <i>Antioxidants</i> , 2021, 10, 631.	5.1	0
17	Sex Differences in Lipid Profile across the Life Span in Patients with Type 2 Diabetes: A Primary Care-Based Study. <i>Journal of Clinical Medicine</i> , 2021, 10, 1775.	2.4	4
18	Circulating Trimethylamine N-Oxide Is Associated with Increased Risk of Cardiovascular Mortality in Type-2 Diabetes: Results from a Dutch Diabetes Cohort (ZODIAC-59). <i>Journal of Clinical Medicine</i> , 2021, 10, 2269.	2.4	10

#	ARTICLE	IF	CITATIONS
19	MO589 PLASMA CONCENTRATIONS OF TRIMETHYLAMINE N-OXIDE, AND ITS DIETARY DETERMINANTS ARE ASSOCIATED WITH INCREASED RISK OF GRAFT FAILURE. <i>Nephrology Dialysis Transplantation</i> , 2021, 36, .	0.7	0
20	High-Density Lipoprotein Anti-Inflammatory Capacity and Incident Cardiovascular Events. <i>Circulation</i> , 2021, 143, 1935-1945.	1.6	67
21	Adiposity and the development of dyslipidemia in APOE ϵ 2 homozygous subjects: A longitudinal analysis in two population-based cohorts. <i>Atherosclerosis</i> , 2021, 325, 57-62.	0.8	8
22	Genome-Wide Association Study and Identification of a Protective Missense Variant on Lipoprotein(a) Concentration. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2021, 41, 1792-1800.	2.4	29
23	Circulating trimethylamine N-oxide is associated with all-cause mortality in subjects with nonalcoholic fatty liver disease. <i>Liver International</i> , 2021, 41, 2371-2382.	3.9	31
24	Nonalcoholic fatty liver disease, circulating ketone bodies and all-cause mortality in a general population-based cohort. <i>European Journal of Clinical Investigation</i> , 2021, 51, e13627.	3.4	20
25	Triglyceride-rich lipoprotein and LDL particle subfractions and their association with incident type 2 diabetes: the PREVEND study. <i>Cardiovascular Diabetology</i> , 2021, 20, 156.	6.8	23
26	Association of Thyroid Dysfunction With Cognitive Function. <i>JAMA Internal Medicine</i> , 2021, 181, 1440.	5.1	51
27	Temporal Course of Plasma Trimethylamine N-Oxide (TMAO) Levels in ST-Elevation Myocardial Infarction. <i>Journal of Clinical Medicine</i> , 2021, 10, 5677.	2.4	4
28	Low Detection Rates of Genetic FH in Cohort of Patients With Severe Hypercholesterolemia in the United Arab Emirates. <i>Frontiers in Genetics</i> , 2021, 12, 809256.	2.3	1
29	Cigarette smoking is associated with higher thyroid hormone and lower TSH levels: the PREVEND study. <i>Endocrine</i> , 2020, 67, 613-622.	2.3	32
30	Hypomagnesaemia and its determinants in a contemporary primary care cohort of persons with type 2 diabetes. <i>Endocrine</i> , 2020, 67, 80-86.	2.3	16
31	Efficacy of β -Blockers on Hemodynamic Control during Pheochromocytoma Resection: A Randomized Controlled Trial. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2020, 105, 2381-2391.	3.6	85
32	Lipoprotein(a): The Renaissance of an Enigmatic Lipoprotein. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2020, 105, e896-e898.	3.6	3
33	The Systemic Redox Status Is Maintained in Non-Smoking Type 2 Diabetic Subjects Without Cardiovascular Disease: Association with Elevated Triglycerides and Large VLDL. <i>Journal of Clinical Medicine</i> , 2020, 9, 49.	2.4	8
34	Vitamin B6, Inflammation, and Cardiovascular Outcome in a Population-Based Cohort: The Prevention of Renal and Vascular End-Stage Disease (PREVEND) Study. <i>Nutrients</i> , 2020, 12, 2711.	4.1	7
35	Plasma C-Peptide and Risk of Developing Type 2 Diabetes in the General Population. <i>Journal of Clinical Medicine</i> , 2020, 9, 3001.	2.4	14
36	HDL Cholesterol Efflux Capacity is Impaired in Severe Short-Term Hypothyroidism Despite Increased HDL Cholesterol. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2020, 105, e3355-e3362.	3.6	13

#	ARTICLE	IF	CITATIONS
37	A Newly Developed Diabetes Risk Index, Based on Lipoprotein Subfractions and Branched Chain Amino Acids, is Associated with Incident Type 2 Diabetes Mellitus in the PREVEND Cohort. <i>Journal of Clinical Medicine</i> , 2020, 9, 2781.	2.4	21
38	Plasma branched chain amino acids are lower in short-term profound hypothyroidism and increase in response to thyroid hormone supplementation. <i>Scandinavian Journal of Clinical and Laboratory Investigation</i> , 2020, 80, 562-566.	1.2	5
39	Characterization of LP-Z Lipoprotein Particles and Quantification in Subjects with Liver Disease Using a Newly Developed NMR-Based Assay. <i>Journal of Clinical Medicine</i> , 2020, 9, 2915.	2.4	18
40	An individual participant data analysis of prospective cohort studies on the association between subclinical thyroid dysfunction and depressive symptoms. <i>Scientific Reports</i> , 2020, 10, 19111.	3.3	19
41	High-Density Lipoprotein Particles and Their Relationship to Posttransplantation Diabetes Mellitus in Renal Transplant Recipients. <i>Biomolecules</i> , 2020, 10, 481.	4.0	9
42	Obesity, adipokines and COVID-19. <i>European Journal of Clinical Investigation</i> , 2020, 50, e13313.	3.4	11
43	Associations Between High-Density Lipoprotein Particles and Ischemic Events by Vascular Domain, Sex, and Ethnicity. <i>Circulation</i> , 2020, 142, 657-669.	1.6	49
44	Oxidative stress is associated with suspected non-alcoholic fatty liver disease and all-cause mortality in the general population. <i>Liver International</i> , 2020, 40, 2148-2159.	3.9	28
45	Lipoprotein insulin resistance score and branched-chain amino acids increase after adrenalectomy for unilateral aldosterone-producing adenoma: a preliminary study. <i>Endocrine</i> , 2020, 68, 420-426.	2.3	1
46	High Plasma Branched-Chain Amino Acids Are Associated with Higher Risk of Post-Transplant Diabetes Mellitus in Renal Transplant Recipients. <i>Journal of Clinical Medicine</i> , 2020, 9, 511.	2.4	6
47	P1650HIGH-DENSITY LIPOPROTEIN PARTICLES AND THEIR RELATIONSHIP TO POSTTRANSPLANTATION DIABETES IN RENAL TRANSPLANT RECIPIENTS. <i>Nephrology Dialysis Transplantation</i> , 2020, 35, .	0.7	0
48	Endogenous urinary glucocorticoid metabolites and mortality in prednisolone-treated renal transplant recipients. <i>Clinical Transplantation</i> , 2020, 34, e13824.	1.6	7
49	Association of Plasma Concentration of Vitamin B ₁₂ With All-Cause Mortality in the General Population in the Netherlands. <i>JAMA Network Open</i> , 2020, 3, e1919274.	5.9	45
50	Ketone Bodies Are Mildly Elevated in Subjects with Type 2 Diabetes Mellitus and Are Inversely Associated with Insulin Resistance as Measured by the Lipoprotein Insulin Resistance Index. <i>Journal of Clinical Medicine</i> , 2020, 9, 321.	2.4	40
51	Plasma ApoE elevations are associated with NAFLD: The PREVEND Study. <i>PLoS ONE</i> , 2019, 14, e0220659.	2.5	11
52	Subclinical thyroid dysfunction and depressive symptoms: protocol for a systematic review and individual participant data meta-analysis of prospective cohort studies. <i>BMJ Open</i> , 2019, 9, e029716.	1.9	3
53	HDL (High-Density Lipoprotein) Cholesterol Efflux Capacity Is Associated With Incident Cardiovascular Disease in the General Population. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2019, 39, 1874-1883.	2.4	78
54	Autoantibodies to Apolipoprotein A-1 as Independent Predictors of Cardiovascular Mortality in Renal Transplant Recipients. <i>Journal of Clinical Medicine</i> , 2019, 8, 948.	2.4	10

#	ARTICLE	IF	CITATIONS
55	Blood Mg ²⁺ is more closely associated with hyperglycaemia than with hypertriacylglycerolaemia: the PREVEND study. <i>Diabetologia</i> , 2019, 62, 1732-1734.	6.3	5
56	Research update for articles published in EJCI in 2017. <i>European Journal of Clinical Investigation</i> , 2019, 49, e13163.	3.4	0
57	High Betaine, a Trimethylamine N-Oxide Related Metabolite, Is Prospectively Associated with Low Future Risk of Type 2 Diabetes Mellitus in the PREVEND Study. <i>Journal of Clinical Medicine</i> , 2019, 8, 1813.	2.4	27
58	Plasma Levels of Retinol Binding Protein 4 Relate to Large VLDL and Small LDL Particles in Subjects with and without Type 2 Diabetes. <i>Journal of Clinical Medicine</i> , 2019, 8, 1792.	2.4	24
59	Prescription of statins in suspected non-alcoholic fatty liver disease and high cardiovascular risk, a population-based study. <i>Liver International</i> , 2019, 39, 1343-1354.	3.9	18
60	Non-Alcoholic Fatty Liver Disease and Risk of Incident Type 2 Diabetes: Role of Circulating Branched-Chain Amino Acids. <i>Nutrients</i> , 2019, 11, 705.	4.1	67
61	Antibodies Against the C-Terminus of ApoA-1 Are Inversely Associated with Cholesterol Efflux Capacity and HDL Metabolism in Subjects with and without Type 2 Diabetes Mellitus. <i>International Journal of Molecular Sciences</i> , 2019, 20, 732.	4.1	11
62	Lower Plasma Magnesium, Measured by Nuclear Magnetic Resonance Spectroscopy, is Associated with Increased Risk of Developing Type 2 Diabetes Mellitus in Women: Results from a Dutch Prospective Cohort Study. <i>Journal of Clinical Medicine</i> , 2019, 8, 169.	2.4	16
63	Application of non-HDL cholesterol for population-based cardiovascular risk stratification: results from the Multinational Cardiovascular Risk Consortium. <i>Lancet, The</i> , 2019, 394, 2173-2183.	13.7	177
64	Higher Sodium Intake Assessed by 24 Hour Urinary Sodium Excretion Is Associated with Non-Alcoholic Fatty Liver Disease: The PREVEND Cohort Study. <i>Journal of Clinical Medicine</i> , 2019, 8, 2157.	2.4	16
65	Lipoprotein insulin resistance index, a high-throughput measure of insulin resistance, is associated with incident type II diabetes mellitus in the Prevention of Renal and Vascular End-Stage Disease study. <i>Journal of Clinical Lipidology</i> , 2019, 13, 129-137.e1.	1.5	31
66	Primary aldosteronism is associated with decreased low-density and high-density lipoprotein particle concentrations and increased GlycA, a pro-inflammatory glycoprotein biomarker. <i>Clinical Endocrinology</i> , 2019, 90, 79-87.	2.4	13
67	Low thyroid function is not associated with an accelerated deterioration in renal function. <i>Nephrology Dialysis Transplantation</i> , 2019, 34, 650-659.	0.7	31
68	Unenhanced CT imaging is highly sensitive to exclude pheochromocytoma: a multicenter study. <i>European Journal of Endocrinology</i> , 2018, 178, 431-437.	3.7	44
69	A novel NMR-based assay to measure circulating concentrations of branched-chain amino acids: Elevation in subjects with type 2 diabetes mellitus and association with carotid intima media thickness. <i>Clinical Biochemistry</i> , 2018, 54, 92-99.	1.9	71
70	The HDL anti-inflammatory function is impaired in the context of low-normal free thyroxine in diabetic and nondiabetic individuals. <i>Clinical Endocrinology</i> , 2018, 88, 752-754.	2.4	4
71	Plasma Branched-Chain Amino Acids and Risk of Incident Type 2 Diabetes: Results from the PREVEND Prospective Cohort Study. <i>Journal of Clinical Medicine</i> , 2018, 7, 513.	2.4	60
72	The Relation Between Thyroid Function and Anemia: A Pooled Analysis of Individual Participant Data. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2018, 103, 3658-3667.	3.6	39

#	ARTICLE	IF	CITATIONS
73	Plasma lecithin:cholesterol acyltransferase and phospholipid transfer protein activity independently associate with nonalcoholic fatty liver disease. <i>European Journal of Clinical Investigation</i> , 2018, 48, e12988.	3.4	17
74	Disturbed Vitamin A Metabolism in Non-Alcoholic Fatty Liver Disease (NAFLD). <i>Nutrients</i> , 2018, 10, 29.	4.1	138
75	Research update for articles published in <sc>EJCI</sc> in 2016. <i>European Journal of Clinical Investigation</i> , 2018, 48, e13016.	3.4	0
76	Emerging role of dopamine in neovascularization of pheochromocytoma and paraganglioma. <i>FASEB Journal</i> , 2017, 31, 2226-2240.	0.5	12
77	Urinary prednisolone excretion is a determinant of serum hepcidin levels in renal transplant recipients. <i>American Journal of Hematology</i> , 2017, 92, E173-E175.	4.1	2
78	High Serum PCSK9 Is Associated With Increased Risk of New-Onset Diabetes After Transplantation in Renal Transplant Recipients. <i>Diabetes Care</i> , 2017, 40, 894-901.	8.6	27
79	Branched Chain Amino Acids Are Associated with Insulin Resistance Independent of Leptin and Adiponectin in Subjects with Varying Degrees of Glucose Tolerance. <i>Metabolic Syndrome and Related Disorders</i> , 2017, 15, 183-186.	1.3	49
80	Downregulation of cholesteryl ester transfer protein by glucocorticoids: a randomised study on <sc>HDL</sc>. <i>European Journal of Clinical Investigation</i> , 2017, 47, 494-503.	3.4	15
81	TMAO is Associated with Mortality: Impact of Modestly Impaired Renal Function. <i>Scientific Reports</i> , 2017, 7, 13781.	3.3	96
82	Calculating the optimal surveillance for head and neck paraganglioma in SDHB-mutation carriers. <i>Familial Cancer</i> , 2017, 16, 123-130.	1.9	30
83	Soluble Vascular Cell Adhesion Molecules May be Protective of Future Cardiovascular Disease Risk: Findings from the PREVEND Prospective Cohort Study. <i>Journal of Atherosclerosis and Thrombosis</i> , 2017, 24, 804-818.	2.0	21
84	Prevalence and determinants of non-alcoholic fatty liver disease in lifelines: A large Dutch population cohort. <i>PLoS ONE</i> , 2017, 12, e0171502.	2.5	74
85	The anti-inflammatory function of HDL is impaired in type 2 diabetes: role of hyperglycemia, paraoxonase-1 and low grade inflammation. <i>Cardiovascular Diabetology</i> , 2017, 16, 132.	6.8	71
86	Somatosensory function in patients with secondary adrenal insufficiency treated with two different doses of hydrocortisoneâ€”Results from a randomized controlled trial. <i>PLoS ONE</i> , 2017, 12, e0180326.	2.5	0
87	GlycA, a marker of acute phase glycoproteins, and the risk of incident type 2 diabetes mellitus: PREVEND study. <i>Clinica Chimica Acta</i> , 2016, 452, 10-17.	1.1	80
88	The predictive value of the antioxidative function of HDL for cardiovascular disease and graft failure in renal transplant recipients. <i>Atherosclerosis</i> , 2016, 249, 181-185.	0.8	12
89	Research update for articles published in <sc>EJCI</sc> in 2014. <i>European Journal of Clinical Investigation</i> , 2016, 46, 880-894.	3.4	2
90	Higher hydrocortisone dose increases bilirubin in hypopituitary patientsâ€”results from an <sc>RCT</sc>. <i>European Journal of Clinical Investigation</i> , 2016, 46, 475-480.	3.4	7

#	ARTICLE	IF	CITATIONS
91	Effects of Hydrocortisone on the Regulation of Blood Pressure: Results From a Randomized Controlled Trial. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2016, 101, 3691-3699.	3.6	34
92	HDL function is impaired in acute myocardial infarction independent of plasma HDL cholesterol levels. <i>Journal of Clinical Lipidology</i> , 2016, 10, 1318-1328.	1.5	50
93	No influence of antihypertensive agents on plasma free metanephrines. <i>Clinical Biochemistry</i> , 2016, 49, 1368-1371.	1.9	10
94	Impaired HDL cholesterol efflux in metabolic syndrome is unrelated to glucose tolerance status: the CODAM study. <i>Scientific Reports</i> , 2016, 6, 27367.	3.3	53
95	Association of hepcidin \leq 25 with survival after kidney transplantation. <i>European Journal of Clinical Investigation</i> , 2016, 46, 994-1001.	3.4	6
96	Dopamine concentration in blood platelets is elevated in patients with head and neck paragangliomas. <i>Clinical Chemistry and Laboratory Medicine</i> , 2016, 54, 1395-401.	2.3	9
97	HDL Cholesterol Efflux Predicts Graft Failure in Renal Transplant Recipients. <i>Journal of the American Society of Nephrology: JASN</i> , 2016, 27, 595-603.	6.1	71
98	Mass spectrometric quantification of salivary metanephrines—A study in healthy subjects. <i>Clinical Biochemistry</i> , 2016, 49, 983-988.	1.9	10
99	Effect of Metformin Treatment on Lipoprotein Subfractions in Non-Diabetic Patients with Acute Myocardial Infarction: A Glycometabolic Intervention as Adjunct to Primary Coronary Intervention in ST Elevation Myocardial Infarction (GIPS-III) Trial. <i>PLoS ONE</i> , 2016, 11, e0145719.	2.5	13
100	Circulating Total Bilirubin and Risk of Incident Cardiovascular Disease in the General Population. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2015, 35, 716-724.	2.4	96
101	Increased Risk of Atrial Fibrillation After Treatment for Differentiated Thyroid Carcinoma. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2015, 100, 4563-4569.	3.6	39
102	The effects of two different doses of hydrocortisone on cognition in patients with secondary adrenal insufficiency — Results from a randomized controlled trial. <i>Psychoneuroendocrinology</i> , 2015, 55, 36-47.	2.7	23
103	The Relationship of the Anti-Oxidant Bilirubin with Free Thyroxine Is Modified by Insulin Resistance in Euthyroid Subjects. <i>PLoS ONE</i> , 2014, 9, e90886.	2.5	21
104	Genome Wide Association Identifies Common Variants at the SERPINA6/SERPINA1 Locus Influencing Plasma Cortisol and Corticosteroid Binding Globulin. <i>PLoS Genetics</i> , 2014, 10, e1004474.	3.5	105
105	The HDL anti-inflammatory function is impaired in myocardial infarction and may predict new cardiac events independent of HDL cholesterol. <i>Clinica Chimica Acta</i> , 2014, 433, 34-38.	1.1	32
106	Skin Autofluorescence Is Associated With 5-Year Mortality and Cardiovascular Events in Patients With Peripheral Artery Disease. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2014, 34, 933-938.	2.4	78
107	Pleiotropic Effects of Lipid Genes on Plasma Glucose, HbA1c, and HOMA-IR Levels. <i>Diabetes</i> , 2014, 63, 3149-3158.	0.6	61
108	Combined Effects of Smoking and Alcohol on Metabolic Syndrome: The LifeLines Cohort Study. <i>PLoS ONE</i> , 2014, 9, e96406.	2.5	73

#	ARTICLE	IF	CITATIONS
109	Hdl Cholesterol Efflux Predicts Graft Failure But Not Cardiovascular And Overall Mortality In Renal Transplant Recipients. , 2014, , .		0
110	Role of HDL Cholesterol and Estimates of HDL Particle Composition in Future Development of Type 2 Diabetes in the General Population: The PREVEND Study. Journal of Clinical Endocrinology and Metabolism, 2013, 98, E1352-E1359.	3.6	98
111	Assessing the functional properties of high-density lipoproteins: an emerging concept in cardiovascular research. Biomarkers in Medicine, 2013, 7, 457-472.	1.4	62
112	The Interaction of Age and Type 2 Diabetes on Executive Function and Memory in Persons Aged 35 Years or Older. PLoS ONE, 2013, 8, e82991.	2.5	22
113	PS4 - 22. HDL-cholesterol, Apolipoprotein A-I/A-II, and HDL-cholesterol particle composition for the risk of developing type 2 diabetes in the community: the PREVEND Study. Nederlands Tijdschrift Voor Diabetologie, 2012, 10, 113-113.	0.0	0
114	Research update for articles published in EJCI in 2009. European Journal of Clinical Investigation, 2011, 41, 1149-1163.	3.4	0
115	Genetic Variation at the <i>Phospholipid Transfer Protein</i> Locus Affects Its Activity and High-Density Lipoprotein Size and Is a Novel Marker of Cardiovascular Disease Susceptibility. Circulation, 2010, 122, 470-477.	1.6	86
116	High plasma C-reactive protein (CRP) is related to low paraoxonase (PON) activity independently of high leptin and low adiponectin in type 2 diabetes mellitus. Clinical Endocrinology, 2009, 70, 221-226.	2.4	52
117	Glucocorticoid replacement is associated with hypertriglyceridaemia, elevated glucose and higher non-HDL cholesterol and may diminish the association of HDL cholesterol with the <i>ApoA</i> CETP promoter polymorphism in GH-receiving hypopituitary patients. Clinical Endocrinology, 2008, 69, 359-366.	2.4	14
118	Common variation in the CETP gene and the implications for cardiovascular disease and its treatment: an updated analysis. Pharmacogenomics, 2008, 9, 747-763.	1.3	48
119	Renin-angiotensin-aldosterone responsiveness to low sodium and blood pressure reactivity to angiotensin-II are unrelated to cholesteryl ester transfer protein mass in healthy subjects. Expert Opinion on Therapeutic Targets, 2008, 12, 1321-1328.	3.4	4
120	Plasma Lecithin: Cholesterol Acyltransferase Activity Is Elevated in Metabolic Syndrome and Is an Independent Marker of Increased Carotid Artery Intima Media Thickness. Journal of Clinical Endocrinology and Metabolism, 2008, 93, 4860-4866.	3.6	54
121	High plasma cholesteryl ester transfer protein levels may favour reduced incidence of cardiovascular events in men with low triglycerides. European Heart Journal, 2007, 28, 1012-1018.	2.2	68
122	Lower plasma adiponectin is a marker of increased intima-media thickness associated with type 2 diabetes mellitus and with male gender. European Journal of Endocrinology, 2007, 156, 387-394.	3.7	54
123	Carotid artery intima media thickness is inversely related to serum free thyroxine in euthyroid subjects. Clinical Endocrinology, 2007, 67, 668-673.	2.4	67
124	The Effect of Cholesteryl Ester Transfer Protein A Promoter Polymorphism on High-Density Lipoprotein Cholesterol Is Dependent on Serum Triglycerides. Journal of Clinical Endocrinology and Metabolism, 2005, 90, 4198-4204.	3.6	67