

Josep Maria Haro

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4303299/publications.pdf>

Version: 2024-02-01

671
papers

99,466
citations

1531

109
h-index

324

295
g-index

724
all docs

724
docs citations

724
times ranked

120049
citing authors

#	ARTICLE	IF	CITATIONS
1	Disability-adjusted life years (DALYs) for 291 diseases and injuries in 21 regions, 1990â€“2010: a systematic analysis for the Global Burden of Disease Study 2010. Lancet, The, 2012, 380, 2197-2223.	6.3	7,061
2	Years lived with disability (YLDs) for 1160 sequelae of 289 diseases and injuries 1990â€“2010: a systematic analysis for the Global Burden of Disease Study 2010. Lancet, The, 2012, 380, 2163-2196.	6.3	6,376
3	Global, regional, and national ageâ€“sex specific all-cause and cause-specific mortality for 240 causes of death, 1990â€“2013: a systematic analysis for the Global Burden of Disease Study 2013. Lancet, The, 2015, 385, 117-171.	6.3	5,847
4	Global, regional, and national incidence, prevalence, and years lived with disability for 328 diseases and injuries for 195 countries, 1990â€“2016: a systematic analysis for the Global Burden of Disease Study 2016. Lancet, The, 2017, 390, 1211-1259.	6.3	5,578
5	Global, regional, and national incidence, prevalence, and years lived with disability for 301 acute and chronic diseases and injuries in 188 countries, 1990â€“2013: a systematic analysis for the Global Burden of Disease Study 2013. Lancet, The, 2015, 386, 743-800.	6.3	4,951
6	Prevalence, Severity, and Unmet Need for Treatment of Mental Disorders in the World Health Organization World Mental Health Surveys. JAMA - Journal of the American Medical Association, 2004, 291, 2581.	3.8	2,696
7	Global, regional, and national burden of neurological disorders, 1990â€“2016: a systematic analysis for the Global Burden of Disease Study 2016. Lancet Neurology, The, 2019, 18, 459-480.	4.9	2,625
8	Global, regional, and national comparative risk assessment of 79 behavioural, environmental and occupational, and metabolic risks or clusters of risks in 188 countries, 1990â€“2013: a systematic analysis for the Global Burden of Disease Study 2013. Lancet, The, 2015, 386, 2287-2323.	6.3	2,184
9	Alcohol use and burden for 195 countries and territories, 1990â€“2016: a systematic analysis for the Global Burden of Disease Study 2016. Lancet, The, 2018, 392, 1015-1035.	6.3	2,005
10	Cross-national prevalence and risk factors for suicidal ideation, plans and attempts. British Journal of Psychiatry, 2008, 192, 98-105.	1.7	1,993
11	Childhood adversities and adult psychopathology in the WHO World Mental Health Surveys. British Journal of Psychiatry, 2010, 197, 378-385.	1.7	1,854
12	Global, Regional, and National Cancer Incidence, Mortality, Years of Life Lost, Years Lived With Disability, and Disability-Adjusted Life-Years for 29 Cancer Groups, 1990 to 2017. JAMA Oncology, 2019, 5, 1749.	3.4	1,691
13	Global, regional, and national disability-adjusted life years (DALYs) for 306 diseases and injuries and healthy life expectancy (HALE) for 188 countries, 1990â€“2013: quantifying the epidemiological transition. Lancet, The, 2015, 386, 2145-2191.	6.3	1,544
14	Global, regional, and national burden of Alzheimer's disease and other dementias, 1990â€“2016: a systematic analysis for the Global Burden of Disease Study 2016. Lancet Neurology, The, 2019, 18, 88-106.	4.9	1,512
15	Use of mental health services for anxiety, mood, and substance disorders in 17 countries in the WHO world mental health surveys. Lancet, The, 2007, 370, 841-850.	6.3	1,271
16	Lifetime prevalence and age-of-onset distributions of mental disorders in the World Health Organization's World Mental Health Survey Initiative. World Psychiatry, 2007, 6, 168-76.	4.8	1,239
17	Cross-national prevalence and correlates of adult attention-deficit hyperactivity disorder. British Journal of Psychiatry, 2007, 190, 402-409.	1.7	1,118
18	Global, regional, and national burden of traumatic brain injury and spinal cord injury, 1990â€“2016: a systematic analysis for the Global Burden of Disease Study 2016. Lancet Neurology, The, 2019, 18, 56-87.	4.9	1,064

#	ARTICLE	IF	CITATIONS
19	Prevalence of mental disorders in Europe: results from the European Study of the Epidemiology of Mental Disorders (ESEMEd) project. <i>Acta Psychiatrica Scandinavica</i> , 2004, 109, 21-27.	2.2	1,053
20	The Prevalence and Correlates of Binge Eating Disorder in the World Health Organization World Mental Health Surveys. <i>Biological Psychiatry</i> , 2013, 73, 904-914.	0.7	1,025
21	Cross-National Associations Between Gender and Mental Disorders in the World Health Organization World Mental Health Surveys. <i>Archives of General Psychiatry</i> , 2009, 66, 785.	13.8	1,018
22	Concordance of the Composite International Diagnostic Interview Version 3.0 (CIDI 3.0) with standardized clinical assessments in the WHO World Mental Health Surveys. <i>International Journal of Methods in Psychiatric Research</i> , 2006, 15, 167-180.	1.1	949
23	Common Chronic Pain Conditions in Developed and Developing Countries: Gender and Age Differences and Comorbidity With Depression-Anxiety Disorders. <i>Journal of Pain</i> , 2008, 9, 883-891.	0.7	918
24	The global burden of disease attributable to alcohol and drug use in 195 countries and territories, 1990-2016: a systematic analysis for the Global Burden of Disease Study 2016. <i>Lancet Psychiatry</i> , the, 2018, 5, 987-1012.	3.7	885
25	Mental disorders among college students in the World Health Organization World Mental Health Surveys. <i>Psychological Medicine</i> , 2016, 46, 2955-2970.	2.7	866
26	Trauma and PTSD in the WHO World Mental Health Surveys. <i>HÅrre Utbildning</i> , 2017, 8, 1353383.	1.4	801
27	Global, regional, and national incidence and mortality for HIV, tuberculosis, and malaria during 1990-2013: a systematic analysis for the Global Burden of Disease Study 2013. <i>Lancet, The</i> , 2014, 384, 1005-1070.	6.3	786
28	Posttraumatic stress disorder in the World Mental Health Surveys. <i>Psychological Medicine</i> , 2017, 47, 2260-2274.	2.7	675
29	Undertreatment of people with major depressive disorder in 21 countries. <i>British Journal of Psychiatry</i> , 2017, 210, 119-124.	1.7	664
30	Cross-National Analysis of the Associations among Mental Disorders and Suicidal Behavior: Findings from the WHO World Mental Health Surveys. <i>PLoS Medicine</i> , 2009, 6, e1000123.	3.9	592
31	The association between social relationships and depression: A systematic review. <i>Journal of Affective Disorders</i> , 2015, 175, 53-65.	2.0	552
32	The Clinical Global Impression-Schizophrenia scale: a simple instrument to measure the diversity of symptoms present in schizophrenia. <i>Acta Psychiatrica Scandinavica</i> , 2003, 107, 16-23.	2.2	508
33	The descriptive epidemiology of DSM-IV Adult ADHD in the World Health Organization World Mental Health Surveys. <i>ADHD Attention Deficit and Hyperactivity Disorders</i> , 2017, 9, 47-65.	1.7	507
34	Healthcare Access and Quality Index based on mortality from causes amenable to personal health care in 195 countries and territories, 1990-2015: a novel analysis from the Global Burden of Disease Study 2015. <i>Lancet, The</i> , 2017, 390, 231-266.	6.3	480
35	Mental disorders among persons with chronic back or neck pain: Results from the world mental health surveys. <i>Pain</i> , 2007, 129, 332-342.	2.0	471
36	Risk factors for suicidality in Europe: Results from the ESEMED study. <i>Journal of Affective Disorders</i> , 2007, 101, 27-34.	2.0	465

#	ARTICLE	IF	CITATIONS
37	Estimates of global, regional, and national incidence, prevalence, and mortality of HIV, 1980–2015: the Global Burden of Disease Study 2015. <i>Lancet HIV</i> , 2016, 3, e361-e387.	2.1	461
38	Depression–anxiety relationships with chronic physical conditions: Results from the World Mental Health surveys. <i>Journal of Affective Disorders</i> , 2007, 103, 113-120.	2.0	414
39	Development of Lifetime Comorbidity in the World Health Organization World Mental Health Surveys. <i>Archives of General Psychiatry</i> , 2011, 68, 90.	13.8	410
40	Global Multimorbidity Patterns: A Cross-Sectional, Population-Based, Multi-Country Study. <i>Journals of Gerontology - Series A Biological Sciences and Medical Sciences</i> , 2016, 71, 205-214.	1.7	404
41	The global, regional, and national burden of stomach cancer in 195 countries, 1990–2017: a systematic analysis for the Global Burden of Disease study 2017. <i>The Lancet Gastroenterology and Hepatology</i> , 2020, 5, 42-54.	3.7	390
42	Psychotic Experiences in the General Population. <i>JAMA Psychiatry</i> , 2015, 72, 697.	6.0	387
43	The epidemiology of eating disorders in six European countries: Results of the ESEMeD-WMH project. <i>Journal of Psychiatric Research</i> , 2009, 43, 1125-1132.	1.5	373
44	Age differences in the prevalence and co-morbidity of DSM-IV major depressive episodes: results from the WHO World Mental Health Survey Initiative. <i>Depression and Anxiety</i> , 2010, 27, 351-364.	2.0	363
45	Days out of role due to common physical and mental conditions: results from the WHO World Mental Health surveys. <i>Molecular Psychiatry</i> , 2011, 16, 1234-1246.	4.1	360
46	Predictors and clinical consequences of non-adherence with antipsychotic medication in the outpatient treatment of schizophrenia. <i>Psychiatry Research</i> , 2010, 176, 109-113.	1.7	359
47	Delay and failure in treatment seeking after first onset of mental disorders in the World Health Organization's World Mental Health Survey Initiative. <i>World Psychiatry</i> , 2007, 6, 177-85.	4.8	351
48	Obesity and mental disorders in the general population: results from the world mental health surveys. <i>International Journal of Obesity</i> , 2008, 32, 192-200.	1.6	348
49	Childhood Predictors of Adult Attention-Deficit/Hyperactivity Disorder: Results from the World Health Organization World Mental Health Survey Initiative. <i>Biological Psychiatry</i> , 2009, 65, 46-54.	0.7	346
50	Use of mental health services in Europe: results from the European Study of the Epidemiology of Mental Disorders (ESEMeD) project. <i>Acta Psychiatrica Scandinavica</i> , 2004, 109, 47-54.	2.2	333
51	CUMULATIVE TRAUMAS AND RISK THRESHOLDS: 12-MONTH PTSD IN THE WORLD MENTAL HEALTH (WMH) SURVEYS. <i>Depression and Anxiety</i> , 2014, 31, 130-142.	2.0	329
52	Disability and treatment of specific mental and physical disorders across the world. <i>British Journal of Psychiatry</i> , 2008, 192, 368-375.	1.7	326
53	Treatment gap for anxiety disorders is global: Results of the World Mental Health Surveys in 21 countries. <i>Depression and Anxiety</i> , 2018, 35, 195-208.	2.0	318
54	Assessing depression in primary care with the PHQ-9: Can it be carried out over the telephone?. <i>Journal of General Internal Medicine</i> , 2005, 20, 738-742.	1.3	298

#	ARTICLE	IF	CITATIONS
55	Disability and quality of life impact of mental disorders in Europe: results from the European Study of the Epidemiology of Mental Disorders (ESEMED) project. <i>Acta Psychiatrica Scandinavica</i> , 2004, 109, 38-46.	2.2	285
56	Association of Childhood Adversities and Early-Onset Mental Disorders With Adult-Onset Chronic Physical Conditions. <i>Archives of General Psychiatry</i> , 2011, 68, 838.	13.8	281
57	Childhood adversities as risk factors for onset and persistence of suicidal behaviour. <i>British Journal of Psychiatry</i> , 2010, 197, 20-27.	1.7	278
58	The role of socio-economic status in depression: results from the COURAGE (aging survey in Europe). <i>BMC Public Health</i> , 2016, 16, 1098.	1.2	278
59	Loneliness and depression in the elderly: the role of social network. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2017, 52, 381-390.	1.6	274
60	Cross-sectional Comparison of the Epidemiology of <i>DSM-5</i> Generalized Anxiety Disorder Across the Globe. <i>JAMA Psychiatry</i> , 2017, 74, 465.	6.0	274
61	Socio-economic variations in the mental health treatment gap for people with anxiety, mood, and substance use disorders: results from the WHO World Mental Health (WMH) surveys. <i>Psychological Medicine</i> , 2018, 48, 1560-1571.	2.7	274
62	The prevalence and effects of adult attention-deficit/hyperactivity disorder (ADHD) on the performance of workers: results from the WHO World Mental Health Survey Initiative. <i>Occupational and Environmental Medicine</i> , 2008, 65, 835-842.	1.3	262
63	The relation between multiple pains and mental disorders: Results from the World Mental Health Surveys. <i>Pain</i> , 2008, 135, 82-91.	2.0	260
64	The cross-national epidemiology of social anxiety disorder: Data from the World Mental Health Survey Initiative. <i>BMC Medicine</i> , 2017, 15, 143.	2.3	258
65	Anxiety, depression, loneliness and social network in the elderly: Longitudinal associations from The Irish Longitudinal Study on Ageing (TILDA). <i>Journal of Affective Disorders</i> , 2019, 246, 82-88.	2.0	252
66	Mental disorders among adults with asthma: results from the World Mental Health Survey. <i>General Hospital Psychiatry</i> , 2007, 29, 123-133.	1.2	251
67	Dissociation in Posttraumatic Stress Disorder: Evidence from the World Mental Health Surveys. <i>Biological Psychiatry</i> , 2013, 73, 302-312.	0.7	251
68	Mental-physical co-morbidity and its relationship with disability: results from the World Mental Health Surveys. <i>Psychological Medicine</i> , 2009, 39, 33-43.	2.7	239
69	Population level of unmet need for mental healthcare in Europe. <i>British Journal of Psychiatry</i> , 2007, 190, 299-306.	1.7	222
70	Main traumatic events in Europe: PTSD in the European study of the epidemiology of mental disorders survey. <i>Journal of Traumatic Stress</i> , 2008, 21, 455-462.	1.0	218
71	Mental disorders among persons with heart disease – results from World Mental Health surveys. <i>General Hospital Psychiatry</i> , 2007, 29, 325-334.	1.2	204
72	Are loneliness and social isolation associated with cognitive decline?. <i>International Journal of Geriatric Psychiatry</i> , 2019, 34, 1613-1622.	1.3	202

#	ARTICLE	IF	CITATIONS
73	Parent psychopathology and offspring mental disorders: Results from the WHO World Mental Health Surveys. <i>British Journal of Psychiatry</i> , 2012, 200, 290-299.	1.7	201
74	How well can post-traumatic stress disorder be predicted from pre-trauma risk factors? An exploratory study in the WHO World Mental Health Surveys. <i>World Psychiatry</i> , 2014, 13, 265-274.	4.8	194
75	Comparison of Population Health Status in Six European Countries. <i>Medical Care</i> , 2009, 47, 255-261.	1.1	191
76	Global Mortality From Firearms, 1990-2016. <i>JAMA - Journal of the American Medical Association</i> , 2018, 320, 792.	3.8	189
77	The Mental Component of the Short-Form 12 Health Survey (SF-12) as a Measure of Depressive Disorders in the General Population: Results with Three Alternative Scoring Methods. <i>Value in Health</i> , 2013, 16, 564-573.	0.1	187
78	Association of perceived stigma and mood and anxiety disorders: results from the World Mental Health Surveys. <i>Acta Psychiatrica Scandinavica</i> , 2008, 118, 305-314.	2.2	182
79	Factors associated with skeletal muscle mass, sarcopenia, and sarcopenic obesity in older adults: a multi-continent study. <i>Journal of Cachexia, Sarcopenia and Muscle</i> , 2016, 7, 312-321.	2.9	182
80	Association of <i>DSM-IV</i> Posttraumatic Stress Disorder With Traumatic Experience Type and History in the World Health Organization World Mental Health Surveys. <i>JAMA Psychiatry</i> , 2017, 74, 270.	6.0	179
81	Chronic Physical Conditions and Their Association With First Onset of Suicidal Behavior in the World Mental Health Surveys. <i>Psychosomatic Medicine</i> , 2010, 72, 712-719.	1.3	178
82	The Bidirectional Associations Between Psychotic Experiences and DSM-IV Mental Disorders. <i>American Journal of Psychiatry</i> , 2016, 173, 997-1006.	4.0	176
83	Development of EMPRO: A Tool for the Standardized Assessment of Patient-Reported Outcome Measures. <i>Value in Health</i> , 2008, 11, 700-708.	0.1	166
84	Chronic Conditions and Sleep Problems among Adults Aged 50 years or over in Nine Countries: A Multi-Country Study. <i>PLoS ONE</i> , 2014, 9, e114742.	1.1	163
85	Mapping 123 million neonatal, infant and child deaths between 2000 and 2017. <i>Nature</i> , 2019, 574, 353-358.	13.7	161
86	Estimating treatment coverage for people with substance use disorders: an analysis of data from the World Mental Health Surveys. <i>World Psychiatry</i> , 2017, 16, 299-307.	4.8	160
87	Mental health research priorities for Europe. <i>Lancet Psychiatry</i> , 2015, 2, 1036-1042.	3.7	158
88	12-Month comorbidity patterns and associated factors in Europe: results from the European Study of the Epidemiology of Mental Disorders (ESEMED) project. <i>Acta Psychiatrica Scandinavica</i> , 2004, 109, 28-37.	2.2	157
89	Associations of serious mental illness with earnings: results from the WHO World Mental Health surveys. <i>British Journal of Psychiatry</i> , 2010, 197, 114-121.	1.7	156
90	Using Smartphones and Wearable Devices to Monitor Behavioral Changes During COVID-19. <i>Journal of Medical Internet Research</i> , 2020, 22, e19992.	2.1	155

#	ARTICLE	IF	CITATIONS
91	Mental disorders among persons with diabetesâ€”Results from the World Mental Health Surveys. <i>Journal of Psychosomatic Research</i> , 2008, 65, 571-580.	1.2	154
92	Perceived stigma among individuals with common mental disorders. <i>Journal of Affective Disorders</i> , 2009, 118, 180-186.	2.0	152
93	Stratified medicine for mental disorders. <i>European Neuropsychopharmacology</i> , 2014, 24, 5-50.	0.3	152
94	Cross-national clinical and functional remission rates: Worldwide Schizophrenia Outpatient Health Outcomes (W-SOHO) study. <i>British Journal of Psychiatry</i> , 2011, 199, 194-201.	1.7	148
95	Treatment adequacy for anxiety and depressive disorders in six European countries. <i>British Journal of Psychiatry</i> , 2007, 190, 172-173.	1.7	147
96	Rates and predictors of remission and recovery during 3â€”years in 392 neverâ€”treated patients with schizophrenia. <i>Acta Psychiatrica Scandinavica</i> , 2008, 118, 220-229.	2.2	145
97	Cross-national epidemiology of panic disorder and panic attacks in the world mental health surveys. <i>Depression and Anxiety</i> , 2016, 33, 1155-1177.	2.0	144
98	Psychotropic drug utilization in Europe: results from the European Study of the Epidemiology of Mental Disorders (ESEMeD) project. <i>Acta Psychiatrica Scandinavica</i> , 2004, 109, 55-64.	2.2	143
99	Bullying Victimization and Suicide Attempt Among Adolescents Aged 12â€”15 Years From 48 Countries. <i>Journal of the American Academy of Child and Adolescent Psychiatry</i> , 2019, 58, 907-918.e4.	0.3	140
100	Prevalence of mental disorders in primary care: results from the diagnosis and treatment of mental disorders in primary care study (DASMAP). <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2010, 45, 201-210.	1.6	137
101	Sampling and methods of the European Study of the Epidemiology of Mental Disorders (ESEMeD) project. <i>Acta Psychiatrica Scandinavica</i> , 2004, 109, 8-20.	2.2	136
102	Gender and the relationship between marital status and first onset of mood, anxiety and substance use disorders. <i>Psychological Medicine</i> , 2010, 40, 1495-1505.	2.7	133
103	Mental health impact of the first wave of COVID-19 pandemic on Spanish healthcare workers: A large cross-sectional survey. <i>Revista De Psiquiatr�a Y Salud Mental</i> , 2021, 14, 90-105.	1.0	133
104	Mental disorders and termination of education in high-income and low- and middle-income countries: epidemiological study. <i>British Journal of Psychiatry</i> , 2009, 194, 411-417.	1.7	129
105	The GERAS Study: A Prospective Observational Study of Costs and Resource Use in Community Dwellers with Alzheimer's Disease in Three European Countries â€” Study Design and Baseline Findings. <i>Journal of Alzheimer's Disease</i> , 2013, 36, 385-399.	1.2	129
106	Clinical factors influencing the prescription of antidepressants and benzodiazepines. <i>Journal of Affective Disorders</i> , 2008, 110, 84-93.	2.0	128
107	Impact of Multimorbidity on Disability and Quality of Life in the Spanish Older Population. <i>PLoS ONE</i> , 2014, 9, e111498.	1.1	122
108	Loneliness, Social Networks, and Health: A Cross-Sectional Study in Three Countries. <i>PLoS ONE</i> , 2016, 11, e0145264.	1.1	121

#	ARTICLE	IF	CITATIONS
109	Social relationships, loneliness, and mental health among older men and women in Ireland: A prospective community-based study. <i>Journal of Affective Disorders</i> , 2016, 204, 59-69.	2.0	117
110	Research priorities for public mental health in Europe: recommendations of the ROAMER project. <i>European Journal of Public Health</i> , 2015, 25, 249-254.	0.1	113
111	La carga de enfermedad en España: resultados del Estudio de la Carga Global de las Enfermedades 2016. <i>Medicina Clínica</i> , 2018, 151, 171-190.	0.3	113
112	Remission and Relapse in the Outpatient Care of Schizophrenia. <i>Journal of Clinical Psychopharmacology</i> , 2006, 26, 571-578.	0.7	112
113	Recovery in the outpatient setting: 36-month results from the Schizophrenia Outpatients Health Outcomes (SOHO) study. <i>Schizophrenia Research</i> , 2009, 108, 223-230.	1.1	111
114	The state of the art in European research on reducing social exclusion and stigma related to mental health: A systematic mapping of the literature. <i>European Psychiatry</i> , 2014, 29, 381-389.	0.1	110
115	Post-traumatic stress disorder associated with natural and human-made disasters in the World Mental Health Surveys. <i>Psychological Medicine</i> , 2017, 47, 227-241.	2.7	110
116	ROAMER: roadmap for mental health research in Europe. <i>International Journal of Methods in Psychiatric Research</i> , 2014, 23, 1-14.	1.1	109
117	The association between childhood adversities and subsequent first onset of psychotic experiences: a cross-national analysis of 23 998 respondents from 17 countries. <i>Psychological Medicine</i> , 2017, 47, 1230-1245.	2.7	108
118	Three-year antipsychotic effectiveness in the outpatient care of schizophrenia: Observational versus randomized studies results. <i>European Neuropsychopharmacology</i> , 2007, 17, 235-244.	0.3	106
119	Built Environment and Elderly Population Health: A Comprehensive Literature Review. <i>Clinical Practice and Epidemiology in Mental Health</i> , 2014, 10, 103-115.	0.6	106
120	Obsessive-compulsive symptom dimensions in the general population: Results from an epidemiological study in six European countries. <i>Journal of Affective Disorders</i> , 2010, 124, 291-299.	2.0	105
121	Differences in Lifetime Use of Services for Mental Health Problems in Six European Countries. <i>Psychiatric Services</i> , 2007, 58, 213-220.	1.1	103
122	Gender differences in response to antipsychotic treatment in outpatients with schizophrenia. <i>Psychiatry Research</i> , 2007, 153, 225-231.	1.7	103
123	Associations between mental disorders and subsequent onset of hypertension. <i>General Hospital Psychiatry</i> , 2014, 36, 142-149.	1.2	102
124	The Burden of Mental Disorders in the Eastern Mediterranean Region, 1990-2013. <i>PLoS ONE</i> , 2017, 12, e0169575.	1.1	102
125	Remote assessment of disease and relapse in major depressive disorder (RADAR-MDD): a multi-centre prospective cohort study protocol. <i>BMC Psychiatry</i> , 2019, 19, 72.	1.1	99
126	The European Schizophrenia Outpatient Health Outcomes (SOHO) study: rationale, methods and recruitment. <i>Acta Psychiatrica Scandinavica</i> , 2003, 107, 222-232.	2.2	98

#	ARTICLE	IF	CITATIONS
127	Epidemiology of major depressive episode in a southern European country: Results from the ESEMeD-Spain project. <i>Journal of Affective Disorders</i> , 2010, 120, 76-85.	2.0	98
128	Health status of the advanced elderly in six european countries: results from a representative survey using EQ-5D and SF-12. <i>Health and Quality of Life Outcomes</i> , 2010, 8, 143.	1.0	98
129	Clinical and economic consequences of medication non-adherence in the treatment of patients with a manic/mixed episode of bipolar disorder: Results from the European Mania in Bipolar Longitudinal Evaluation of Medication (EMBLEM) Study. <i>Psychiatry Research</i> , 2011, 190, 110-114.	1.7	96
130	Childhood Adversity, Early-Onset Depressive/Anxiety Disorders, and Adult-Onset Asthma. <i>Psychosomatic Medicine</i> , 2008, 70, 1035-1043.	1.3	91
131	Validation of an instrument to evaluate quality of life in the aging population: WHOQOL-AGE. <i>Health and Quality of Life Outcomes</i> , 2013, 11, 177.	1.0	89
132	Population prevalence of edentulism and its association with depression and self-rated health. <i>Scientific Reports</i> , 2016, 6, 37083.	1.6	89
133	Negative perceptions of ageing predict the onset and persistence of depression and anxiety: Findings from a prospective analysis of the Irish Longitudinal Study on Ageing (TILDA). <i>Journal of Affective Disorders</i> , 2016, 199, 132-138.	2.0	88
134	Barriers to and Facilitators of Engagement With mHealth Technology for Remote Measurement and Management of Depression: Qualitative Analysis. <i>JMIR MHealth and UHealth</i> , 2019, 7, e11325.	1.8	88
135	The European Study of the Epidemiology of Mental Disorders (ESEMeD/MHEDEA 2000) project: rationale and methods. <i>International Journal of Methods in Psychiatric Research</i> , 2002, 11, 55-67.	1.1	85
136	Effectiveness of antipsychotic treatment for schizophrenia: 6-month results of the Pan-European Schizophrenia Outpatient Health Outcomes (SOHO) study. <i>Acta Psychiatrica Scandinavica</i> , 2005, 111, 220-231.	2.2	83
137	Long-term Course of Hospitalization for Schizophrenia: Part II. Change With Passage of Time. <i>Schizophrenia Bulletin</i> , 1992, 18, 229-241.	2.3	82
138	Relationship of insight with medication adherence and the impact on outcomes in patients with schizophrenia and bipolar disorder: results from a 1-year European outpatient observational study. <i>BMC Psychiatry</i> , 2015, 15, 189.	1.1	82
139	Trauma and psychotic experiences: transnational data from the World Mental Health Survey. <i>British Journal of Psychiatry</i> , 2017, 211, 373-380.	1.7	82
140	Latent class analysis of multimorbidity patterns and associated outcomes in Spanish older adults: a prospective cohort study. <i>BMC Geriatrics</i> , 2017, 17, 186.	1.1	82
141	The World Health Organization World Mental Health Survey Initiative. <i>Epidemiologia E Psichiatria Sociale</i> , 2006, 15, 161-166.	1.0	81
142	Mental disorders among persons with arthritis: results from the World Mental Health Surveys. <i>Psychological Medicine</i> , 2008, 38, 1639-1650.	2.7	81
143	Is major depression adequately diagnosed and treated by general practitioners? Results from an epidemiological study. <i>General Hospital Psychiatry</i> , 2010, 32, 201-209.	1.2	81
144	Factors associated with suicidal ideation and attempts in Spain for different age groups. Prevalence before and after the onset of the economic crisis. <i>Journal of Affective Disorders</i> , 2014, 163, 1-9.	2.0	81

#	ARTICLE	IF	CITATIONS
145	Gender differences in response to antidepressant treatment prescribed in primary care. Does menopause make a difference?. <i>Journal of Affective Disorders</i> , 2006, 93, 53-60.	2.0	79
146	Depression comorbid with tuberculosis and its impact on health status: cross-sectional analysis of community-based data from 48 low- and middle-income countries. <i>BMC Medicine</i> , 2017, 15, 209.	2.3	79
147	Suicidal thoughts and behaviors among college students and same-aged peers: results from the World Health Organization World Mental Health Surveys. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2018, 53, 279-288.	1.6	79
148	Comparison of Different Valuation Methods for Population Health Status Measured by the EQ-5D in Three European Countries. <i>Value in Health</i> , 2009, 12, 750-758.	0.1	78
149	Psychometric properties of the twelve item World Health Organization Disability Assessment Schedule II (WHO-DAS II) in Spanish primary care patients with a first major depressive episode. <i>Journal of Affective Disorders</i> , 2010, 121, 52-58.	2.0	78
150	Met and Unmet Needs of Schizophrenia Patients in a Spanish Sample. <i>Schizophrenia Bulletin</i> , 2003, 29, 201-210.	2.3	77
151	Role of common mental and physical disorders in partial disability around the world. <i>British Journal of Psychiatry</i> , 2012, 200, 454-461.	1.7	77
152	Caregiver Burden in Alzheimer's Disease: Differential Associations in Adult-Child and Spousal Caregivers in the GERAS Observational Study. <i>Dementia and Geriatric Cognitive Disorders Extra</i> , 2014, 4, 51-64.	0.6	77
153	Advanced analytical methodologies for measuring healthy ageing and its determinants, using factor analysis and machine learning techniques: the ATHLOS project. <i>Scientific Reports</i> , 2017, 7, 43955.	1.6	77
154	Family burden related to mental and physical disorders in the world: results from the WHO World Mental Health (WMH) surveys. <i>Revista Brasileira De Psiquiatria</i> , 2013, 35, 115-125.	0.9	76
155	Analysis of burden in caregivers of people with Alzheimer's disease using self-report and supervision hours. <i>Journal of Nutrition, Health and Aging</i> , 2014, 18, 677-684.	1.5	76
156	Neurocognitive performance and negative symptoms: Are they equal in explaining disability in schizophrenia outpatients?. <i>Schizophrenia Research</i> , 2006, 87, 246-253.	1.1	75
157	Evidence for three distinct classes of 'typical', 'psychotic' and 'dual' mania: results from the EMBLEM study. <i>Acta Psychiatrica Scandinavica</i> , 2006, 113, 112-120.	2.2	75
158	Association Between Psychotic Experiences and Subsequent Suicidal Thoughts and Behaviors. <i>JAMA Psychiatry</i> , 2017, 74, 1136.	6.0	75
159	Food insecurity (hunger) and suicide attempts among 179,771 adolescents attending school from 9 high-income, 31 middle-income, and 4 low-income countries: A cross-sectional study. <i>Journal of Affective Disorders</i> , 2019, 248, 91-98.	2.0	75
160	The SOHO (Schizophrenia Outpatient Health Outcome) Study. <i>CNS Drugs</i> , 2006, 20, 293-301.	2.7	74
161	The association of relationship quality and social networks with depression, anxiety, and suicidal ideation among older married adults: Findings from a cross-sectional analysis of the Irish Longitudinal Study on Ageing (TILDA). <i>Journal of Affective Disorders</i> , 2015, 179, 134-141.	2.0	74
162	Adjustment disorders in primary care: prevalence, recognition and use of services. <i>British Journal of Psychiatry</i> , 2012, 201, 137-142.	1.7	73

#	ARTICLE	IF	CITATIONS
163	Multimorbidity Patterns in a National Representative Sample of the Spanish Adult Population. PLoS ONE, 2014, 9, e84794.	1.1	73
164	The flavonoid quercetin induces acute vasodilator effects in healthy volunteers: Correlation with beta-glucuronidase activity. Pharmacological Research, 2014, 89, 11-18.	3.1	73
165	The association between socioeconomic status and depression among older adults in Finland, Poland and Spain: A comparative cross-sectional study of distinct measures and pathways. Journal of Affective Disorders, 2018, 241, 311-318.	2.0	73
166	Functional impairment in patients with major depressive disorder: the 2-year PERFORM study. Neuropsychiatric Disease and Treatment, 2018, Volume 14, 239-249.	1.0	72
167	Which somatic symptoms are associated with an unfavorable course in Asian patients with major depressive disorder?. Journal of Affective Disorders, 2013, 149, 182-188.	2.0	71
168	Determinants of Health and Disability in Ageing Population: The COURAGE in Europe Project (Collaborative Research on Ageing in Europe). Clinical Psychology and Psychotherapy, 2014, 21, 193-198.	1.4	70
169	Successful aging, dietary habits and health status of elderly individuals: A k-dimensional approach within the multi-national MEDIS study. Experimental Gerontology, 2014, 60, 57-63.	1.2	70
170	Thirty-day suicidal thoughts and behaviors among hospital workers during the first wave of the Spain COVID-19 outbreak. Depression and Anxiety, 2021, 38, 528-544.	2.0	70
171	Associations between DSM-IV mental disorders and diabetes mellitus: a role for impulse control disorders and depression. Diabetologia, 2014, 57, 699-709.	2.9	69
172	Post-traumatic stress disorder associated with sexual assault among women in the WHO World Mental Health Surveys. Psychological Medicine, 2018, 48, 155-167.	2.7	69
173	Perceived Need for Mental Health Care and Service Use Among Adults in Western Europe: Results of the ESEMeD Project. Psychiatric Services, 2009, 60, 1051-1058.	1.1	68
174	Incidence of Extrapyrarnidal Symptoms and Tardive Dyskinesia in Schizophrenia. Journal of Clinical Psychopharmacology, 2010, 30, 531-540.	0.7	68
175	Cross-national differences in the prevalence and correlates of burden among older family caregivers in the World Health Organization World Mental Health (WMH) Surveys. Psychological Medicine, 2013, 43, 865-879.	2.7	68
176	Diez años de investigación traslacional colaborativa en enfermedades mentales: el CIBERSAM. Revista De Psiquiatría Y Salud Mental, 2019, 12, 1-8.	1.0	68
177	Skeletal muscle mass in relation to 10 year cardiovascular disease incidence among middle aged and older adults: the ATTICA study. Journal of Epidemiology and Community Health, 2020, 74, 26-31.	2.0	68
178	The 12-item World Health Organization Disability Assessment Schedule II (WHO-DAS II): a nonparametric item response analysis. BMC Medical Research Methodology, 2010, 10, 45.	1.4	67
179	All-cause mortality and multimorbidity in older adults: The role of social support and loneliness. Experimental Gerontology, 2017, 99, 120-126.	1.2	67
180	Machine learning methodologies versus cardiovascular risk scores, in predicting disease risk. BMC Medical Research Methodology, 2018, 18, 179.	1.4	67

#	ARTICLE	IF	CITATIONS
181	Comorbidity within mental disorders: a comprehensive analysis based on 145 990 survey respondents from 27 countries. <i>Epidemiology and Psychiatric Sciences</i> , 2020, 29, e153.	1.8	67
182	Olanzapine vs. other antipsychotics in actual out-patient settings: six months tolerability results from the European Schizophrenia Out-patient Health Outcomes study. <i>Acta Psychiatrica Scandinavica</i> , 2005, 111, 232-243.	2.2	65
183	Falls in older aged adults in 22 European countries: incidence, mortality and burden of disease from 1990 to 2017. <i>Injury Prevention</i> , 2020, 26, i67-i74.	1.2	65
184	Early Cannabis Use and Estimated Risk of Later Onset of Depression Spells: Epidemiologic Evidence From the Population-based World Health Organization World Mental Health Survey Initiative. <i>American Journal of Epidemiology</i> , 2010, 172, 149-159.	1.6	64
185	Country-level and individual correlates of overweight and obesity among primary school children: a cross-sectional study in seven European countries. <i>BMC Public Health</i> , 2015, 15, 475.	1.2	64
186	Determinants of Quality of Life in Ageing Populations: Results from a Cross-Sectional Study in Finland, Poland and Spain. <i>PLoS ONE</i> , 2016, 11, e0159293.	1.1	64
187	Depression, sleep problems, and perceived stress among informal caregivers in 58 low-, middle-, and high-income countries: A cross-sectional analysis of community-based surveys. <i>Journal of Psychiatric Research</i> , 2018, 96, 115-123.	1.5	64
188	Influence of gender on social outcome in schizophrenia. <i>Acta Psychiatrica Scandinavica</i> , 2002, 106, 337-342.	2.2	63
189	Worldwide-Schizophrenia Outpatient Health Outcomes (W-SOHO): baseline characteristics of pan-regional observational data from more than 17,000 patients*. <i>International Journal of Clinical Practice</i> , 2009, 63, 1578-1588.	0.8	63
190	The cost of relapse in patients with schizophrenia in the European SOHO (Schizophrenia Outpatient) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 835-841.	2.5	63
191	Burden of chronic physical conditions and mental disorders in primary care. <i>British Journal of Psychiatry</i> , 2010, 196, 302-309.	1.7	63
192	Mild cognitive impairment in a Spanish representative sample: prevalence and associated factors. <i>International Journal of Geriatric Psychiatry</i> , 2016, 31, 858-867.	1.3	63
193	The role of muscle mass and body fat on disability among older adults: A cross-national analysis. <i>Experimental Gerontology</i> , 2015, 69, 27-35.	1.2	62
194	Determinants of societal costs in Alzheimer's disease: GERAS study baseline results. <i>Alzheimer's and Dementia</i> , 2015, 11, 933-945.	0.4	62
195	Long-term follow-up on health-related quality of life in major depressive disorder: a 2-year European cohort study. <i>Neuropsychiatric Disease and Treatment</i> , 2018, Volume 14, 1339-1350.	1.0	62
196	Treatment adequacy of anxiety and depressive disorders: Primary versus specialised care in Spain. <i>Journal of Affective Disorders</i> , 2006, 96, 9-20.	2.0	60
197	2-year course of bipolar disorder type I patients in outpatient care: Factors associated with remission and functional recovery. <i>European Neuropsychopharmacology</i> , 2011, 21, 287-293.	0.3	60
198	Relationship between living alone and common mental disorders in the 1993, 2000 and 2007 National Psychiatric Morbidity Surveys. <i>PLoS ONE</i> , 2019, 14, e0215182.	1.1	60

#	ARTICLE	IF	CITATIONS
199	Association between mental disorders and subsequent adult onset asthma. <i>Journal of Psychiatric Research</i> , 2014, 59, 179-188.	1.5	58
200	The association between psychotic experiences and disability: results from the <scp>WHO</scp> World Mental Health Surveys. <i>Acta Psychiatrica Scandinavica</i> , 2017, 136, 74-84.	2.2	58
201	Utilisation of mental health services and costs of patients with schizophrenia in three areas of Spain. <i>British Journal of Psychiatry</i> , 1998, 173, 334-340.	1.7	57
202	Do needs, symptoms or disability of outpatients with schizophrenia influence family burden?. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2008, 43, 612-618.	1.6	57
203	Are there gender differences in service use for mental disorders across countries in the European Union? Results from the EU-World Mental Health survey. <i>Journal of Epidemiology and Community Health</i> , 2014, 68, 649-656.	2.0	57
204	The association between obesity and back pain in nine countries: a cross-sectional study. <i>BMC Public Health</i> , 2015, 15, 123.	1.2	57
205	Decrease in olfactory and taste receptor expression in the dorsolateral prefrontal cortex in chronic schizophrenia. <i>Journal of Psychiatric Research</i> , 2015, 60, 109-116.	1.5	57
206	Rapid-cycling bipolar I disorder: Course and treatment outcome of a large sample across Europe. <i>Journal of Psychiatric Research</i> , 2008, 42, 1068-1075.	1.5	56
207	Diabetes mellitus and its association with central obesity and disability among older adults: A global perspective. <i>Experimental Gerontology</i> , 2015, 64, 70-77.	1.2	56
208	The associations between psychotic experiences and substance use and substance use disorders: findings from the World Health Organization World Mental Health surveys. <i>Addiction</i> , 2018, 113, 924-934.	1.7	56
209	Olanzapine versus haloperidol in the treatment of acute mania: clinical outcomes, health-related quality of life and work status. <i>International Clinical Psychopharmacology</i> , 2002, 17, 227-237.	0.9	55
210	Therapeutic Effectiveness and Patient Satisfaction after 6 Months of Treatment with Tadalafil, Sildenafil, and Vardenafil: Results from the Erectile Dysfunction Observational Study (EDOS). <i>European Urology</i> , 2007, 51, 541-550.	0.9	54
211	A descriptive case-register study of delusional disorder. <i>European Psychiatry</i> , 2008, 23, 125-133.	0.1	54
212	Factor structure, internal consistency and construct validity of the Sheehan Disability Scale in a Spanish primary care sample. <i>Journal of Evaluation in Clinical Practice</i> , 2010, 16, 895-901.	0.9	54
213	Assessment of Medication Adherence in a Cohort of Patients with Bipolar Disorder. <i>Pharmacopsychiatry</i> , 2010, 43, 263-270.	1.7	54
214	Health and happiness: cross-sectional household surveys in Finland, Poland and Spain. <i>Bulletin of the World Health Organization</i> , 2014, 92, 716-725.	1.5	54
215	Social network typologies and mortality risk among older people in China, India, and Latin America: A 10/66 Dementia Research Group population-based cohort study. <i>Social Science and Medicine</i> , 2015, 147, 134-143.	1.8	54
216	Social inequalities in mental health: results from the EU contribution to the World Mental Health Surveys Initiative. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2013, 48, 173-181.	1.6	53

#	ARTICLE	IF	CITATIONS
217	Health-related quality-of-life instruments for Alzheimer's disease and mixed dementia. <i>International Psychogeriatrics</i> , 2013, 25, 691-706.	0.6	53
218	Global mortality from dementia: Application of a new method and results from the Global Burden of Disease Study 2019. <i>Alzheimer's and Dementia: Translational Research and Clinical Interventions</i> , 2021, 7, e12200.	1.8	53
219	The associations of earlier trauma exposures and history of mental disorders with PTSD after subsequent traumas. <i>Molecular Psychiatry</i> , 2018, 23, 1892-1899.	4.1	52
220	Trajectories of Verbal Episodic Memory in Middle-aged and Older Adults: Evidence from the English Longitudinal Study of Ageing. <i>Journal of the American Geriatrics Society</i> , 2017, 65, 1274-1281.	1.3	51
221	Including information about co-morbidity in estimates of disease burden: results from the World Health Organization World Mental Health Surveys. <i>Psychological Medicine</i> , 2011, 41, 873-886.	2.7	49
222	n-3 Fatty acids, Mediterranean diet and cognitive function in normal aging: A systematic review. <i>Experimental Gerontology</i> , 2017, 91, 39-50.	1.2	49
223	Mild cognitive impairment diagnosed with the new DSM-5 criteria: prevalence and associations with non-cognitive psychopathology. <i>Acta Psychiatrica Scandinavica</i> , 2015, 131, 29-39.	2.2	48
224	Mediterranean diet and depression among older individuals: The multinational MEDIS study. <i>Experimental Gerontology</i> , 2018, 110, 67-72.	1.2	48
225	The Relation between Body Mass Index, Mental Health, and Functional Disability: A European Population Perspective. <i>Canadian Journal of Psychiatry</i> , 2008, 53, 679-688.	0.9	47
226	The transcription factor SP4 is reduced in postmortem cerebellum of bipolar disorder subjects: control by depolarization and lithium. <i>Bipolar Disorders</i> , 2011, 13, 474-485.	1.1	47
227	Gender Differences in Psychotropic Use Across Europe: Results From a Large Cross-Sectional, Population-Based Study. <i>European Psychiatry</i> , 2015, 30, 778-788.	0.1	47
228	Complementary and alternative medicine contacts by persons with mental disorders in 25 countries: results from the World Mental Health Surveys. <i>Epidemiology and Psychiatric Sciences</i> , 2018, 27, 552-567.	1.8	47
229	A comparative analysis of role attainment and impairment in binge-eating disorder and bulimia nervosa: results from the WHO World Mental Health Surveys. <i>Epidemiology and Psychiatric Sciences</i> , 2014, 23, 27-41.	1.8	46
230	Mild cognitive impairment is associated with falls among older adults: Findings from the Irish Longitudinal Study on Ageing (TILDA). <i>Experimental Gerontology</i> , 2016, 75, 42-47.	1.2	46
231	Frequent attendance in primary care: comparison and implications of different definitions. <i>British Journal of General Practice</i> , 2010, 60, e49-e55.	0.7	45
232	Risk and functional significance of psychotic experiences among individuals with depression in 44 low- and middle-income countries. <i>Psychological Medicine</i> , 2016, 46, 2655-2665.	2.7	45
233	The epidemiology of drug use disorders cross-nationally: Findings from the WHO's World Mental Health Surveys. <i>International Journal of Drug Policy</i> , 2019, 71, 103-112.	1.6	45
234	Differential impact of transient and chronic loneliness on health status. A longitudinal study. <i>Psychology and Health</i> , 2020, 35, 177-195.	1.2	45

#	ARTICLE	IF	CITATIONS
235	The European Schizophrenia Outpatient Health Outcomes Study: baseline findings across country and treatment. <i>Acta Psychiatrica Scandinavica</i> , 2003, 107, 7-15.	2.2	44
236	Pharmacological treatment and other predictors of treatment outcomes in previously untreated patients with schizophrenia: results from the European Schizophrenia Outpatient Health Outcomes (SOHO) study. <i>International Clinical Psychopharmacology</i> , 2005, 20, 199-205.	0.9	44
237	Dog-Assisted Therapy in the Treatment of Chronic Schizophrenia Inpatients. <i>Anthrozoos</i> , 2009, 22, 149-159.	0.7	44
238	Use of general practitioners versus mental health professionals in six European countries: the decisive role of the organization of mental health-care systems. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2013, 48, 137-149.	1.6	44
239	Factors associated with failure to achieve remission and with relapse after remission in patients with major depressive disorder in the PERFORM study. <i>Neuropsychiatric Disease and Treatment</i> , 2017, Volume 13, 2151-2165.	1.0	44
240	Burden of injury along the development spectrum: associations between the Socio-demographic Index and disability-adjusted life year estimates from the Global Burden of Disease Study 2017. <i>Injury Prevention</i> , 2020, 26, i12-i26.	1.2	44
241	Changes in depression and suicidal ideation under severe lockdown restrictions during the first wave of the COVID-19 pandemic in Spain: a longitudinal study in the general population. <i>Epidemiology and Psychiatric Sciences</i> , 2021, 30, e49.	1.8	44
242	Pain and depression in older people: Comorbidity and patterns of help seeking. <i>Journal of Affective Disorders</i> , 2009, 117, 193-196.	2.0	43
243	Why Do Clinicians Maintain Antidepressants in Some Patients With Acute Mania?. <i>Journal of Clinical Psychiatry</i> , 2010, 71, 1000-1006.	1.1	43
244	The association between obesity and severe disability among adults aged 50 or over in nine high-income, middle-income and low-income countries: a cross-sectional study. <i>BMJ Open</i> , 2015, 5, e007313-e007313.	0.8	42
245	Subclinical psychosis and suicidal behavior in England: Findings from the 2007 Adult Psychiatric Morbidity Survey. <i>Schizophrenia Research</i> , 2015, 168, 62-67.	1.1	42
246	Remote Assessment of Disease and Relapse in Major Depressive Disorder (RADAR-MDD): recruitment, retention, and data availability in a longitudinal remote measurement study. <i>BMC Psychiatry</i> , 2022, 22, 136.	1.1	42
247	Methodological Aspects in the Assessment of Treatment Effects in Observational Health Outcomes Studies. <i>Applied Health Economics and Health Policy</i> , 2006, 5, 11-25.	1.0	41
248	Antipsychotic type and correlates of antipsychotic treatment discontinuation in the outpatient treatment of schizophrenia. <i>European Psychiatry</i> , 2006, 21, 41-47.	0.1	41
249	Implications of modifying the duration requirement of generalized anxiety disorder in developed and developing countries. <i>Psychological Medicine</i> , 2009, 39, 1163.	2.7	41
250	IS ADHD IN CHILDHOOD ASSOCIATED WITH LIFETIME HOARDING SYMPTOMS? AN EPIDEMIOLOGICAL STUDY. <i>Depression and Anxiety</i> , 2013, 30, 741-748.	2.0	41
251	Depression and metabolic syndrome in the older population: A review of evidence. <i>Journal of Affective Disorders</i> , 2018, 237, 56-64.	2.0	41
252	Use of Primary and Specialized Mental Health Care for a Major Depressive Episode in Spain by ESEMeD Respondents. <i>Psychiatric Services</i> , 2011, 62, 152-161.	1.1	40

#	ARTICLE	IF	CITATIONS
253	Cannabis and bipolar disorder: does quitting cannabis use during manic/mixed episode improve clinical/functional outcomes?. <i>Acta Psychiatrica Scandinavica</i> , 2015, 131, 100-110.	2.2	40
254	Factors associated with long-term impact on informal caregivers during Alzheimer's disease dementia progression: 36-month results from GERAS. <i>International Psychogeriatrics</i> , 2020, 32, 267-277.	0.6	40
255	Predictors of the course of illness in outpatients with schizophrenia: A prospective three year study. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2008, 32, 1287-1292.	2.5	39
256	What is the relative importance of self reported psychotic symptoms in epidemiological studies? Results from the ESEMeD-Catalonia Study. <i>Schizophrenia Research</i> , 2008, 102, 261-269.	1.1	39
257	Analysis of Sp transcription factors in the postmortem brain of chronic schizophrenia: A pilot study of relationship to negative symptoms. <i>Journal of Psychiatric Research</i> , 2013, 47, 926-934.	1.5	39
258	Cohort Profile: The Ageing Trajectories of Health - Longitudinal Opportunities and Synergies (ATHLOS) project. <i>International Journal of Epidemiology</i> , 2019, 48, 1052-1053i.	0.9	39
259	The need for a behavioural science focus in research on mental health and mental disorders. <i>International Journal of Methods in Psychiatric Research</i> , 2014, 23, 28-40.	1.1	38
260	Visual impairment and multimorbidity in a representative sample of the Spanish population. <i>BMC Public Health</i> , 2014, 14, 815.	1.2	38
261	Complex Interaction Between Symptoms, Social Factors, and Gender in Social Functioning in a Community-Dwelling Sample of Schizophrenia. <i>Psychiatric Quarterly</i> , 2011, 82, 261-274.	1.1	37
262	The burden of disease in Spain: Results from the Global Burden of Disease 2016. <i>Medicina Clínica (English Edition)</i> , 2018, 151, 171-190.	0.1	37
263	The burden of disease in Greece, health loss, risk factors, and health financing, 2000-2016: an analysis of the Global Burden of Disease Study 2016. <i>Lancet Public Health</i> , The, 2018, 3, e395-e406.	4.7	37
264	Effects of transient and chronic loneliness on major depression in older adults: A longitudinal study. <i>International Journal of Geriatric Psychiatry</i> , 2021, 36, 76-85.	1.3	37
265	Parental Bonding and Suicidality in Adulthood. <i>Australian and New Zealand Journal of Psychiatry</i> , 2007, 41, 66-73.	1.3	36
266	Adverse parenting as a risk factor in the occurrence of anxiety disorders. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2008, 43, 266-272.	1.6	36
267	Antipsychotic treatment discontinuation in previously untreated patients with schizophrenia: 36-month results from the SOHO study. <i>Journal of Psychiatric Research</i> , 2009, 43, 265-273.	1.5	36
268	High and low suicidality in Europe: A fine-grained comparison of France and Spain within the ESEMeD surveys. <i>Journal of Affective Disorders</i> , 2011, 133, 247-256.	2.0	36
269	Replication of previous genome-wide association studies of psychiatric diseases in a large schizophrenia case-control sample from Spain. <i>Schizophrenia Research</i> , 2014, 159, 107-113.	1.1	36
270	Disease Progression in Mild Dementia due to Alzheimer Disease in an 18-Month Observational Study (GERAS): The Impact on Costs and Caregiver Outcomes. <i>Dementia and Geriatric Cognitive Disorders Extra</i> , 2017, 7, 87-100.	0.6	36

#	ARTICLE	IF	CITATIONS
271	Determinants of Health Trajectories in England and the United States: An Approach to Identify Different Patterns of Healthy Aging. <i>Journals of Gerontology - Series A Biological Sciences and Medical Sciences</i> , 2018, 73, 1512-1518.	1.7	36
272	Relationship between subjective well-being and healthy lifestyle behaviours in older adults: a longitudinal study. <i>Aging and Mental Health</i> , 2020, 24, 611-619.	1.5	36
273	Treatment-seeking Behavior of Erectile Dysfunction Patients in Europe: Results of the Erectile Dysfunction Observational Study. <i>Journal of Sexual Medicine</i> , 2006, 3, 530-540.	0.3	35
274	Association between traumatic events and post-traumatic stress disorder: results from the ESEMeD-Spain study. <i>Epidemiology and Psychiatric Sciences</i> , 2015, 24, 172-183.	1.8	35
275	Identifying factors of activities of daily living important for cost and caregiver outcomes in Alzheimer's disease. <i>International Psychogeriatrics</i> , 2016, 28, 247-259.	0.6	35
276	The protective properties of Act-Belong-Commit indicators against incident depression, anxiety, and cognitive impairment among older Irish adults: Findings from a prospective community-based study. <i>Experimental Gerontology</i> , 2017, 91, 79-87.	1.2	35
277	Relationship Between Major Depression Symptom Severity and Sleep Collected Using a Wristband Wearable Device: Multicenter Longitudinal Observational Study. <i>JMIR MHealth and UHealth</i> , 2021, 9, e24604.	1.8	35
278	Empirical redefinition of delusional disorder and its phenomenology: the DELIREMP study. <i>Comprehensive Psychiatry</i> , 2013, 54, 243-255.	1.5	34
279	Relation between jumping to conclusions and cognitive functioning in people with schizophrenia in contrast with healthy participants. <i>Schizophrenia Research</i> , 2014, 159, 211-217.	1.1	34
280	Post-traumatic stress disorder associated with life-threatening motor vehicle collisions in the WHO World Mental Health Surveys. <i>BMC Psychiatry</i> , 2016, 16, 257.	1.1	34
281	Gender-related differences in the multi-pathway effect of social determinants on quality of life in older age—the COURAGE in Europe project. <i>Quality of Life Research</i> , 2017, 26, 1865-1878.	1.5	34
282	Dietary inflammatory potential is linked to cardiovascular disease risk burden in the US adult population. <i>International Journal of Cardiology</i> , 2017, 240, 409-413.	0.8	34
283	Estimating the prevalence of schizophrenia in Spain using a disease model. <i>Schizophrenia Research</i> , 2006, 86, 194-201.	1.1	33
284	Symptomatic remission in previously untreated patients with schizophrenia: 2-year results from the SOHO study. <i>Psychopharmacology</i> , 2007, 191, 1015-1022.	1.5	33
285	Marginal structural models might overcome confounding when analyzing multiple treatment effects in observational studies. <i>Journal of Clinical Epidemiology</i> , 2008, 61, 525-530.	2.4	33
286	Predictors of switching from mania to depression in a large observational study across Europe (EMBLEM). <i>Journal of Affective Disorders</i> , 2009, 118, 118-123.	2.0	33
287	Tolerability of outpatient antipsychotic treatment: 36-month results from the European Schizophrenia Outpatient Health Outcomes (SOHO) study. <i>European Neuropsychopharmacology</i> , 2009, 19, 542-550.	0.3	33
288	Association of headache with childhood adversity and mental disorder: cross-national study. <i>British Journal of Psychiatry</i> , 2009, 194, 111-116.	1.7	33

#	ARTICLE	IF	CITATIONS
289	Clinical features, comorbidity, and cognitive impairment in elderly bipolar patients. <i>Neuropsychiatric Disease and Treatment</i> , 2016, 12, 1203.	1.0	33
290	Posttraumatic stress disorder associated with unexpected death of a loved one: Cross-national findings from the world mental health surveys. <i>Depression and Anxiety</i> , 2017, 34, 315-326.	2.0	33
291	Post-traumatic stress symptoms are associated with physical multimorbidity: Findings from the Adult Psychiatric Morbidity Survey 2007. <i>Journal of Affective Disorders</i> , 2018, 232, 385-392.	2.0	33
292	Long-term trajectories of depressive symptoms in old age: Relationships with sociodemographic and health-related factors. <i>Journal of Affective Disorders</i> , 2019, 246, 329-337.	2.0	33
293	Are environmental pollution and biodiversity levels associated to the spread and mortality of COVID-19? A four-month global analysis. <i>Environmental Pollution</i> , 2021, 271, 116326.	3.7	33
294	Longitudinal Relationships Between Positive Affect, Loneliness, and Suicide Ideation: Age-specific Factors in a General Population. <i>Suicide and Life-Threatening Behavior</i> , 2019, 49, 90-103.	0.9	33
295	Emotional Distress in Disaster Victims. <i>Journal of Nervous and Mental Disease</i> , 1993, 181, 388-393.	0.5	32
296	Gender differences in delusional disorder: Evidence from an outpatient sample. <i>Psychiatry Research</i> , 2010, 177, 235-239.	1.7	32
297	Premarital mental disorders and physical violence in marriage: cross-national study of married couples. <i>British Journal of Psychiatry</i> , 2011, 199, 330-337.	1.7	32
298	Is cognitive impairment associated with suicidality? A population-based study. <i>European Neuropsychopharmacology</i> , 2015, 25, 203-213.	0.3	32
299	Psychotic symptoms and smoking in 44 countries. <i>Acta Psychiatrica Scandinavica</i> , 2016, 133, 497-505.	2.2	32
300	Epidemiology of depression with psychotic experiences and its association with chronic physical conditions in 47 low- and middle-income countries. <i>Psychological Medicine</i> , 2017, 47, 531-542.	2.7	32
301	Previous Mental Disorders and Subsequent Onset of Chronic Back or Neck Pain: Findings From 19 Countries. <i>Journal of Pain</i> , 2018, 19, 99-110.	0.7	32
302	Loneliness and depression among older European adults: The role of perceived neighborhood built environment. <i>Health and Place</i> , 2020, 62, 102280.	1.5	32
303	Development of a common scale for measuring healthy ageing across the world: results from the ATHLOS consortium. <i>International Journal of Epidemiology</i> , 2021, 50, 880-892.	0.9	32
304	Thirty-day suicidal thoughts and behaviours in the Spanish adult general population during the first wave of the Spain COVID-19 pandemic. <i>Epidemiology and Psychiatric Sciences</i> , 2021, 30, e19.	1.8	32
305	Suicide ideation across reproductive life cycle of women Results from a European epidemiological study. <i>Journal of Affective Disorders</i> , 2009, 116, 144-147.	2.0	31
306	Semaphorin and plexin gene expression is altered in the prefrontal cortex of schizophrenia patients with and without auditory hallucinations. <i>Psychiatry Research</i> , 2015, 229, 850-857.	1.7	31

#	ARTICLE	IF	CITATIONS
307	Cognitive reserve is associated with quality of life: A population-based study. <i>Experimental Gerontology</i> , 2017, 87, 67-73.	1.2	31
308	Cross-national prevalence and factors associated with suicide ideation and attempts in older and young-and-middle age people. <i>Aging and Mental Health</i> , 2020, 24, 1533-1542.	1.5	31
309	Multi-Country Evaluation of Affective Experience: Validation of an Abbreviated Version of the Day Reconstruction Method in Seven Countries. <i>PLoS ONE</i> , 2013, 8, e61534.	1.1	31
310	Utility of the twelve-item World Health Organization Disability Assessment Schedule II (WHO-DAS II) for discriminating depression and severity in Spanish primary care patients. <i>Quality of Life Research</i> , 2010, 19, 97-101.	1.5	30
311	Clinical and Cognitive Correlates of Psychiatric Comorbidity in Delusional Disorder Outpatients. <i>Australian and New Zealand Journal of Psychiatry</i> , 2011, 45, 416-425.	1.3	30
312	Development of the insight scale for affective disorders (ISAD): Modification from the scale to assess unawareness of mental disorder. <i>Journal of Affective Disorders</i> , 2012, 142, 65-71.	2.0	30
313	Recognition of anxiety disorders by the general practitioner: results from the DASMAP Study. <i>General Hospital Psychiatry</i> , 2012, 34, 227-233.	1.2	30
314	Factors associated with active aging in Finland, Poland, and Spain. <i>International Psychogeriatrics</i> , 2014, 26, 1363-1375.	0.6	30
315	Suicidality and its relationship with depression, alcohol disorders and childhood experiences of violence: Results from the ESEMeD study. <i>Journal of Affective Disorders</i> , 2015, 175, 168-174.	2.0	30
316	Sex differences in the course of schizophrenia across diverse regions of the world. <i>Neuropsychiatric Disease and Treatment</i> , 2016, Volume 12, 2927-2939.	1.0	30
317	Mobility Difficulties, Physical Activity, and All-cause Mortality Risk in a Nationally representative Sample of Older Adults. <i>Journals of Gerontology - Series A Biological Sciences and Medical Sciences</i> , 2018, 73, 1272-1279.	1.7	30
318	Cost-effectiveness of different clinical interventions for reducing the burden of schizophrenia in Spain. <i>Acta Psychiatrica Scandinavica</i> , 2006, 114, 29-38.	2.2	29
319	Work impairment in bipolar disorder patients results from a two-year observational study (EMBLEM). <i>European Psychiatry</i> , 2010, 25, 338-344.	0.1	29
320	Recreational drug use among individuals living with HIV in Europe: review of the prevalence, comparison with the general population and HIV guidelines recommendations. <i>Frontiers in Microbiology</i> , 2015, 6, 690.	1.5	29
321	Understanding the impact of persistent symptoms in schizophrenia: Cross-sectional findings from the Pattern study. <i>Schizophrenia Research</i> , 2015, 169, 234-240.	1.1	29
322	The Associations Between Preexisting Mental Disorders and Subsequent Onset of Chronic Headaches: A Worldwide Epidemiologic Perspective. <i>Journal of Pain</i> , 2015, 16, 42-52.	0.7	29
323	The Impact of Depression on the Development of Mild Cognitive Impairment over 3 Years of Follow-Up: A Population-Based Study. <i>Dementia and Geriatric Cognitive Disorders</i> , 2017, 43, 155-169.	0.7	29
324	The role of socio-economic status and neighborhood social capital on loneliness among older adults: evidence from the Sant Boi Aging Study. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2017, 52, 1237-1246.	1.6	29

#	ARTICLE	IF	CITATIONS
325	Association of bullying victimization with overweight and obesity among adolescents from 41 low- and middle-income countries. <i>Pediatric Obesity</i> , 2020, 15, e12571.	1.4	29
326	Use of concomitant medication with antipsychotic treatment in outpatients with schizophrenia: Results from the European Schizophrenia Outpatients Health Outcomes (SOHO) study. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2005, 29, 972-982.	2.5	28
327	Needs and its relation to symptom dimensions in a sample of outpatients with schizophrenia. <i>Schizophrenia Research</i> , 2005, 75, 129-134.	1.1	28
328	Patterns of switching phosphodiesterase type 5 inhibitors in the treatment of erectile dysfunction: results from the Erectile Dysfunction Observational Study. <i>International Journal of Clinical Practice</i> , 2007, 61, 1850-1862.	0.8	28
329	The Schizophrenia Outpatient Health Outcomes (SOHO) study: 3-year results of antipsychotic treatment discontinuation and related clinical factors in Spain. <i>European Psychiatry</i> , 2008, 23, 1-7.	0.1	28
330	Cross-National Comparisons of Sex Differences in Opportunities to Use Alcohol or Drugs, and the Transitions to Use. <i>Substance Use and Misuse</i> , 2011, 46, 1169-1178.	0.7	28
331	The role of negative symptoms in the context of cognitive remediation for schizophrenia. <i>Schizophrenia Research</i> , 2013, 150, 58-63.	1.1	28
332	Cognitive correlates of verbal memory and verbal fluency in schizophrenia, and differential effects of various clinical symptoms between male and female patients. <i>Schizophrenia Research</i> , 2013, 147, 81-85.	1.1	28
333	Recovery from DSM-IV post-traumatic stress disorder in the WHO World Mental Health surveys. <i>Psychological Medicine</i> , 2018, 48, 437-450.	2.7	28
334	Dietary Patterns and Their Association with Anxiety Symptoms among Older Adults: The ATTICA Study. <i>Nutrients</i> , 2019, 11, 1250.	1.7	28
335	Mediterranean diet and wellbeing: evidence from a nationwide survey. <i>Psychology and Health</i> , 2019, 34, 321-335.	1.2	28
336	Adequacy of Antidepressant Treatment in Spanish Primary Care: A Naturalistic Six-Month Follow-Up Study. <i>Psychiatric Services</i> , 2008, 59, 78-83.	1.1	27
337	Reduced expression of SP1 and SP4 transcription factors in peripheral blood mononuclear cells in first-episode psychosis. <i>Journal of Psychiatric Research</i> , 2013, 47, 1608-1614.	1.5	27
338	Predictors and consequences of adherence to the treatment of pediatric patients with attention-deficit/hyperactivity disorder in Central Europe and East Asia. <i>Patient Preference and Adherence</i> , 2013, 7, 987.	0.8	27
339	The burden of disease in Spain: results from the global burden of disease study 2010. <i>BMC Medicine</i> , 2014, 12, 236.	2.3	27
340	A dimensional comparison between delusional disorder, schizophrenia and schizoaffective disorder. <i>Schizophrenia Research</i> , 2015, 169, 248-254.	1.1	27
341	Proportion of patients without mental disorders being treated in mental health services worldwide. <i>British Journal of Psychiatry</i> , 2015, 206, 101-109.	1.7	27
342	How useful is the EQ-5D in assessing the impact of caring for people with Alzheimer's disease?. <i>Health and Quality of Life Outcomes</i> , 2017, 15, 16.	1.0	27

#	ARTICLE	IF	CITATIONS
343	Psychotic experiences and general medical conditions: a cross-national analysis based on 28 002 respondents from 16 countries in the WHO World Mental Health Surveys. <i>Psychological Medicine</i> , 2018, 48, 2730-2739.	2.7	27
344	Psychotic experiences and religiosity: data from the WHO World Mental Health Surveys. <i>Acta Psychiatrica Scandinavica</i> , 2018, 137, 306-315.	2.2	27
345	Cross-national patterns of substance use disorder treatment and associations with mental disorder comorbidity in the WHO World Mental Health Surveys. <i>Addiction</i> , 2019, 114, 1446-1459.	1.7	27
346	Association of Cohort and Individual Substance Use With Risk of Transitioning to Drug Use, Drug Use Disorder, and Remission From Disorder. <i>JAMA Psychiatry</i> , 2019, 76, 708.	6.0	27
347	Disability Mediates the Impact of Common Conditions on Perceived Health. <i>PLoS ONE</i> , 2013, 8, e65858.	1.1	27
348	Predictability of rehospitalization for schizophrenia. <i>European Archives of Psychiatry and Clinical Neuroscience</i> , 1994, 244, 241-246.	1.8	26
349	REPYFLEC cognitive remediation group training in schizophrenia. <i>Schizophrenia Research</i> , 2012, 142, 137-144.	1.1	26
350	Symptomatic remission and patient quality of life in an observational study of schizophrenia: Is there a relationship?. <i>Psychiatry Research</i> , 2014, 220, 163-169.	1.7	26
351	Altered CSNK1E, FABP4 and NEFH protein levels in the dorsolateral prefrontal cortex in schizophrenia. <i>Schizophrenia Research</i> , 2016, 177, 88-97.	1.1	26
352	Direct and indirect costs and resource use in dementia care: A cross-sectional study in patients living at home. <i>International Journal of Nursing Studies</i> , 2016, 55, 39-49.	2.5	26
353	Brain structure in women at risk of postpartum psychosis: an MRI study. <i>Translational Psychiatry</i> , 2017, 7, 1286.	2.4	26
354	Understanding the course of persistent symptoms in schizophrenia: Longitudinal findings from the pattern study. <i>Psychiatry Research</i> , 2018, 267, 56-62.	1.7	26
355	Determinants of time to institutionalisation and related healthcare and societal costs in a community-based cohort of patients with Alzheimer's disease dementia. <i>European Journal of Health Economics</i> , 2019, 20, 343-355.	1.4	26
356	Predicting Depressive Symptom Severity Through Individuals' Nearby Bluetooth Device Count Data Collected by Mobile Phones: Preliminary Longitudinal Study. <i>JMIR MHealth and UHealth</i> , 2021, 9, e29840.	1.8	26
357	Deficits in Executive and Memory Processes in Delusional Disorder: A Case-Control Study. <i>PLoS ONE</i> , 2013, 8, e67341.	1.1	26
358	Longitudinal Relationships Between Depressive Symptom Severity and Phone-Measured Mobility: Dynamic Structural Equation Modeling Study. <i>JMIR Mental Health</i> , 2022, 9, e34898.	1.7	26
359	Cost-Utility Analysis of Treatment with Olanzapine Compared with Other Antipsychotic Treatments in Patients with Schizophrenia in the Pan-European SOHO Study. <i>Pharmacoeconomics</i> , 2008, 26, 341-358.	1.7	25
360	Brain alterations in low-frequency fluctuations across multiple bands in obsessive compulsive disorder. <i>Brain Imaging and Behavior</i> , 2017, 11, 1690-1706.	1.1	25

#	ARTICLE	IF	CITATIONS
361	Physical multimorbidity and subjective cognitive complaints among adults in the United Kingdom: a cross-sectional community-based study. <i>Scientific Reports</i> , 2019, 9, 12417.	1.6	25
362	Multimorbidity patterns in low-middle and high income regions: a multiregion latent class analysis using ATHLOS harmonised cohorts. <i>BMJ Open</i> , 2020, 10, e034441.	0.8	25
363	A framework for evidence-based mental health care and policy. <i>Acta Psychiatrica Scandinavica</i> , 2006, 114, 5-11.	2.2	24
364	Determinants of Voluntary vs. Involuntary Admission in Bipolar Disorder and the Impact of Adherence. <i>Pharmacopsychiatry</i> , 2008, 41, 29-36.	1.7	24
365	Validaci3n al castellano de la escala de estigma social: Community Attitudes towards Mental Illness en poblaci3n adolescente. <i>Revista De PsiquiatrAa Y Salud Mental</i> , 2016, 9, 150-157.	1.0	24
366	Social determinants, health status and 10-year mortality among 10,906 older adults from the English longitudinal study of aging: the ATHLOS project. <i>BMC Public Health</i> , 2018, 18, 1357.	1.2	24
367	Stability of clinically relevant depression symptoms in old4age across 11 cohorts: a multi4state study. <i>Acta Psychiatrica Scandinavica</i> , 2019, 140, 541-551.	2.2	24
368	Childhood generalized specific phobia as an early marker of internalizing psychopathology across the lifespan: results from the World Mental Health Surveys. <i>BMC Medicine</i> , 2019, 17, 101.	2.3	24
369	The association of religiosity with suicidal ideation and suicide attempts in the United Kingdom. <i>Acta Psychiatrica Scandinavica</i> , 2019, 139, 164-173.	2.2	24
370	Assessment of remission in schizophrenia with the CGI and CGI-SCH scales. <i>Acta Psychiatrica Scandinavica</i> , 2007, 115, 163-164.	2.2	23
371	Comorbid painful physical symptoms and anxiety: Prevalence, work loss and help-seeking. <i>Journal of Affective Disorders</i> , 2008, 109, 264-272.	2.0	23
372	Increased SP4 and SP1 transcription factor expression in the postmortem hippocampus of chronic schizophrenia. <i>Journal of Psychiatric Research</i> , 2014, 58, 189-196.	1.5	23
373	What Drives Country Differences in Cost of Alzheimer4s Disease? An Explanation from Resource Use in the GERAS Study. <i>Journal of Alzheimer's Disease</i> , 2017, 57, 797-812.	1.2	23
374	The Effect of Exclusive Olive Oil Consumption on Successful Aging: A Combined Analysis of the ATTICA and MEDIS Epidemiological Studies. <i>Foods</i> , 2019, 8, 25.	1.9	23
375	Toward measuring effective treatment coverage: critical bottlenecks in quality- and user-adjusted coverage for major depressive disorder. <i>Psychological Medicine</i> , 2022, 52, 1948-1958.	2.7	23
376	Chronic mania revisited: Factors associated with treatment non-response during prospective follow-up of a large European cohort (EMBLEM). <i>World Journal of Biological Psychiatry</i> , 2008, 9, 313-320.	1.3	22
377	Validity and Reliability of the Hamilton Depression Rating Scale (5 Items) for Manic and Mixed Bipolar Disorders. <i>Journal of Nervous and Mental Disease</i> , 2009, 197, 682-686.	0.5	22
378	The Cost of Relapse for Patients with a Manic/Mixed Episode of Bipolar Disorder in the EMBLEM Study. <i>Pharmacoeconomics</i> , 2010, 28, 555-566.	1.7	22

#	ARTICLE	IF	CITATIONS
379	Exploring the excess mortality due to depressive symptoms in a community-based sample: The role of Alzheimer's Disease. <i>Journal of Affective Disorders</i> , 2016, 202, 163-170.	2.0	22
380	Anti-NMDAR antibodies in new-onset psychosis. Positive results in an HIV-infected patient. <i>Brain, Behavior, and Immunity</i> , 2016, 56, 56-60.	2.0	22
381	Act-Belong-Commit Indicators Promote Mental Health and Wellbeing among Irish Older Adults. <i>American Journal of Health Behavior</i> , 2018, 42, 31-45.	0.6	22
382	Relationship between attention-deficit hyperactivity disorder symptoms and problem gambling: A mediation analysis of influential factors among 7,403 individuals from the UK. <i>Journal of Behavioral Addictions</i> , 2018, 7, 781-791.	1.9	22
383	Mortality in unipolar depression preceding and following chronic somatic diseases. <i>Acta Psychiatrica Scandinavica</i> , 2018, 138, 500-508.	2.2	22
384	A comparison of <i>DSM</i> 5 and <i>DSM</i> IV agoraphobia in the World Mental Health Surveys. <i>Depression and Anxiety</i> , 2019, 36, 499-510.	2.0	22
385	Psychotic experiences and subjective cognitive complaints among 224 842 people in 48 low- and middle-income countries. <i>Epidemiology and Psychiatric Sciences</i> , 2020, 29, e11.	1.8	22
386	Sociodemographic Indicators of Health Status Using a Machine Learning Approach and Data from the English Longitudinal Study of Aging (ELSA). <i>Medical Science Monitor</i> , 2019, 25, 1994-2001.	0.5	22
387	Psychotic-Like Experiences and Nonsuicidal Self-Injury in England: Results from a National Survey. <i>PLoS ONE</i> , 2015, 10, e0145533.	1.1	22
388	Financing Mental Health Care in Spain: Context and critical issues. <i>European Journal of Psychiatry</i> , 2006, 20, .	0.7	22
389	Cost?utility of selective serotonin reuptake inhibitors for depression in primary care in Catalonia. <i>Acta Psychiatrica Scandinavica</i> , 2006, 114, 39-47.	2.2	21
390	Service utilization and costs of first-onset schizophrenia in two widely differing health service areas in North-East Spain. <i>Acta Psychiatrica Scandinavica</i> , 2007, 100, 335-343.	2.2	21
391	Remission in the outpatient care of schizophrenia: 3-Year results from the Schizophrenia Outpatients Health Outcomes (SOHO) Study in France. <i>European Psychiatry</i> , 2008, 23, 491-496.	0.1	21
392	Impact of mental disorders and chronic physical conditions in health-related quality of life among primary care patients: results from an epidemiological study. <i>Quality of Life Research</i> , 2009, 18, 1011-1018.	1.5	21
393	Psychotropic prescribing in Catalonia: results from an epidemiological study. <i>Family Practice</i> , 2012, 29, 154-162.	0.8	21
394	Patterns of care and dropout rates from outpatient mental healthcare in low-, middle- and high-income countries from the World Health Organization's World Mental Health Survey Initiative. <i>Psychological Medicine</i> , 2021, 51, 2104-2116.	2.7	21
395	Mental health impact of the first wave of COVID-19 pandemic on Spanish healthcare workers: A large cross-sectional survey. <i>Revista De PsiquiatrÃa Y Salud Mental (English Edition)</i> , 2021, 14, 90-105.	0.2	21
396	Factors Associated With Use of Psychiatrists and Nonpsychiatrist Providers by ESEMeD Respondents in Six European Countries. <i>Psychiatric Services</i> , 2011, 62, 143-151.	1.1	20

#	ARTICLE	IF	CITATIONS
397	Comorbidity of major depressive episode and chronic physical conditions in Spain, a country with low prevalence of depression. <i>General Hospital Psychiatry</i> , 2012, 34, 510-517.	1.2	20
398	Diagnosis delay in first episodes of major depression: A study of primary care patients in Spain. <i>Journal of Affective Disorders</i> , 2013, 150, 1247-1250.	2.0	20
399	Horizon 2020 Priorities in Clinical Mental Health Research: Results of a Consensus-Based ROAMER Expert Survey. <i>International Journal of Environmental Research and Public Health</i> , 2014, 11, 10915-10939.	1.2	20
400	Towards Horizon 2020: challenges and advances for clinical mental health research – outcome of an expert survey. <i>Neuropsychiatric Disease and Treatment</i> , 2014, 10, 1057.	1.0	20
401	Association of experienced and evaluative well-being with health in nine countries with different income levels: a cross-sectional study. <i>Globalization and Health</i> , 2017, 13, 65.	2.4	20
402	Recovery in patients with major depressive disorder (MDD): results of a 6-month, multinational, observational study. <i>Patient Preference and Adherence</i> , 2017, Volume 11, 1859-1868.	0.8	20
403	The association between psychotic experiences and health-related quality of life: a cross-national analysis based on World Mental Health Surveys. <i>Schizophrenia Research</i> , 2018, 201, 46-53.	1.1	20
404	The association of social support networks and loneliness with negative perceptions of ageing: evidence from the Irish Longitudinal Study on Ageing (TILDA). <i>Ageing and Society</i> , 2019, 39, 1070-1090.	1.2	20
405	Informal Caregiving, Chronic Physical Conditions, and Physical Multimorbidity in 48 Low- and Middle-Income Countries. <i>Journals of Gerontology - Series A Biological Sciences and Medical Sciences</i> , 2020, 75, 1572-1578.	1.7	20
406	Sarcopenia and fall-related injury among older adults in five low- and middle-income countries. <i>Experimental Gerontology</i> , 2021, 147, 111262.	1.2	20
407	Is it always the same? Variability of depressive symptoms across six European countries. <i>Psychiatry Research</i> , 2009, 168, 137-144.	1.7	19
408	Dropping out of mental health treatment among patients with depression and anxiety by type of provider: results of the European Study of the Epidemiology of Mental Disorders. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2011, 46, 273-280.	1.6	19
409	Risk factors for suicide behaviors in the observational schizophrenia outpatient health outcomes (SOHO) study. <i>BMC Psychiatry</i> , 2012, 12, 83.	1.1	19
410	The Validity of the Instrument to Evaluate Social Network in the Ageing Population: The Collaborative Research on Ageing in Europe Social Network Index. <i>Clinical Psychology and Psychotherapy</i> , 2014, 21, 227-241.	1.4	19
411	The state of the art on European well-being research within the area of mental health. <i>International Journal of Clinical and Health Psychology</i> , 2015, 15, 171-179.	2.7	19
412	Exploring the path between depression, anxiety and 10-year cardiovascular disease incidence, among apparently healthy Greek middle-aged adults: The ATTICA study. <i>Maturitas</i> , 2017, 106, 73-79.	1.0	19
413	<p>Patient-reported depression severity and cognitive symptoms as determinants of functioning in patients with major depressive disorder: a secondary analysis of the 2-year prospective PERFORM study</p>. <i>Neuropsychiatric Disease and Treatment</i> , 2019, Volume 15, 2313-2323.	1.0	19
414	Fruit and Vegetable Consumption and Sarcopenia among Older Adults in Low- and Middle-Income Countries. <i>Nutrients</i> , 2020, 12, 706.	1.7	19

#	ARTICLE	IF	CITATIONS
415	Exploring the effect of loneliness on all-cause mortality: Are there differences between older adults and younger and middle-aged adults?. <i>Social Science and Medicine</i> , 2020, 258, 113087.	1.8	19
416	Association between Food Insecurity and Sarcopenia among Adults Aged ≥65 Years in Low- and Middle-Income Countries. <i>Nutrients</i> , 2021, 13, 1879.	1.7	19
417	Multimorbidity and obesity in older adults from six low- and middle-income countries. <i>Preventive Medicine</i> , 2021, 153, 106816.	1.6	19
418	Influence of age at onset on social functioning in outpatients with schizophrenia. <i>European Journal of Psychiatry</i> , 2006, 20, .	0.7	19
419	The Association Between Home Stay and Symptom Severity in Major Depressive Disorder: Preliminary Findings From a Multicenter Observational Study Using Geolocation Data From Smartphones. <i>JMIR MHealth and UHealth</i> , 2022, 10, e28095.	1.8	19
420	Effectiveness and cost-effectiveness of antidepressant treatment in primary health care: A six-month randomised study comparing fluoxetine to imipramine. <i>Journal of Affective Disorders</i> , 2006, 91, 153-163.	2.0	18
421	Stability and treatment outcome of distinct classes of mania. <i>European Psychiatry</i> , 2008, 23, 360-367.	0.1	18
422	Overview of the findings from the European SOHO study. <i>Expert Review of Neurotherapeutics</i> , 2008, 8, 873-880.	1.4	18
423	Incidence of stressful life events and influence of sociodemographic and clinical variables on the onset of first-episode psychosis. <i>Psychiatry Research</i> , 2016, 245, 108-115.	1.7	18
424	Eficacia, eficiencia y efectividad en el tratamiento multidimensional de la esquizofrenia: proyecto Rethinking. <i>Revista De Psiquiatría Y Salud Mental</i> , 2017, 10, 4-20.	1.0	18
425	The association between pain and suicidal behavior in an English national sample: The role of psychopathology. <i>Journal of Psychiatric Research</i> , 2018, 98, 39-46.	1.5	18
426	Anti-Inflammatory Nutrition and Successful Ageing in Elderly Individuals: The Multinational MEDIS Study. <i>Gerontology</i> , 2018, 64, 3-10.	1.4	18
427	Adaptation and validation of the quality-of-life scale: Satisfaction with Life Domains Scale by Baker and Intagliata. <i>Comprehensive Psychiatry</i> , 2009, 50, 76-80.	1.5	17
428	Empirically driven definitions of "good," "moderate," and "poor" levels of functioning in the treatment of schizophrenia. <i>Quality of Life Research</i> , 2013, 22, 2085-2094.	1.5	17
429	Three-factor model of premorbid adjustment in a sample with chronic schizophrenia and first-episode psychosis. <i>Schizophrenia Research</i> , 2013, 151, 252-258.	1.1	17
430	Which somatic symptoms are associated with an unfavorable course in Chinese patients with major depressive disorder?. <i>Asia-Pacific Psychiatry</i> , 2015, 7, 427-435.	1.2	17
431	Predictors of remission in the treatment of major depressive disorder: real-world evidence from a 6-month prospective observational study. <i>Neuropsychiatric Disease and Treatment</i> , 2015, 11, 197.	1.0	17
432	Associations between DSM-IV mental disorders and subsequent non-fatal, self-reported stroke. <i>Journal of Psychosomatic Research</i> , 2015, 79, 130-136.	1.2	17

#	ARTICLE	IF	CITATIONS
433	La carga de la enfermedad en España 2010: trastornos neurológicos, mentales y relacionados con el consumo de sustancias. Revista De Psiquiatría Y Salud Mental, 2015, 8, 207-217.	1.0	17
434	Risk factors and topographies for self-harm behaviour in a sample of adults with intellectual developmental disorders. Journal of Intellectual Disability Research, 2018, 62, 1018-1029.	1.2	17
435	Epidemiology of panic attacks, panic disorder and the moderating role of age: Results from a population-based study. Journal of Affective Disorders, 2018, 241, 627-633.	2.0	17
436	Pain rates in general population for the period 1991–2015 and 10-years prediction: results from a multi-continent age-period-cohort analysis. Journal of Headache and Pain, 2020, 21, 52.	2.5	17
437	Course of depression and cognitive decline at 3-year follow-up: The role of age of onset.. Psychology and Aging, 2019, 34, 475-485.	1.4	17
438	Effectiveness of olanzapine monotherapy and olanzapine combination treatment in the long term following acute mania – Results of a two year observational study in bipolar disorder (EMBLEM). Journal of Affective Disorders, 2011, 131, 320-329.	2.0	16
439	Regional differences in treatment response and three year course of schizophrenia across the world. Journal of Psychiatric Research, 2012, 46, 856-864.	1.5	16
440	Cognitive profiles of three clusters of patients with a first-episode psychosis. Schizophrenia Research, 2013, 150, 151-156.	1.1	16
441	Degree of cognitive impairment and mortality: a 17-year follow-up in a community study. Epidemiology and Psychiatric Sciences, 2015, 24, 503-511.	1.8	16
442	Which Are the Most Burdensome Functioning Areas in Depression? A Cross-National Study. Frontiers in Psychology, 2016, 7, 1342.	1.1	16
443	Study of positive and negative feedback sensitivity in psychosis using the Wisconsin Card Sorting Test. Comprehensive Psychiatry, 2016, 68, 119-128.	1.5	16
444	Psychotic-like experiences and disordered eating in the English general population. Psychiatry Research, 2016, 241, 26-34.	1.7	16
445	Estimating the COVID-19 Spread Through Real-time Population Mobility Patterns: Surveillance in Low- and Middle-Income Countries. Journal of Medical Internet Research, 2021, 23, e22999.	2.1	16
446	Patterns and correlates of patient-reported helpfulness of treatment for common mental and substance use disorders in the WHO World Mental Health Surveys. World Psychiatry, 2022, 21, 272-286.	4.8	16
447	Alcohol abuse in developed and developing countries in the World Mental Health Surveys: Socially defined consequences or psychiatric disorder?. American Journal on Addictions, 2014, 23, 145-155.	1.3	15
448	Visual encoding impairment in patients with schizophrenia: Contribution of reduced working memory span, decreased processing speed, and affective symptoms.. Neuropsychology, 2015, 29, 17-24.	1.0	15
449	Weight Perception, Satisfaction, Control, and Low Energy Dietary Reporting in the US Adult Population: Results from the National Health and Nutrition Examination Survey 2007-2012. Journal of the Academy of Nutrition and Dietetics, 2016, 116, 579-589.	0.4	15
450	Spanish validation of the social stigma scale: Community Attitudes towards Mental Illness. Revista De Psiquiatría Y Salud Mental (English Edition), 2016, 9, 150-157.	0.2	15

#	ARTICLE	IF	CITATIONS
451	Real-world Effectiveness of Antipsychotics for the Treatment of Negative Symptoms in Patients with Schizophrenia with Predominantly Negative Symptoms. <i>Pharmacopsychiatry</i> , 2017, 50, 56-63.	1.7	15
452	Costs and Burden Associated With Loss of Labor Productivity in Informal Caregivers of People With Dementia. <i>Journal of Occupational and Environmental Medicine</i> , 2018, 60, 449-456.	0.9	15
453	A Cross-National Investigation of Hallucination-Like Experiences in 10 Countries: The E-CLECTIC Study. <i>Schizophrenia Bulletin</i> , 2019, 45, S43-S55.	2.3	15
454	Validation in Spanish population of the family objective and subjective burden interview (ECFOS-II) for relatives of patients with schizophrenia. <i>Actas Espanolas De Psiquiatria</i> , 2007, 35, 372-81.	0.1	15
455	Comparison of treatment discontinuation and hospitalization among nonadherent patients initiating depot or oral typical antipsychotic medications. <i>International Clinical Psychopharmacology</i> , 2012, 27, 275-282.	0.9	14
456	Factors Associated With Components of Arterial Pressure Among Older Individuals (the Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 547 Td (M Consumption. <i>Journal of Clinical Hypertension</i> , 2014, 16, 645-651.	1.0	14
457	Factors Related to Unemployment in Europe. A Cross-Sectional Study from the COURAGE Survey in Finland, Poland and Spain. <i>International Journal of Environmental Research and Public Health</i> , 2018, 15, 722.	1.2	14
458	Cognition and functionality in delusional disorder. <i>European Psychiatry</i> , 2019, 55, 52-60.	0.1	14
459	Assessment of service use patterns in out-patients with schizophrenia: a Spanish study. <i>Acta Psychiatrica Scandinavica</i> , 2006, 114, 12-18.	2.2	13
460	Open-label olanzapine treatment in bipolar I disorder: clinical and work functional outcomes. <i>Current Medical Research and Opinion</i> , 2006, 22, 961-966.	0.9	13
461	Observational study designs for bipolar disorder â€” What can they tell us about treatment in acute mania?. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2009, 33, 715-721.	2.5	13
462	Impact of depressive symptoms on subjective well-being: the importance of patient-reported outcomes in schizophrenia. <i>Patient Preference and Adherence</i> , 2011, 5, 471.	0.8	13
463	Estudio cualitativo sobre la autopercepci3n del estigma social en personas con esquizofrenia. <i>Revista De La Asociaci3n Espa±ola De NeuropsiquiatrAa</i> , 2011, 31, 477-489.	0.1	13
464	Differential 3-Year Effects of First- Versus Second-Generation Antipsychotics on Subjective Well-Being in Schizophrenia Using Marginal Structural Models. <i>Journal of Clinical Psychopharmacology</i> , 2011, 31, 226-230.	0.7	13
465	Gender differences in mental health expectancies in early- and midlife in six European countries. <i>British Journal of Psychiatry</i> , 2013, 202, 294-300.	1.7	13
466	Do concomitant pain symptoms in patients with major depression affect quality of life even when taking into account baseline depression severity?. <i>Patient Preference and Adherence</i> , 2013, 7, 463.	0.8	13
467	Mood and anxiety disorders across the adult lifespan: a European perspective. <i>Psychological Medicine</i> , 2014, 44, 707-722.	2.7	13
468	Antidepressant medication treatment patterns in Asian patients with major depressive disorder. <i>Patient Preference and Adherence</i> , 2015, 9, 421.	0.8	13

#	ARTICLE	IF	CITATIONS
469	Development and validation of the Self-Stigma Questionnaire (SSQ) for people with schizophrenia and its relation to social functioning. <i>Comprehensive Psychiatry</i> , 2015, 62, 93-99.	1.5	13
470	Positive Affect Is Inversely Associated with Mortality in Individuals without Depression. <i>Frontiers in Psychology</i> , 2016, 7, 1040.	1.1	13
471	How to deal with missing longitudinal data in cost of illness analysis in Alzheimer's disease? suggestions from the GERAS observational study. <i>BMC Medical Research Methodology</i> , 2016, 16, 83.	1.4	13
472	National funding for mental health research in Finland, France, Spain and the United Kingdom. <i>European Neuropsychopharmacology</i> , 2017, 27, 892-899.	0.3	13
473	Measurement invariance of the WHOQOL-AGE questionnaire across three European countries. <i>Quality of Life Research</i> , 2018, 27, 1015-1025.	1.5	13
474	Fruit and Vegetable Consumption and Potential Moderators Associated with All-Cause Mortality in a Representative Sample of Spanish Older Adults. <i>Nutrients</i> , 2019, 11, 1794.	1.7	13
475	Moving more, ageing happy: findings from six low- and middle-income countries. <i>Age and Ageing</i> , 2021, 50, 488-497.	0.7	13
476	Relationship between sexual orientation and psychotic experiences in the general population in England. <i>Psychological Medicine</i> , 2021, 51, 138-146.	2.7	13
477	Mediators of the socioeconomic status and life satisfaction relationship in older adults: a multi-country structural equation modeling approach. <i>Aging and Mental Health</i> , 2021, 25, 585-592.	1.5	13
478	Socioeconomic inequalities in physical and cognitive functioning: cross-sectional evidence from 37 cohorts across 28 countries in the ATHLOS project. <i>Journal of Epidemiology and Community Health</i> , 2021, 75, 980-986.	2.0	13
479	The association of detachment with affective disorder symptoms during the COVID-19 lockdown: The role of living situation and social support. <i>Journal of Affective Disorders</i> , 2021, 292, 464-470.	2.0	13
480	Un estudio empírico de los correlatos psicosociales y clínicos del trastorno delirante: el estudio DELIREMP. <i>Revista De Psiquiatría Y Salud Mental</i> , 2009, 2, 72-82.	1.0	12
481	Association of childhood adversities with the first onset of mental disorders in Spain: results from the ESEMeD project. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2013, 48, 371-384.	1.6	12
482	Functional similarity of facial emotion processing between people with a first episode of psychosis and healthy subjects. <i>Schizophrenia Research</i> , 2013, 149, 35-41.	1.1	12
483	Subclinical psychosis and pain in an English national sample: The role of common mental disorders. <i>Schizophrenia Research</i> , 2016, 175, 209-215.	1.1	12
484	Real-world outcomes in patients with depression treated with duloxetine or a selective serotonin reuptake inhibitor in Asia. <i>Asia-Pacific Psychiatry</i> , 2016, 8, 51-59.	1.2	12
485	Measurement invariance of the Spanish Launay-Slade Hallucinations Scale-Extended version between putatively healthy controls and people diagnosed with a mental disorder. <i>International Journal of Methods in Psychiatric Research</i> , 2018, 27, e1741.	1.1	12
486	Predictive capacity of prodromal symptoms in first-episode psychosis of recent onset. <i>Microbial Biotechnology</i> , 2019, 13, 414-424.	0.9	12

#	ARTICLE	IF	CITATIONS
487	FOXP2 expression and gray matter density in the male brains of patients with schizophrenia. <i>Brain Imaging and Behavior</i> , 2021, 15, 1403-1411.	1.1	12
488	Informal caregiving and physical activity among 204,315 adults in 38 low- and middle-income countries: A cross-sectional study. <i>Preventive Medicine</i> , 2020, 132, 106007.	1.6	12
489	Four symptom dimensions in outpatients with schizophrenia. <i>Comprehensive Psychiatry</i> , 2006, 47, 384-388.	1.5	11
490	Sexual Intercourse Frequency in Men Presenting for Treatment of Erectile Dysfunction: Results from the Pan-European Erectile Dysfunction Observational Study. <i>Journal of Sex and Marital Therapy</i> , 2007, 33, 3-18.	1.0	11
491	An updated global picture of cigarette smoking persistence among adults. <i>Journal of Epidemiology and Global Health</i> , 2012, 2, 135.	1.1	11
492	Spanish validation of the Premorbid Adjustment Scale (PAS-S). <i>Comprehensive Psychiatry</i> , 2013, 54, 187-194.	1.5	11
493	The Role of Energy Balance in Successful Aging Among Elderly Individuals. <i>Journal of Aging and Health</i> , 2015, 27, 1375-1391.	0.9	11
494	Comparison of clinical outcomes with orodispersible versus standard oral olanzapine tablets in nonadherent patients with schizophrenia or bipolar disorder. <i>Patient Preference and Adherence</i> , 2017, Volume 11, 1019-1025.	0.8	11
495	Sociodemographic Factors Associated With Changes in Successful Aging in Spain: A Follow-Up Study. <i>Journal of Aging and Health</i> , 2018, 30, 1244-1262.	0.9	11
496	Association between intelligence quotient and violence perpetration in the English general population. <i>Psychological Medicine</i> , 2019, 49, 1316-1323.	2.7	11
497	Can the Positive and Negative Syndrome scale (PANSS) differentiate treatment-resistant from non-treatment-resistant schizophrenia? A factor analytic investigation based on data from the Pattern cohort study. <i>Psychiatry Research</i> , 2019, 276, 210-217.	1.7	11
498	Determinants of Subjective Wellbeing Trajectories in Older Adults: A Growth Mixture Modeling Approach. <i>Journal of Happiness Studies</i> , 2021, 22, 709-726.	1.9	11
499	Episodic Memory and Verbal Fluency Tasks: Normative Data from Nine Nationally Representative Samples. <i>Journal of the International Neuropsychological Society</i> , 2021, 27, 89-98.	1.2	11
500	Social functioning in schizophrenia: what is the influence of gender?. <i>European Journal of Psychiatry</i> , 2007, 21, .	0.7	11
501	Methodological approach for assessing the cost-effectiveness of treatments using longitudinal observational data: The SOHO study. <i>International Journal of Technology Assessment in Health Care</i> , 2006, 22, 460-468.	0.2	10
502	Seeking Help for Mental Health Problems outside the Conventional Health Care System: Results from the European Study of the Epidemiology of Mental Disorders (ESEMED). <i>Canadian Journal of Psychiatry</i> , 2010, 55, 586-597.	0.9	10
503	Abnormal functioning of the semantic network in schizophrenia patients with thought disorganization. An exemplar production task. <i>Psychiatry Research</i> , 2013, 205, 1-6.	1.7	10
504	Decreased processing speed might account for working memory span deficit in schizophrenia, and might mediate the associations between working memory span and clinical symptoms. <i>European Psychiatry</i> , 2014, 29, 473-478.	0.1	10

#	ARTICLE	IF	CITATIONS
505	Determinants of mobility in populations of older adults: Results from a cross-sectional study in Finland, Poland and Spain. <i>Maturitas</i> , 2018, 115, 84-91.	1.0	10
506	Perceived helpfulness of treatment for generalized anxiety disorder: a World Mental Health Surveys report. <i>BMC Psychiatry</i> , 2021, 21, 392.	1.1	10
507	The social, psychopathological and consumer context of rate of symptom improvement in acute mania. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2007, 42, 631-638.	1.6	9
508	Translation of Randomised Controlled Trial Findings into Clinical Practice: Comparison of Olanzapine and Valproate in the EMBLEM Study. <i>Pharmacopsychiatry</i> , 2009, 42, 145-152.	1.7	9
509	Reasons and Outcomes of Olanzapine Dose Adjustments in the Outpatient Treatment of Schizophrenia. <i>Pharmacopsychiatry</i> , 2009, 42, 135-140.	1.7	9
510	Fluoxetine and imipramine: are there differences in cost-utility for depression in primary care?. <i>Journal of Evaluation in Clinical Practice</i> , 2009, 15, 195-203.	0.9	9
511	Gender differences in outcomes of acute mania: a 12-month follow-up study. <i>Archives of Women's Mental Health</i> , 2011, 14, 107-113.	1.2	9
512	Antipsychotic Monotherapy and Polypharmacy in the Treatment of Outpatients With Schizophrenia in the European Schizophrenia Outpatient Health Outcomes Study. <i>Journal of Nervous and Mental Disease</i> , 2012, 200, 637-643.	0.5	9
513	Subjective well-being in schizophrenia: A randomised controlled open-label 12-month non-inferiority study comparing quetiapine XR with risperidone (RECOVER). <i>European Neuropsychopharmacology</i> , 2013, 23, 1257-1269.	0.3	9
514	The joint association of depression and cognitive function with severe disability among community-dwelling older adults in Finland, Poland and Spain. <i>Experimental Gerontology</i> , 2016, 76, 39-45.	1.2	9
515	Development and validation of the Questionnaire of Stressful Life Events (QSLE). <i>Journal of Psychiatric Research</i> , 2017, 95, 213-223.	1.5	9
516	Furthering the Evidence of the Effectiveness of Employment Strategies for People with Mental Disorders in Europe: A Systematic Review. <i>International Journal of Environmental Research and Public Health</i> , 2018, 15, 838.	1.2	9
517	European mental health research resources: Picture and recommendations of the ROAMER project. <i>European Neuropsychopharmacology</i> , 2019, 29, 179-194.	0.3	9
518	Predictors of pain in general ageing populations: results from a multi-country analysis based on ATHLOS harmonized database. <i>Journal of Headache and Pain</i> , 2020, 21, 45.	2.5	9
519	Associations between sedentary behavior and happiness: An analysis of influential factors among middle-aged and older adults from six low- and middle-income countries. <i>Maturitas</i> , 2021, 143, 157-164.	1.0	9
520	From childhood financial hardship to late-life depression: socioeconomic pathways. <i>Aging and Mental Health</i> , 2021, 25, 86-93.	1.5	9
521	Perceived helpfulness of treatment for specific phobia: Findings from the World Mental Health Surveys. <i>Journal of Affective Disorders</i> , 2021, 288, 199-209.	2.0	9
522	Analysis of burden in caregivers of people with Alzheimer's disease using self-report and supervision hours. <i>Journal of Nutrition, Health and Aging</i> , 0, . .	1.5	9

#	ARTICLE	IF	CITATIONS
523	No significant association between COVID-19 diagnosis and the incidence of depression and anxiety disorder? A retrospective cohort study conducted in Germany. <i>Journal of Psychiatric Research</i> , 2022, 147, 79-84.	1.5	9
524	The effect of loneliness and social support on the course of major depressive disorder among adults aged 50 years and older: A longitudinal study. <i>Depression and Anxiety</i> , 2022, 39, 147-155.	2.0	9
525	Variabilidad en el gen COMT y modificaci3n del riesgo de esquizofrenia conferido por consumo de cannabis. <i>Revista De PsiquiatrAa Y Salud Mental</i> , 2009, 2, 89-94.	1.0	8
526	Prevalence and correlates of respiratory and non-respiratory panic attacks in the general population. <i>Journal of Affective Disorders</i> , 2011, 131, 330-338.	2.0	8
527	Measuring use of services for mental health problems in epidemiological surveys. <i>International Journal of Methods in Psychiatric Research</i> , 2011, 20, 182-191.	1.1	8
528	Clinical consequences of switching from olanzapine to risperidone and vice versa in outpatients with schizophrenia: 36-month results from the worldwide schizophrenia outpatients health outcomes (W-SOHO) study. <i>BMC Psychiatry</i> , 2012, 12, 218.	1.1	8
529	Factors Predicting Hostility in Outpatients With Schizophrenia. <i>Journal of Nervous and Mental Disease</i> , 2013, 201, 464-470.	0.5	8
530	Association of auditory-verbal and visual hallucinations with impaired and improved recognition of colored pictures.. <i>Neuropsychology</i> , 2015, 29, 667-674.	1.0	8
531	Impact of Negative Symptoms on Quality of Life in Patients with Schizophrenia. <i>Value in Health</i> , 2015, 18, A836-A837.	0.1	8
532	Measurement Invariance of the Day Reconstruction Method: Results from the COURAGE in Europe Project. <i>Journal of Happiness Studies</i> , 2016, 17, 1769-1787.	1.9	8
533	Health-related quality of life in outpatients with schizophrenia: factors that determine changes over time. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2018, 53, 239-248.	1.6	8
534	The impact of socioeconomic status on the association between biomedical and psychosocial well-being and all-cause mortality in older Spanish adults. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2018, 53, 259-268.	1.6	8
535	Trial exclusion criteria and their impact on the estimation of antipsychotic drugs effect: A case study using the SOHO database. <i>Schizophrenia Research</i> , 2018, 193, 146-153.	1.1	8
536	Attention deficit hyperactivity disorder symptoms and disordered eating in the English general population. <i>International Journal of Eating Disorders</i> , 2018, 51, 942-952.	2.1	8
537	<p></p>Correlation of Health-Related Quality of Life in Clinically Stable Outpatients with Schizophrenia</p>. <i>Neuropsychiatric Disease and Treatment</i> , 2019, Volume 15, 3475-3486.	1.0	8
538	Metabolic dysregulation in older adults with depression and loneliness: The ATHLOS study. <i>Psychoneuroendocrinology</i> , 2021, 123, 104918.	1.3	8
539	Sarcopenia and Mild Cognitive Impairment in Older Adults from Six Low- and Middle-Income Countries. <i>Journal of Alzheimer's Disease</i> , 2021, 82, 1745-1754.	1.2	8
540	Antidepressant use in low- middle- and high-income countries: a World Mental Health Surveys report. <i>Psychological Medicine</i> , 2023, 53, 1583-1591.	2.7	8

#	ARTICLE	IF	CITATIONS
541	Kynurenine pathway in post-mortem prefrontal cortex and cerebellum in schizophrenia: relationship with monoamines and symptomatology. <i>Journal of Neuroinflammation</i> , 2021, 18, 198.	3.1	8
542	Assessment instruments: standardization of the European Service Mapping Schedule (ESMS) in Spain. <i>Acta Psychiatrica Scandinavica</i> , 2000, 102, 24-32.	2.2	8
543	Gender differences in incipient psychosis. <i>European Journal of Psychiatry</i> , 2010, 24, .	0.7	8
544	Psychiatric disorders in primary health care clinics one year after a major latin american disaster. <i>Stress and Health</i> , 1991, 7, 25-32.	0.7	7
545	Evaluation of Affect in Mexico and Spain: Psychometric Properties and Usefulness of an Abbreviated Version of the Day Reconstruction Method. <i>Journal of Happiness Studies</i> , 2014, 15, 915-935.	1.9	7
546	Impact of painful physical symptoms on depression outcomes in elderly Asian patients. <i>International Psychogeriatrics</i> , 2015, 27, 305-312.	0.6	7
547	Determinants of the components of arterial pressure among older adults " The role of anthropometric and clinical factors: A multi-continent study. <i>Atherosclerosis</i> , 2015, 238, 240-249.	0.4	7
548	Clinical outcomes of patients with major depressive disorder treated with either duloxetine, escitalopram, fluoxetine, paroxetine, or sertraline. <i>Neuropsychiatric Disease and Treatment</i> , 2018, Volume 14, 2473-2484.	1.0	7
549	Cross-cultural comparison of symptom networks in late-life major depressive disorder: Yoruba Africans and the Spanish Population. <i>International Journal of Geriatric Psychiatry</i> , 2020, 35, 1060-1068.	1.3	7
550	Perceived helpfulness of bipolar disorder treatment: Findings from the World Health Organization World Mental Health Surveys. <i>Bipolar Disorders</i> , 2021, 23, 565-583.	1.1	7
551	Is there a combined effect of depression and cognitive reserve on cognitive function? Findings from a population-based study. <i>Psychology and Health</i> , 2022, 37, 1132-1147.	1.2	7
552	Multimorbidity and Subjective Cognitive Complaints: Findings from 48 Low- and Middle-Income Countries of the World Health Survey 2002-2004. <i>Journal of Alzheimer's Disease</i> , 2021, 81, 1737-1747.	1.2	7
553	The Burden of schizophrenia. <i>European Neuropsychopharmacology</i> , 2022, 57, 33-35.	0.3	7
554	Perceived helpfulness of treatment for social anxiety disorder: findings from the WHO World Mental Health Surveys. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2022, 57, 2079-2095.	1.6	7
555	Predictors of engagement with remote sensing technologies for symptom measurement in Major Depressive Disorder. <i>Journal of Affective Disorders</i> , 2022, 310, 106-115.	2.0	7
556	Unit costs in international economic evaluations: resource costing of the Schizophrenia Outpatient Health Outcomes Study. <i>Acta Psychiatrica Scandinavica</i> , 2003, 107, 41-47.	2.2	6
557	Evaluaci3n telef3nica de los sistemas de activaci3n e inhibici3n conductual. <i>Revista De Psiquiatr3a Y Salud Mental</i> , 2009, 2, 66-71.	1.0	6
558	Association between annual earnings and mental disorders in Spain: individual and societal-level estimates. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2012, 47, 1717-1725.	1.6	6

#	ARTICLE	IF	CITATIONS
559	Patient-Reported Cognitive Dysfunction Negatively Impacts Functioning in Patients with Major Depressive Disorder – Preliminary Findings from the Perform Study. <i>Value in Health</i> , 2013, 16, A543-A544.	0.1	6
560	Predictors of psychiatric hospitalization during 6 months of maintenance treatment with olanzapine long-acting injection: post hoc analysis of a randomized, double-blind study. <i>BMC Psychiatry</i> , 2013, 13, 224.	1.1	6
561	Patient characteristics associated with treatment initiation among paediatric patients with Attention-Deficit/Hyperactivity Disorder symptoms in a naturalistic setting in Central Europe and East Asia. <i>BMC Psychiatry</i> , 2014, 14, 304.	1.1	6
562	Desarrollo profesional en investigación traslacional en neurociencias y salud mental: educación y formación dentro del Centro de Investigación Biomédica en Red en Salud Mental. <i>Revista De Psiquiatría Y Salud Mental</i> , 2015, 8, 65-74.	1.0	6
563	Associations between DSM-IV mental disorders and subsequent onset of arthritis. <i>Journal of Psychosomatic Research</i> , 2016, 82, 11-16.	1.2	6
564	A novel approach to measuring response and remission in schizophrenia in clinical trials. <i>Schizophrenia Research</i> , 2017, 190, 123-128.	1.1	6
565	Child death and maternal psychosis-like experiences in 44 low- and middle-income countries: The role of depression. <i>Schizophrenia Research</i> , 2017, 183, 41-46.	1.1	6
566	Health conditions and role limitation in three European Regions: a public-health perspective. <i>Gaceta Sanitaria</i> , 2017, 31, 2-10.	0.6	6
567	The association between problem gambling and psychotic experiences: Findings from the Adult Psychiatric Morbidity Survey 2007. <i>Schizophrenia Research</i> , 2018, 201, 79-84.	1.1	6
568	Characteristics of patients with depression initiating or switching antidepressant treatment: baseline analyses of the PERFORM cohort study. <i>BMC Psychiatry</i> , 2018, 18, 80.	1.1	6
569	Development of a Combined Sensory-Cognitive Measure Based on the Common Cause Hypothesis: Heterogeneous Trajectories and Associated Risk Factors. <i>Gerontologist</i> , The, 2020, 60, e357-e366.	2.3	6
570	Association between intelligence quotient and disability: The role of socioeconomic status. <i>Annals of Physical and Rehabilitation Medicine</i> , 2020, 63, 296-301.	1.1	6
571	The relationship between all-cause mortality and depression in different gender and age groups of the Spanish population. <i>Journal of Affective Disorders</i> , 2020, 266, 424-428.	2.0	6
572	Cognitive function associated with different diagnoses of anxiety disorders over the lifespan: Results from a Spanish representative sample. <i>Journal of Anxiety Disorders</i> , 2020, 75, 102296.	1.5	6
573	Associations between cataract and multimorbidity: a cross-sectional study of 23,089 adults from Spain. <i>Eye</i> , 2021, 35, 791-798.	1.1	6
574	The Relationship Between Plasma Haloperidol Concentrations and Clinical Results. <i>Archives of General Psychiatry</i> , 1996, 53, 1167.	13.8	5
575	Effects of common mental disorders and physical conditions on role functioning in Spain. <i>Gaceta Sanitaria</i> , 2013, 27, 480-486.	0.6	5
576	Changes in adherence and treatment costs following initiation of oral or depot typical antipsychotics among previously non-adherent patients with schizophrenia. <i>Human Psychopharmacology</i> , 2013, 28, 438-446.	0.7	5

#	ARTICLE	IF	CITATIONS
577	Long-term functional improvements in the 2-year treatment of schizophrenia outpatients with olanzapine long-acting injection. <i>Neuropsychiatric Disease and Treatment</i> , 2014, 10, 1125.	1.0	5
578	Health-related quality of life in the Cambridge City over-75s Cohort (CC75C): development of a dementia-specific scale and descriptive analyses. <i>BMC Geriatrics</i> , 2014, 14, 18.	1.1	5
579	The Spanish Burden of Disease 2010: Neurological, mental and substance use disorders. <i>Revista De Psiquiatría Y Salud Mental (English Edition)</i> , 2015, 8, 207-217.	0.2	5
580	Chinese patients with major depression: Do concomitant pain symptoms affect quality of life independently of severity of depression?. <i>International Journal of Psychiatry in Clinical Practice</i> , 2015, 19, 174-181.	1.2	5
581	Psychometric Properties of Spanish Adaptation of the PDD-MRS Scale in Adults with Intellectual Developmental Disorders: The EVTEA-DI Scale. <i>Journal of Autism and Developmental Disorders</i> , 2018, 48, 1566-1578.	1.7	5
582	Health-related quality of life associated with different symptoms in women and in men who suffer from schizophrenia. <i>Archives of Women's Mental Health</i> , 2019, 22, 357-365.	1.2	5
583	CIBERSAM: Ten years of collaborative translational research in mental disorders. <i>Revista De Psiquiatría Y Salud Mental (English Edition)</i> , 2019, 12, 1-8.	0.2	5
584	The relationship of childhood sexual and physical abuse with adulthood disability. <i>Annals of Physical and Rehabilitation Medicine</i> , 2020, 63, 332-339.	1.1	5
585	The Association Between Substance Use Disorder and Depression During the COVID-19 Lockdown in Spain and the Moderating Role of Social Support: a Cross-Sectional Study. <i>International Journal of Mental Health and Addiction</i> , 2023, 21, 1157-1167.	4.4	5
586	Prevalence of and factors associated with long-term sick leave in working-age adults with chronic low back pain in Germany. <i>International Archives of Occupational and Environmental Health</i> , 2022, 95, 1549-1556.	1.1	5
587	Cohort Profile: The Spanish Longitudinal Study on Ageing and Health (<i>Edad Con Salud</i>). <i>International Journal of Epidemiology</i> , 2022, 51, e189-e199.	0.9	5
588	Methimazole treatment reduces cardiac hypertrophy and mortality without a concomitant reduction in blood pressure in established Goldblatt two-kidney one clip hypertension. <i>Experientia</i> , 1992, 48, 755-758.	1.2	4
589	Clinical consequences of switching antipsychotic drugs in outpatients with schizophrenia: 36-month results from the European Schizophrenia Outpatient Health Outcomes study. <i>International Clinical Psychopharmacology</i> , 2008, 23, 203-208.	0.9	4
590	An empirical study of psychosocial and clinical correlates of delusional disorder: the DELIREMP study. <i>Revista De Psiquiatría Y Salud Mental (English Edition)</i> , 2009, 2, 72-82.	0.2	4
591	Comparison of olanzapine and risperidone in the EMBLEM Study: translation of randomized controlled trial findings into clinical practice. <i>International Clinical Psychopharmacology</i> , 2010, 25, 257-263.	0.9	4
592	Schizophrenia Outpatient Health Outcomes study: twelve-month findings. <i>Journal of Pragmatic and Observational Research</i> , 2012, 3, 27.	1.1	4
593	Usefulness of 2 Questions About Age and Year of Birth in the Case-Finding of Dementia. <i>Journal of the American Medical Directors Association</i> , 2013, 14, 627.e7-627.e12.	1.2	4
594	Comparing Resource Use In Alzheimer's Disease Across Three European Countries - 18-Month Results of The Geras Study. <i>Value in Health</i> , 2015, 18, A759-A760.	0.1	4

#	ARTICLE	IF	CITATIONS
595	Phosphorylation of Transcription Factor Specificity Protein 4 Is Increased in Peripheral Blood Mononuclear Cells of First-Episode Psychosis. <i>PLoS ONE</i> , 2015, 10, e0125115.	1.1	4
596	To be happy and behave in a healthier way. A longitudinal study about gender differences in the older population. <i>Psychology and Health</i> , 2021, , 1-17.	1.2	4
597	The role of social support, detachment, and depressive and anxiety symptoms in suicidal thoughts and behaviours during the Covid-19 lockdown: Potential pathways. <i>Current Psychology</i> , 2021, , 1-12.	1.7	4
598	Are retired people higher in experiential wellbeing than working older adults? A time use approach.. <i>Emotion</i> , 2020, 20, 1411-1422.	1.5	4
599	Should Unexplained Painful Physical Symptoms be Considered within the Spectrum of Depressive Symptoms?. <i>Clinical Practice and Epidemiology in Mental Health</i> , 2015, 11, 130-139.	0.6	4
600	Assessment of remission in schizophrenia with the CGI and CGIâ€‘SCH scales. <i>Acta Psychiatrica Scandinavica</i> , 2008, 117, 156-156.	2.2	3
601	Functioning in patients with major depression treated with duloxetine or a selective serotonin reuptake inhibitor in East Asia. <i>Neuropsychiatric Disease and Treatment</i> , 2016, 12, 383.	1.0	3
602	Factors associated with and impact of pain persistence in Asian patients with depression: a 3-month, prospective observational study. <i>International Journal of Psychiatry in Clinical Practice</i> , 2017, 21, 29-35.	1.2	3
603	Frailty and health status of older individuals in three European countries: The COURAGE cross-sectional study. <i>Experimental Gerontology</i> , 2018, 106, 137-144.	1.2	3
604	Impact of pain and remission in the functioning of patients with depression in <sc>M</sc>ainland <sc>C</sc>hina, <sc>T</sc>aiwan, and <sc>H</sc>ong <sc>K</sc>ong. <i>Asia-Pacific Psychiatry</i> , 2018, 10, e12295.	1.2	3
605	Functioning profiles in a nationally representative cohort of Spanish older adults: A latent class study. <i>Health and Social Care in the Community</i> , 2020, 28, 2190-2198.	0.7	3
606	The new Horizon Europe programme 2021â€‘2028: Should the gap between the burden of mental disorders and the funding of mental health research be filled?. <i>European Journal of Psychiatry</i> , 2020, 34, 44-46.	0.7	3
607	Trajectories of Immediate and Delayed Verbal Memory in the Spanish General Population of Middle-aged and Older Adults. <i>Brain Sciences</i> , 2020, 10, 249.	1.1	3
608	Selective serotonin reuptake inhibitors: are there differences in effectiveness in primary care?. <i>Primary Care and Community Psychiatry</i> , 2007, 11, 113-120.	0.4	3
609	Perceived helpfulness of treatment for alcohol use disorders: Findings from the World Mental Health Surveys. <i>Drug and Alcohol Dependence</i> , 2021, 229, 109158.	1.6	3
610	Previous disorders and depression outcomes in individuals with 12-month major depressive disorder in the World Mental Health surveys. <i>Epidemiology and Psychiatric Sciences</i> , 2021, 30, e70.	1.8	3
611	A divisive hierarchical clustering methodology for enhancing the ensemble prediction power in large scale population studies: the ATHLOS project. <i>Health Information Science and Systems</i> , 2022, 10, 6.	3.4	3
612	PHS50 Health Care Resource Use and Cost in Alzheimer's Disease in Three European Countries â€‘ Baseline Findings of the GERAS Study. <i>Value in Health</i> , 2012, 15, A527.	0.1	2

#	ARTICLE	IF	CITATIONS
613	Relationship of Insight with Medication Adherence and the Impact on Outcomes in Patients with Schizophrenia and Bipolar Disorder: Results From A 1-Year European Outpatient Observational Study. <i>Value in Health</i> , 2014, 17, A455.	0.1	2
614	Partial Likelihood Estimation of IRT Models with Censored Lifetime Data: An Application to Mental Disorders in the ESEMeD Surveys. <i>Psychometrika</i> , 2014, 79, 470-488.	1.2	2
615	Health-related quality of life in patients with depression treated with duloxetine or a selective serotonin reuptake inhibitor in a naturalistic outpatient setting. <i>Patient Preference and Adherence</i> , 2015, 9, 1481.	0.8	2
616	Impact of anxiety symptoms on outcomes of depression: an observational study in Asian patients. <i>Neuropsychiatric Disease and Treatment</i> , 2016, 12, 795.	1.0	2
617	Specificity proteins 1 and 4, hippocampal volume and first-episode psychosis. <i>British Journal of Psychiatry</i> , 2016, 208, 591-592.	1.7	2
618	Is Parental Longevity Associated With the Cardiovascular Risk and the Successful Aging of Their Offspring? Results From the Multinational MEDIS Study. <i>Angiology</i> , 2017, 68, 124-131.	0.8	2
619	The "RCT augmentation" a novel simulation method to add patient heterogeneity into phase III trials. <i>BMC Medical Research Methodology</i> , 2018, 18, 75.	1.4	2
620	Retrospective analysis of patients with advanced or metastatic gastric cancer in Colombia. <i>Journal of Medical Economics</i> , 2019, 22, 891-900.	1.0	2
621	The role of ageing in the wish to be dead: disentangling age, period and cohort effects in suicide ideation in European population. <i>Epidemiology and Psychiatric Sciences</i> , 2021, 30, e17.	1.8	2
622	Association of objective and subjective far vision impairment with perceived stress among older adults in six low- and middle-income countries. <i>Eye</i> , 2022, 36, 1274-1280.	1.1	2
623	Functional Impairment and Painful Physical Symptoms in Patients with Major Depressive Disorder Treated with Antidepressants: Real-World Evidence from the Middle East. <i>Clinical Practice and Epidemiology in Mental Health</i> , 2017, 13, 145-155.	0.6	2
624	Late-life disability trajectories in Yoruba Nigerians and the Spanish population: a state space model in continuous time. <i>Aging and Mental Health</i> , 2022, 26, 2447-2453.	1.5	2
625	Types of Racism and Twitter Users'™ Responses Amid the COVID-19 Outbreak: Content Analysis. <i>JMIR Formative Research</i> , 2022, 6, e29183.	0.7	2
626	Olanzapine versus haloperidol as initial treatment in acute mania: Prospective comparison of work functional outcomes. <i>European Neuropsychopharmacology</i> , 2002, 12, 307.	0.3	1
627	Clinical and functional status over the first three months of treatment: Results from the schizophrenia outpatient health outcomes (SOHO) study. <i>Schizophrenia Research</i> , 2003, 60, 282.	1.1	1
628	0428 EFFECTIVENESS OF ANTIPSYCHOTIC TREATMENT IN NEVER TREATED PATIENTS WITH SCHIZOPHRENIA: 36-MONTH RESULTS FROM THE SCHIZOPHRENIA OUTPATIENTS HEALTH OUTCOMES (SOHO) STUDY. <i>Schizophrenia Research</i> , 2006, 86, S131-S132.	1.1	1
629	Evaluation of behavioural activation and inhibition systems by telephone. <i>Revista De Psiquiatria Y Salud Mental (English Edition)</i> , 2009, 2, 66-71.	0.2	1
630	The Burden of Treatment Change in Major Depressive Disorder: Comparison of Switch Versus Non-Switch Patients in the Perform Study. <i>Value in Health</i> , 2013, 16, A543.	0.1	1

#	ARTICLE	IF	CITATIONS
631	Comparative Effectiveness in Terms of Treatment Discontinuation of Orodispersable Versus. Standard Oral Olanzapine Tablets in Non-Adherent Patients: Results from a 1-Year European Outpatient Observational Study. Value in Health, 2014, 17, A766.	0.1	1
632	How Useful are Eq-5d And Zbi In Assessing The Impact of Caring for Alzheimerâ€™s Disease Patients?. Value in Health, 2015, 18, A760.	0.1	1
633	Sibship size, birth order and psychotic experiences: Evidence from 43 low- and middle-income countries. Schizophrenia Research, 2018, 201, 406-412.	1.1	1
634	Associations of physical and psychiatric conditions with chronic fatigue syndrome in Germany: an exploratory case-control study. Psychological Medicine, 2022, 52, 780-786.	2.7	1
635	Transdiagnostic development of internalizing psychopathology throughout the life course up to age 45: a World Mental Health Surveys report. Psychological Medicine, 2020, , 1-10.	2.7	1
636	Determinants of verbal fluency trajectories among older adults from the English Longitudinal Study of Aging. Applied Neuropsychology Adult, 2021, , 1-10.	0.7	1
637	Personalidad internalizante y externalizante y efectos subjetivos en una muestra de adolescentes consumidores de cnnabis. Revista De Psicología De La Salud, 2016, 28, 231.	0.2	1
638	Retrospective analysis of patients with locally advanced or metastatic gastric cancer in Argentina. Medwave, 2019, 19, e7693-e7693.	0.2	1
639	Determinants of effective treatment coverage for major depressive disorder in the WHO World Mental Health Surveys. International Journal of Mental Health Systems, 2022, 16, .	1.1	1
640	Statistical models for population heterogeneity in epidemiological longitudinal studies: Person centred latent class analysis to identify casemix groups in the schizophrenia outpatient health outcomes study. European Neuropsychopharmacology, 2002, 12, 314.	0.3	0
641	The treatment of schizophrenia across Europe: Between country baseline differences in the schizophrenia outpatient health outcomes (SOHO) study. European Neuropsychopharmacology, 2002, 12, 317.	0.3	0
642	PMH27 12-MONTH COST-EFFECTIVENESS ANALYSIS OF ORAL ANTIPSYCHOTIC TREATMENTS IN PATIENTS WITH SCHIZOPHRENIA IN THE PAN-EUROPEAN SOHO (SCHIZOPHRENIA OUTPATIENT HEALTH OUTCOMES) STUDY. Value in Health, 2005, 8, A204.	0.1	0
643	PMH31 12 MONTH COST-UTILITY ANALYSIS OF ORAL ANTIPSYCHOTIC TREATMENTS IN PATIENTS WITH SCHIZOPHRENIA IN THE PAN-EUROPEAN SOHO (SCHIZOPHRENIA OUTPATIENT HEALTH OUTCOMES) STUDY. Value in Health, 2005, 8, A205.	0.1	0
644	P.2.e.006 Outcomes of acute mania: 12 months results from the European mania in bipolar longitudinal evaluation of medication (EMBLEM) study. European Neuropsychopharmacology, 2006, 16, S349.	0.3	0
645	P.3.a.042 Clinical course, social functioning, quality of life & medication tolerability in previously untreated patients with schizophrenia: 36-month results from the European schizophrenia outpatient health outcomes (SOHO) study. European Neuropsychopharmacology, 2006, 16, S387-S388.	0.3	0
646	PMH35 THE COST OF RELAPSE IN PATIENTS WITH SCHIZOPHRENIA IN THE PAN-EUROPEAN SOHO (SCHIZOPHRENIA OUTPATIENT HEALTH OUTCOMES) STUDY. Value in Health, 2006, 9, A75.	0.1	0
647	Number needed to treat (NNT) for all-cause medication discontinuation in catie compared to the schizophrenia outpatients health outcomes (SOHO) study. European Psychiatry, 2007, 22, S130.	0.1	0
648	Previously untreated patients with schizophrenia : The nnt for all causes of treatment discontinuation and the nnh for weight gain. European Psychiatry, 2008, 23, S166.	0.1	0

#	ARTICLE	IF	CITATIONS
649	Religion and Spiritual Advice as Surrogate Care Provision for Mental Health Problems in the ESEMED Survey. <i>European Psychiatry</i> , 2009, 24, .	0.1	0
650	PMH25 CLINICAL AND ECONOMIC CONSEQUENCES OF MEDICATION NONADHERENCE IN THE TREATMENT OF PATIENTS WITH A MANIC/ MIXED EPISODE OF BIPOLAR DISORDER: RESULTS FROM THE EUROPEAN MANIA IN BIPOLAR LONGITUDINAL EVALUATION OF MEDICATION (EMBLEM) STUDY. <i>Value in Health</i> , 2009, 12, A355.	0.1	0
651	PND48 Health State Values in Alzheimer'S Disease " Baseline Results From a Large Prospective Observational Study in Three European Countries (GERAS). <i>Value in Health</i> , 2012, 15, A554.	0.1	0
652	Clinical Outcomes of Patients with Major Depressive Disorder Treated with Either Duloxetine or Selective Serotonin Reuptake Inhibitors in Mexico. <i>Value in Health</i> , 2013, 16, A694.	0.1	0
653	P.3.a.009 SP1 and SP4 transcription factors are increased in the postmortem hippocampus of subjects with chronic schizophrenia. <i>European Neuropsychopharmacology</i> , 2013, 23, S426-S427.	0.3	0
654	Factors Associated with Pain Persistence in Patient with Depression During a 3 "Month Follow-Up Period. <i>Value in Health</i> , 2013, 16, A544.	0.1	0
655	Description of Agitation and Crisis Interventions in a Psychiatric Hospital in Spain: a Qualitative Study. <i>Value in Health</i> , 2014, 17, A463.	0.1	0
656	Use of Services and Cost of Agitation and Containment in Psychiatric Hospitals: a Systematic Review. <i>Value in Health</i> , 2014, 17, A460.	0.1	0
657	Predictors Pf Remission in the Treatment of Depression in the Middle East: Real-World Evidence From A 6-Month Prospective Observational Study. <i>Value in Health</i> , 2014, 17, A454.	0.1	0
658	Symtomatic Factors in Patients With Major Depressive Disorder (MDD): Results from an Observational Study. <i>Value in Health</i> , 2014, 17, A543.	0.1	0
659	How to deal with missing longitudinal data in cost of illness models in alzheimer's disease " suggestions from the geras study results. <i>Value in Health</i> , 2014, 17, A185.	0.1	0
660	The use of random-effects models to identify health care center-related characteristics modifying the effect of antipsychotic drugs. <i>Clinical Epidemiology</i> , 2017, Volume 9, 689-698.	1.5	0
661	Attention-Deficit Hyperactivity Disorder. , 0, , 195-208.		0
662	The Effects of Trauma Type, Timing, Accumulation, and Sequencing. , 0, , 153-162.		0
663	Enhancing the Human Health Status Prediction: The ATHLOS Project. <i>Applied Artificial Intelligence</i> , 2021, 35, 834-856.	2.0	0
664	Risk of all-cause mortality associated with chronic obstructive pulmonary disease and the role of healthy ageing trajectories: a population-based study of middle-aged and older adults. <i>BMJ Open</i> , 2021, 11, e050947.	0.8	0
665	The Use of Bayesian Hierarchical Methods to Assess Selection and Observer Bias in Observational Health Outcome Studies. <i>SSRN Electronic Journal</i> , 0, , .	0.4	0
666	Estudio SOHO (Variables Relacionadas con la Salud en Pacientes Ambulatorios con Esquizofrenia): resultados a los 3 aÑ±os de la suspensi³n del tratamiento antipsic³tico y factores cl³nicos relacionados con ella en EspaÑ±a. <i>European Psychiatry (Ed EspaÑ±ola)</i> , 2008, 15, 137-144.	0.0	0

#	ARTICLE	IF	CITATIONS
667	Estudio descriptivo de un registro de casos de trastorno delirante. European Psychiatry (Ed) Tj ETQq1 1 0.784314 rgBT /Overlock 10 TFE	0.0	0
668	Factor V Leiden and the 10-year incidence of depression: A retrospective cohort study conducted in Germany. Journal of Psychiatric Research, 2022, 146, 87-91.	1.5	0
669	Launayâ€“Slade Hallucination Scale-Extended: simplifying its interpretation. Psychosis, 0, , 1-10.	0.4	0
670	The associations of physical incapacity and wealth with remaining in paid employment after age 60 in five middle-income and high-income countries. Ageing and Society, 2023, 43, 2994-3017.	1.2	0
671	Psychotic experiences among informal caregivers: findings from 48 low- and middle-income countries. Social Psychiatry and Psychiatric Epidemiology, 0, , .	1.6	0