Josep Maria Haro

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/4303299/publications.pdf

Version: 2024-02-01

671 papers

99,466 citations

109 h-index 295 g-index

724 all docs

724 docs citations

times ranked

724

120049 citing authors

#	Article	IF	CITATIONS
1	The Association Between Substance Use Disorder and Depression During the COVID-19 Lockdown in Spain and the Moderating Role of Social Support: a Cross-Sectional Study. International Journal of Mental Health and Addiction, 2023, 21, 1157-1167.	4.4	5
2	Antidepressant use in low- middle- and high-income countries: a World Mental Health Surveys report. Psychological Medicine, 2023, 53, 1583-1591.	2.7	8
3	The associations of physical incapacity and wealth with remaining in paid employment after age 60 in five middle-income and high-income countries. Ageing and Society, 2023, 43, 2994-3017.	1.2	O
4	Associations of physical and psychiatric conditions with chronic fatigue syndrome in Germany: an exploratory case-control study. Psychological Medicine, 2022, 52, 780-786.	2.7	1
5	Toward measuring effective treatment coverage: critical bottlenecks in quality- and user-adjusted coverage for major depressive disorder. Psychological Medicine, 2022, 52, 1948-1958.	2.7	23
6	Is there a combined effect of depression and cognitive reserve on cognitive function? Findings from a population-based study. Psychology and Health, 2022, 37, 1132-1147.	1.2	7
7	Association of objective and subjective far vision impairment with perceived stress among older adults in six low- and middle-income countries. Eye, 2022, 36, 1274-1280.	1.1	2
8	Late-life disability trajectories in Yoruba Nigerians and the Spanish population: a state space model in continuous time. Aging and Mental Health, 2022, 26, 2447-2453.	1.5	2
9	Factor V Leiden and the 10-year incidence of depression: A retrospective cohort study conducted in Germany. Journal of Psychiatric Research, 2022, 146, 87-91.	1.5	O
10	No significant association between COVID-19 diagnosis and the incidence of depression and anxiety disorder? A retrospective cohort study conducted in Germany. Journal of Psychiatric Research, 2022, 147, 79-84.	1.5	9
11	The Burden of schizophrenia. European Neuropsychopharmacology, 2022, 57, 33-35.	0.3	7
12	The effect of loneliness and social support on the course of major depressive disorder among adults aged 50 years and older: A longitudinal study. Depression and Anxiety, 2022, 39, 147-155.	2.0	9
13	The Association Between Home Stay and Symptom Severity in Major Depressive Disorder: Preliminary Findings From a Multicenter Observational Study Using Geolocation Data From Smartphones. JMIR MHealth and UHealth, 2022, 10, e28095.	1.8	19
14	Types of Racism and Twitter Users' Responses Amid the COVID-19 Outbreak: Content Analysis. JMIR Formative Research, 2022, 6, e29183.	0.7	2
15	Remote Assessment of Disease and Relapse in Major Depressive Disorder (RADAR-MDD): recruitment, retention, and data availability in a longitudinal remote measurement study. BMC Psychiatry, 2022, 22, 136.	1.1	42
16	Prevalence of and factors associated with long-term sick leave in working-age adults with chronic low back pain in Germany. International Archives of Occupational and Environmental Health, 2022, 95, 1549-1556.	1.1	5
17	Perceived helpfulness of treatment for social anxiety disorder: findings from the WHO World Mental Health Surveys. Social Psychiatry and Psychiatric Epidemiology, 2022, 57, 2079-2095.	1.6	7
18	Longitudinal Relationships Between Depressive Symptom Severity and Phone-Measured Mobility: Dynamic Structural Equation Modeling Study. JMIR Mental Health, 2022, 9, e34898.	1.7	26

#	Article	IF	Citations
19	A divisive hierarchical clustering methodology for enhancing the ensemble prediction power in large scale population studies: the ATHLOS project. Health Information Science and Systems, 2022, 10, 6.	3.4	3
20	Patterns and correlates of patientâ€reported helpfulness of treatment for common mental and substance use disorders in the <scp>WHO</scp> World Mental Health Surveys. World Psychiatry, 2022, 21, 272-286.	4.8	16
21	Predictors of engagement with remote sensing technologies for symptom measurement in Major Depressive Disorder. Journal of Affective Disorders, 2022, 310, 106-115.	2.0	7
22	Cohort Profile: The Spanish Longitudinal Study on Ageing and Health (<i>Edad Con Salud</i>). International Journal of Epidemiology, 2022, 51, e189-e199.	0.9	5
23	Determinants of effective treatment coverage for major depressive disorder in the WHO World Mental Health Surveys. International Journal of Mental Health Systems, 2022, 16, .	1.1	1
24	Development of a common scale for measuring healthy ageing across the world: results from the ATHLOS consortium. International Journal of Epidemiology, 2021, 50, 880-892.	0.9	32
25	FOXP2 expression and gray matter density in the male brains of patients with schizophrenia. Brain Imaging and Behavior, 2021, 15, 1403-1411.	1.1	12
26	Patterns of care and dropout rates from outpatient mental healthcare in low-, middle- and high-income countries from the World Health Organization's World Mental Health Survey Initiative. Psychological Medicine, 2021, 51, 2104-2116.	2.7	21
27	Determinants of Subjective Wellbeing Trajectories in Older Adults: A Growth Mixture Modeling Approach. Journal of Happiness Studies, 2021, 22, 709-726.	1.9	11
28	Associations between cataract and multimorbidity: a cross-sectional study of 23,089 adults from Spain. Eye, 2021, 35, 791-798.	1.1	6
29	Effects of transient and chronic loneliness on major depression in older adults: A longitudinal study. International Journal of Geriatric Psychiatry, 2021, 36, 76-85.	1.3	37
30	Metabolic dysregulation in older adults with depression and loneliness: The ATHLOS study. Psychoneuroendocrinology, 2021, 123, 104918.	1.3	8
31	Associations between sedentary behavior and happiness: An analysis of influential factors among middle-aged and older adults from six low- and middle-income countries. Maturitas, 2021, 143, 157-164.	1.0	9
32	Mental health impact of the first wave of COVID-19 pandemic on Spanish healthcare workers: A large cross-sectional survey. Revista De PsiquiatrÃa Y Salud Mental, 2021, 14, 90-105.	1.0	133
33	Moving more, ageing happy: findings from six low- and middle-income countries. Age and Ageing, 2021, 50, 488-497.	0.7	13
34	Episodic Memory and Verbal Fluency Tasks: Normative Data from Nine Nationally Representative Samples. Journal of the International Neuropsychological Society, 2021, 27, 89-98.	1.2	11
35	From childhood financial hardship to late-life depression: socioeconomic pathways. Aging and Mental Health, 2021, 25, 86-93.	1.5	9
36	Relationship between sexual orientation and psychotic experiences in the general population in England. Psychological Medicine, 2021, 51, 138-146.	2.7	13

#	Article	IF	Citations
37	Mediators of the socioeconomic status and life satisfaction relationship in older adults: a multi-country structural equation modeling approach. Aging and Mental Health, 2021, 25, 585-592.	1.5	13
38	Global mortality from dementia: Application of a new method and results from the Global Burden of Disease Study 2019. Alzheimer's and Dementia: Translational Research and Clinical Interventions, 2021, 7, e12200.	1.8	53
39	Changes in depression and suicidal ideation under severe lockdown restrictions during the first wave of the COVID-19 pandemic in Spain: a longitudinal study in the general population. Epidemiology and Psychiatric Sciences, 2021, 30, e49.	1.8	44
40	The role of ageing in the wish to be dead: disentangling age, period and cohort effects in suicide ideation in European population. Epidemiology and Psychiatric Sciences, 2021, 30, e17.	1.8	2
41	Are environmental pollution and biodiversity levels associated to the spread and mortality of COVID-19? A four-month global analysis. Environmental Pollution, 2021, 271, 116326.	3.7	33
42	Thirty-day suicidal thoughts and behaviours in the Spanish adult general population during the first wave of the Spain COVID-19 pandemic. Epidemiology and Psychiatric Sciences, 2021, 30, e19.	1.8	32
43	Perceived helpfulness of bipolar disorder treatment: Findings from the World Health Organization World Mental Health Surveys. Bipolar Disorders, 2021, 23, 565-583.	1.1	7
44	Socioeconomic inequalities in physical and cognitive functioning: cross-sectional evidence from 37 cohorts across 28 countries in the ATHLOS project. Journal of Epidemiology and Community Health, 2021, 75, 980-986.	2.0	13
45	Relationship Between Major Depression Symptom Severity and Sleep Collected Using a Wristband Wearable Device: Multicenter Longitudinal Observational Study. JMIR MHealth and UHealth, 2021, 9, e24604.	1.8	35
46	Mental health impact of the first wave of COVID-19 pandemic on Spanish healthcare workers: A large cross-sectional survey. Revista De PsiquiatrÃa Y Salud Mental (English Edition), 2021, 14, 90-105.	0.2	21
47	Predicting Depressive Symptom Severity Through Individuals' Nearby Bluetooth Device Count Data Collected by Mobile Phones: Preliminary Longitudinal Study. JMIR MHealth and UHealth, 2021, 9, e29840.	1.8	26
48	Association between Food Insecurity and Sarcopenia among Adults Aged \hat{a} %¥65 Years in Low- and Middle-Income Countries. Nutrients, 2021, 13, 1879.	1.7	19
49	Sarcopenia and fall-related injury among older adults in five low- and middle-income countries. Experimental Gerontology, 2021, 147, 111262.	1.2	20
50	Determinants of verbal fluency trajectories among older adults from the English Longitudinal Study of Aging. Applied Neuropsychology Adult, 2021, , 1-10.	0.7	1
51	Estimating the COVID-19 Spread Through Real-time Population Mobility Patterns: Surveillance in Lowand Middle-Income Countries. Journal of Medical Internet Research, 2021, 23, e22999.	2.1	16
52	Multimorbidity and Subjective Cognitive Complaints: Findings from 48 Low- and Middle-Income Countries of the World Health Survey 2002–2004. Journal of Alzheimer's Disease, 2021, 81, 1737-1747.	1.2	7
53	Perceived helpfulness of treatment for specific phobia: Findings from the World Mental Health Surveys. Journal of Affective Disorders, 2021, 288, 199-209.	2.0	9
54	Enhancing the Human Health Status Prediction: The ATHLOS Project. Applied Artificial Intelligence, 2021, 35, 834-856.	2.0	0

#	Article	IF	CITATIONS
55	Risk of all-cause mortality associated with chronic obstructive pulmonary disease and the role of healthy ageing trajectories: a population-based study of middle-aged and older adults. BMJ Open, 2021, 11, e050947.	0.8	0
56	To be happy and behave in a healthier way. A longitudinal study about gender differences in the older population. Psychology and Health, 2021 , , $1-17$.	1.2	4
57	The role of social support, detachment, and depressive and anxiety symptoms in suicidal thoughts and behaviours during the Covid-19 lockdown: Potential pathways. Current Psychology, 2021, , 1-12.	1.7	4
58	Perceived helpfulness of treatment for generalized anxiety disorder: a World Mental Health Surveys report. BMC Psychiatry, 2021, 21, 392.	1.1	10
59	Sarcopenia and Mild Cognitive Impairment in Older Adults from Six Low- and Middle-Income Countries. Journal of Alzheimer's Disease, 2021, 82, 1745-1754.	1.2	8
60	The association of detachment with affective disorder symptoms during the COVID-19 lockdown: The role of living situation and social support. Journal of Affective Disorders, 2021, 292, 464-470.	2.0	13
61	Multimorbidity and obesity in older adults from six low- and middle-income countries. Preventive Medicine, 2021, 153, 106816.	1.6	19
62	Kynurenine pathway in post-mortem prefrontal cortex and cerebellum in schizophrenia: relationship with monoamines and symptomatology. Journal of Neuroinflammation, 2021, 18, 198.	3.1	8
63	Thirtyâ€day suicidal thoughts and behaviors among hospital workers during the first wave of the Spain COVIDâ€19 outbreak. Depression and Anxiety, 2021, 38, 528-544.	2.0	70
64	Perceived helpfulness of treatment for alcohol use disorders: Findings from the World Mental Health Surveys. Drug and Alcohol Dependence, 2021, 229, 109158.	1.6	3
65	Previous disorders and depression outcomes in individuals with 12-month major depressive disorder in the World Mental Health surveys. Epidemiology and Psychiatric Sciences, 2021, 30, e70.	1.8	3
66	Psychotic experiences and subjective cognitive complaints among 224 842 people in 48 low- and middle-income countries. Epidemiology and Psychiatric Sciences, 2020, 29, e11.	1.8	22
67	Relationship between subjective well-being and healthy lifestyle behaviours in older adults: a longitudinal study. Aging and Mental Health, 2020, 24, 611-619.	1.5	36
68	Development of a Combined Sensory-Cognitive Measure Based on the Common Cause Hypothesis: Heterogeneous Trajectories and Associated Risk Factors. Gerontologist, The, 2020, 60, e357-e366.	2.3	6
69	Differential impact of transient and chronic loneliness on health status. A longitudinal study. Psychology and Health, 2020, 35, 177-195.	1.2	45
70	Factors associated with long-term impact on informal caregivers during Alzheimer's disease dementia progression: 36-month results from GERAS. International Psychogeriatrics, 2020, 32, 267-277.	0.6	40
71	Cross-national prevalence and factors associated with suicide ideation and attempts in older and young-and-middle age people. Aging and Mental Health, 2020, 24, 1533-1542.	1.5	31
72	The global, regional, and national burden of stomach cancer in 195 countries, 1990–2017: a systematic analysis for the Global Burden of Disease study 2017. The Lancet Gastroenterology and Hepatology, 2020, 5, 42-54.	3.7	390

#	Article	IF	CITATIONS
73	Association between intelligence quotient and disability: The role of socioeconomic status. Annals of Physical and Rehabilitation Medicine, 2020, 63, 296-301.	1.1	6
74	Association of bullying victimization with overweight and obesity among adolescents from 41 low― and middleâ€income countries. Pediatric Obesity, 2020, 15, e12571.	1.4	29
75	The relationship of childhood sexual and physical abuse with adulthood disability. Annals of Physical and Rehabilitation Medicine, 2020, 63, 332-339.	1.1	5
76	Loneliness and depression among older European adults: The role of perceived neighborhood built environment. Health and Place, 2020, 62, 102280.	1.5	32
77	Skeletal muscle mass in relation to 10 year cardiovascular disease incidence among middle aged and older adults: the ATTICA study. Journal of Epidemiology and Community Health, 2020, 74, 26-31.	2.0	68
78	The relationship between all-cause mortality and depression in different gender and age groups of the Spanish population. Journal of Affective Disorders, 2020, 266, 424-428.	2.0	6
79	Transdiagnostic development of internalizing psychopathology throughout the life course up to age 45: a World Mental Health Surveys report. Psychological Medicine, 2020, , 1-10.	2.7	1
80	Comorbidity within mental disorders: a comprehensive analysis based on 145 990 survey respondents from 27 countries. Epidemiology and Psychiatric Sciences, 2020, 29, e153.	1.8	67
81	Multimorbidity patterns in low-middle and high income regions: a multiregion latent class analysis using ATHLOS harmonised cohorts. BMJ Open, 2020, 10, e034441.	0.8	25
82	Cognitive function associated with different diagnoses of anxiety disorders over the lifespan: Results from a Spanish representative sample. Journal of Anxiety Disorders, 2020, 75, 102296.	1.5	6
83	Predictors of pain in general ageing populations: results from a multi-country analysis based on ATHLOS harmonized database. Journal of Headache and Pain, 2020, 21, 45.	2.5	9
84	Crossâ€cultural comparison of symptom networks in lateâ€life major depressive disorder: Yoruba Africans and the Spanish Population. International Journal of Geriatric Psychiatry, 2020, 35, 1060-1068.	1.3	7
85	Pain rates in general population for the period 1991–2015 and 10-years prediction: results from a multi-continent age-period-cohort analysis. Journal of Headache and Pain, 2020, 21, 52.	2.5	17
86	Functioning profiles in a nationally representative cohort of Spanish older adults: A latent class study. Health and Social Care in the Community, 2020, 28, 2190-2198.	0.7	3
87	Fruit and Vegetable Consumption and Sarcopenia among Older Adults in Low- and Middle-Income Countries. Nutrients, 2020, 12, 706.	1.7	19
88	Exploring the effect of loneliness on all-cause mortality: Are there differences between older adults and younger and middle-aged adults?. Social Science and Medicine, 2020, 258, 113087.	1.8	19
89	The new Horizon Europe programme 2021–2028: Should the gap between the burden of mental disorders and the funding of mental health research be filled?. European Journal of Psychiatry, 2020, 34, 44-46.	0.7	3
90	Informal Caregiving, Chronic Physical Conditions, and Physical Multimorbidity in 48 Low- and Middle-Income Countries. Journals of Gerontology - Series A Biological Sciences and Medical Sciences, 2020, 75, 1572-1578.	1.7	20

#	Article	IF	CITATIONS
91	Informal caregiving and physical activity among 204,315 adults in 38 low- and middle-income countries: A cross-sectional study. Preventive Medicine, 2020, 132, 106007.	1.6	12
92	Falls in older aged adults in 22 European countries: incidence, mortality and burden of disease from 1990 to 2017. Injury Prevention, 2020, 26, i67-i74.	1.2	65
93	Trajectories of Immediate and Delayed Verbal Memory in the Spanish General Population of Middle-aged and Older Adults. Brain Sciences, 2020, 10, 249.	1.1	3
94	Burden of injury along the development spectrum: associations between the Socio-demographic Index and disability-adjusted life year estimates from the Global Burden of Disease Study 2017. Injury Prevention, 2020, 26, i12-i26.	1.2	44
95	Are retired people higher in experiential wellbeing than working older adults? A time use approach Emotion, 2020, 20, 1411-1422.	1.5	4
96	Using Smartphones and Wearable Devices to Monitor Behavioral Changes During COVID-19. Journal of Medical Internet Research, 2020, 22, e19992.	2.1	155
97	Association between intelligence quotient and violence perpetration in the English general population. Psychological Medicine, 2019, 49, 1316-1323.	2.7	11
98	Health-related quality of life associated with different symptoms in women and in men who suffer from schizophrenia. Archives of Women's Mental Health, 2019, 22, 357-365.	1.2	5
99	Fruit and Vegetable Consumption and Potential Moderators Associated with All-Cause Mortality in a Representative Sample of Spanish Older Adults. Nutrients, 2019, 11, 1794.	1.7	13
100	Are loneliness and social isolation associated with cognitive decline?. International Journal of Geriatric Psychiatry, 2019, 34, 1613-1622.	1.3	202
101	Cohort Profile: The Ageing Trajectories of Health – Longitudinal Opportunities and Synergies (ATHLOS) project. International Journal of Epidemiology, 2019, 48, 1052-1053i.	0.9	39
102	The epidemiology of drug use disorders cross-nationally: Findings from the WHO's World Mental Health Surveys. International Journal of Drug Policy, 2019, 71, 103-112.	1.6	45
103	Mapping 123 million neonatal, infant and child deaths between 2000 and 2017. Nature, 2019, 574, 353-358.	13.7	161
104	<p>Patient-reported depression severity and cognitive symptoms as determinants of functioning in patients with major depressive disorder: a secondary analysis of the 2-year prospective PERFORM study</p> . Neuropsychiatric Disease and Treatment, 2019, Volume 15, 2313-2323.	1.0	19
105	Stability of clinically relevant depression symptoms in oldâ€age across 11 cohorts: a multiâ€state study. Acta Psychiatrica Scandinavica, 2019, 140, 541-551.	2.2	24
106	Physical multimorbidity and subjective cognitive complaints among adults in the United Kingdom: a cross-sectional community-based study. Scientific Reports, 2019, 9, 12417.	1.6	25
107	Global, Regional, and National Cancer Incidence, Mortality, Years of Life Lost, Years Lived With Disability, and Disability-Adjusted Life-Years for 29 Cancer Groups, 1990 to 2017. JAMA Oncology, 2019, 5, 1749.	3.4	1,691
108	Food insecurity (hunger) and suicide attempts among 179,771 adolescents attending school from 9 high-income, 31 middle-income, and 4 low-income countries: A cross-sectional study. Journal of Affective Disorders, 2019, 248, 91-98.	2.0	75

#	Article	IF	Citations
109	Childhood generalized specific phobia as an early marker of internalizing psychopathology across the lifespan: results from the World Mental Health Surveys. BMC Medicine, 2019, 17, 101.	2.3	24
110	Dietary Patterns and Their Association with Anxiety Symptoms among Older Adults: The ATTICA Study. Nutrients, 2019, 11, 1250.	1.7	28
111	Retrospective analysis of patients with advanced or metastatic gastric cancer in Colombia. Journal of Medical Economics, 2019, 22, 891-900.	1.0	2
112	Can the Positive and Negative Syndrome scale (PANSS) differentiate treatment-resistant from non-treatment-resistant schizophrenia? A factor analytic investigation based on data from the Pattern cohort study. Psychiatry Research, 2019, 276, 210-217.	1.7	11
113	Relationship between living alone and common mental disorders in the 1993, 2000 and 2007 National Psychiatric Morbidity Surveys. PLoS ONE, 2019, 14, e0215182.	1.1	60
114	Crossâ€national patterns of substance use disorder treatment and associations with mental disorder comorbidity in the WHO World Mental Health Surveys. Addiction, 2019, 114, 1446-1459.	1.7	27
115	CIBERSAM: Ten years of collaborative translational research in mental disorders. Revista De PsiquiatrÃa Y Salud Mental (English Edition), 2019, 12, 1-8.	0.2	5
116	Bullying Victimization and Suicide Attempt Among Adolescents Aged 12–15 Years From 48 Countries. Journal of the American Academy of Child and Adolescent Psychiatry, 2019, 58, 907-918.e4.	0.3	140
117	Association of Cohort and Individual Substance Use With Risk of Transitioning to Drug Use, Drug Use Disorder, and Remission From Disorder. JAMA Psychiatry, 2019, 76, 708.	6.0	27
118	Global, regional, and national burden of neurological disorders, 1990–2016: a systematic analysis for the Global Burden of Disease Study 2016. Lancet Neurology, The, 2019, 18, 459-480.	4.9	2,625
119	A comparison of <i>DSM</i> â€5 and <i>DSM</i> â€W agoraphobia in the World Mental Health Surveys. Depression and Anxiety, 2019, 36, 499-510.	2.0	22
120	A Cross-National Investigation of Hallucination-Like Experiences in 10 Countries: The E-CLECTIC Study. Schizophrenia Bulletin, 2019, 45, S43-S55.	2.3	15
121	The Effect of Exclusive Olive Oil Consumption on Successful Aging: A Combined Analysis of the ATTICA and MEDIS Epidemiological Studies. Foods, 2019, 8, 25.	1.9	23
122	Remote assessment of disease and relapse in major depressive disorder (RADAR-MDD): a multi-centre prospective cohort study protocol. BMC Psychiatry, 2019, 19, 72.	1.1	99
123	<p>Correlation of Health-Related Quality of Life in Clinically Stable Outpatients with Schizophrenia</p> . Neuropsychiatric Disease and Treatment, 2019, Volume 15, 3475-3486.	1.0	8
124	Determinants of time to institutionalisation and related healthcare and societal costs in a community-based cohort of patients with Alzheimer's disease dementia. European Journal of Health Economics, 2019, 20, 343-355.	1.4	26
125	Long-term trajectories of depressive symptoms in old age: Relationships with sociodemographic and health-related factors. Journal of Affective Disorders, 2019, 246, 329-337.	2.0	33
126	Anxiety, depression, loneliness and social network in the elderly: Longitudinal associations from The Irish Longitudinal Study on Ageing (TILDA). Journal of Affective Disorders, 2019, 246, 82-88.	2.0	252

#	Article	IF	CITATIONS
127	European mental health research resources: Picture and recommendations of the ROAMER project. European Neuropsychopharmacology, 2019, 29, 179-194.	0.3	9
128	Global, regional, and national burden of Alzheimer's disease and other dementias, 1990–2016: a systematic analysis for the Global Burden of Disease Study 2016. Lancet Neurology, The, 2019, 18, 88-106.	4.9	1,512
129	Global, regional, and national burden of traumatic brain injury and spinal cord injury, 1990–2016: a systematic analysis for the Global Burden of Disease Study 2016. Lancet Neurology, The, 2019, 18, 56-87.	4.9	1,064
130	Diez años de investigación traslacional colaborativa en enfermedades mentales: el CIBERSAM. Revista De PsiquiatrÃa Y Salud Mental, 2019, 12, 1-8.	1.0	68
131	Cognition and functionality in delusional disorder. European Psychiatry, 2019, 55, 52-60.	0.1	14
132	Mediterranean diet and wellbeing: evidence from a nationwide survey. Psychology and Health, 2019, 34, 321-335.	1.2	28
133	The association of religiosity with suicidal ideation and suicide attempts in the United Kingdom. Acta Psychiatrica Scandinavica, 2019, 139, 164-173.	2.2	24
134	Predictive capacity of prodromal symptoms in firstâ€episode psychosis of recent onset. Microbial Biotechnology, 2019, 13, 414-424.	0.9	12
135	The association of social support networks and loneliness with negative perceptions of ageing: evidence from the Irish Longitudinal Study on Ageing (TILDA). Ageing and Society, 2019, 39, 1070-1090.	1.2	20
136	Course of depression and cognitive decline at 3-year follow-up: The role of age of onset Psychology and Aging, 2019, 34, 475-485.	1.4	17
137	Longitudinal Relationships Between Positive Affect, Loneliness, and Suicide Ideation: Ageâ€Specific Factors in a General Population. Suicide and Life-Threatening Behavior, 2019, 49, 90-103.	0.9	33
138	Sociodemographic Indicators of Health Status Using a Machine Learning Approach and Data from the English Longitudinal Study of Aging (ELSA). Medical Science Monitor, 2019, 25, 1994-2001.	0.5	22
139	Barriers to and Facilitators of Engagement With mHealth Technology for Remote Measurement and Management of Depression: Qualitative Analysis. JMIR MHealth and UHealth, 2019, 7, e11325.	1.8	88
140	Retrospective analysis of patients with locally advanced or metastatic gastric cancer in Argentina. Medwave, 2019, 19, e7693-e7693.	0.2	1
141	Psychotic experiences and general medical conditions: a cross-national analysis based on 28 002 respondents from 16 countries in the WHO World Mental Health Surveys. Psychological Medicine, 2018, 48, 2730-2739.	2.7	27
142	Frailty and health status of older individuals in three European countries: The COURAGE cross-sectional study. Experimental Gerontology, 2018, 106, 137-144.	1.2	3
143	Post-traumatic stress symptoms are associated with physical multimorbidity: Findings from the Adult Psychiatric Morbidity Survey 2007. Journal of Affective Disorders, 2018, 232, 385-392.	2.0	33
144	Depression and metabolic syndrome in the older population: A review of evidence. Journal of Affective Disorders, 2018, 237, 56-64.	2.0	41

#	Article	IF	CITATIONS
145	Understanding the course of persistent symptoms in schizophrenia: Longitudinal findings from the pattern study. Psychiatry Research, 2018, 267, 56-62.	1.7	26
146	Risk factors and topographies for selfâ€injurious behaviour in a sample of adults with intellectual developmental disorders. Journal of Intellectual Disability Research, 2018, 62, 1018-1029.	1.2	17
147	Psychotic experiences and religiosity: data from the <scp>WHO</scp> World Mental Health Surveys. Acta Psychiatrica Scandinavica, 2018, 137, 306-315.	2.2	27
148	Treatment gap for anxiety disorders is global: Results of the World Mental Health Surveys in 21 countries. Depression and Anxiety, 2018, 35, 195-208.	2.0	318
149	Suicidal thoughts and behaviors among college students and same-aged peers: results from the World Health Organization World Mental Health Surveys. Social Psychiatry and Psychiatric Epidemiology, 2018, 53, 279-288.	1.6	79
150	Health-related quality of life in outpatients with schizophrenia: factors that determine changes over time. Social Psychiatry and Psychiatric Epidemiology, 2018, 53, 239-248.	1.6	8
151	Determinants of Health Trajectories in England and the United States: An Approach to Identify Different Patterns of Healthy Aging. Journals of Gerontology - Series A Biological Sciences and Medical Sciences, 2018, 73, 1512-1518.	1.7	36
152	The association between pain and suicidal behavior in an English national sample: The role of psychopathology. Journal of Psychiatric Research, 2018, 98, 39-46.	1.5	18
153	Complementary and alternative medicine contacts by persons with mental disorders in 25 countries: results from the World Mental Health Surveys. Epidemiology and Psychiatric Sciences, 2018, 27, 552-567.	1.8	47
154	The associations between psychotic experiences and substance use and substance use disorders: findings from the World Health Organization World Mental Health surveys. Addiction, 2018, 113, 924-934.	1.7	56
155	The impact of socioeconomic status on the association between biomedical and psychosocial well-being and all-cause mortality in older Spanish adults. Social Psychiatry and Psychiatric Epidemiology, 2018, 53, 259-268.	1.6	8
156	Post-traumatic stress disorder associated with sexual assault among women in the WHO World Mental Health Surveys. Psychological Medicine, 2018, 48, 155-167.	2.7	69
157	Sociodemographic Factors Associated With Changes in Successful Aging in Spain: A Follow-Up Study. Journal of Aging and Health, 2018, 30, 1244-1262.	0.9	11
158	Mobility Difficulties, Physical Activity, and All-cause Mortality Risk in a Nationally representative Sample of Older Adults. Journals of Gerontology - Series A Biological Sciences and Medical Sciences, 2018, 73, 1272-1279.	1.7	30
159	Recovery from DSM-IV post-traumatic stress disorder in the WHO World Mental Health surveys. Psychological Medicine, 2018, 48, 437-450.	2.7	28
160	Trial exclusion criteria and their impact on the estimation of antipsychotic drugs effect: A case study using the SOHO database. Schizophrenia Research, 2018, 193, 146-153.	1.1	8
161	Depression, sleep problems, and perceived stress among informal caregivers in 58 low-, middle-, and high-income countries: A cross-sectional analysis of community-based surveys. Journal of Psychiatric Research, 2018, 96, 115-123.	1.5	64
162	Impact of pain and remission in the functioning of patients with depression in <scp>M</scp> ainland <scp>C</scp> hina, <scp>T</scp> aiwan, and <scp>H</scp> ong <scp>K</scp> ong. Asia-Pacific Psychiatry, 2018, 10, e12295.	1.2	3

#	Article	IF	Citations
163	Previous Mental Disorders and Subsequent Onset of Chronic Back or Neck Pain: Findings From 19 Countries. Journal of Pain, 2018, 19, 99-110.	0.7	32
164	Anti-Inflammatory Nutrition and Successful Ageing in Elderly Individuals: The Multinational MEDIS Study. Gerontology, 2018, 64, 3-10.	1.4	18
165	The associations of earlier trauma exposures and history of mental disorders with PTSD after subsequent traumas. Molecular Psychiatry, 2018, 23, 1892-1899.	4.1	52
166	Psychometric Properties of Spanish Adaptation of the PDD-MRS Scale in Adults with Intellectual Developmental Disorders: The EVTEA-DI Scale. Journal of Autism and Developmental Disorders, 2018, 48, 1566-1578.	1.7	5
167	Socio-economic variations in the mental health treatment gap for people with anxiety, mood, and substance use disorders: results from the WHO World Mental Health (WMH) surveys. Psychological Medicine, 2018, 48, 1560-1571.	2.7	274
168	Measurement invariance of the WHOQOL-AGE questionnaire across three European countries. Quality of Life Research, 2018, 27, 1015-1025.	1.5	13
169	Costs and Burden Associated With Loss of Labor Productivity in Informal Caregivers of People With Dementia. Journal of Occupational and Environmental Medicine, 2018, 60, 449-456.	0.9	15
170	Machine learning methodologies versus cardiovascular risk scores, in predicting disease risk. BMC Medical Research Methodology, 2018, 18, 179.	1.4	67
171	Social determinants, health status and 10-year mortality among 10,906 older adults from the English longitudinal study of aging: the ATHLOS project. BMC Public Health, 2018, 18, 1357.	1.2	24
172	Act-Belong-Commit Indicators Promote Mental Health and Wellbeing among Irish Older Adults. American Journal of Health Behavior, 2018, 42, 31-45.	0.6	22
173	The global burden of disease attributable to alcohol and drug use in 195 countries and territories, 1990–2016: a systematic analysis for the Global Burden of Disease Study 2016. Lancet Psychiatry,the, 2018, 5, 987-1012.	3.7	885
174	Clinical outcomes of patients with major depressive disorder treated with either duloxetine, escitalopram, fluoxetine, paroxetine, or sertraline. Neuropsychiatric Disease and Treatment, 2018, Volume 14, 2473-2484.	1.0	7
175	Measurement invariance of the Spanish Launay–Slade Hallucinations Scaleâ€Extended version between putatively healthy controls and people diagnosed with a mental disorder. International Journal of Methods in Psychiatric Research, 2018, 27, e1741.	1.1	12
176	Relationship between attention-deficit hyperactivity disorder symptoms and problem gambling: A mediation analysis of influential factors among 7,403 individuals from the UK. Journal of Behavioral Addictions, 2018, 7, 781-791.	1.9	22
177	Functional impairment in patients with major depressive disorder: the 2-year PERFORM study. Neuropsychiatric Disease and Treatment, 2018, Volume 14, 239-249.	1.0	72
178	The burden of disease in Spain: Results from the Global Burden of Disease 2016. Medicina ClÃnica (English Edition), 2018, 151, 171-190.	0.1	37
179	Factors Related to Unemployment in Europe. A Cross-Sectional Study from the COURAGE Survey in Finland, Poland and Spain. International Journal of Environmental Research and Public Health, 2018, 15, 722.	1.2	14
180	Attention deficit hyperactivity disorder symptoms and disordered eating in the English general population. International Journal of Eating Disorders, 2018, 51, 942-952.	2.1	8

#	Article	IF	CITATIONS
181	Long-term follow-up on health-related quality of life in major depressive disorder: a 2-year European cohort study. Neuropsychiatric Disease and Treatment, 2018, Volume 14, 1339-1350.	1.0	62
182	Mortality in unipolar depression preceding and following chronic somatic diseases. Acta Psychiatrica Scandinavica, 2018, 138, 500-508.	2.2	22
183	The association between psychotic experiences and health-related quality of life: a cross-national analysis based on World Mental Health Surveys. Schizophrenia Research, 2018, 201, 46-53.	1.1	20
184	The association between problem gambling and psychotic experiences: Findings from the Adult Psychiatric Morbidity Survey 2007. Schizophrenia Research, 2018, 201, 79-84.	1.1	6
185	Mediterranean diet and depression among older individuals: The multinational MEDIS study. Experimental Gerontology, 2018, 110, 67-72.	1.2	48
186	Sibship size, birth order and psychotic experiences: Evidence from 43 low- and middle-income countries. Schizophrenia Research, 2018, 201, 406-412.	1.1	1
187	The "RCT augmentation― a novel simulation method to add patient heterogeneity into phase III trials. BMC Medical Research Methodology, 2018, 18, 75.	1.4	2
188	Furthering the Evidence of the Effectiveness of Employment Strategies for People with Mental Disorders in Europe: A Systematic Review. International Journal of Environmental Research and Public Health, 2018, 15, 838.	1.2	9
189	Determinants of mobility in populations of older adults: Results from a cross-sectional study in Finland, Poland and Spain. Maturitas, 2018, 115, 84-91.	1.0	10
190	La carga de enfermedad en España: resultados del Estudio de la Carga Global de las Enfermedades 2016. Medicina ClÃnica, 2018, 151, 171-190.	0.3	113
191	Characteristics of patients with depression initiating or switching antidepressant treatment: baseline analyses of the PERFORM cohort study. BMC Psychiatry, 2018, 18, 80.	1.1	6
192	Epidemiology of panic attacks, panic disorder and the moderating role of age: Results from a population-based study. Journal of Affective Disorders, 2018, 241, 627-633.	2.0	17
193	Alcohol use and burden for 195 countries and territories, 1990 \hat{a} \in "2016: a systematic analysis for the Global Burden of Disease Study 2016. Lancet, The, 2018, 392, 1015-1035.	6.3	2,005
194	Global Mortality From Firearms, 1990-2016. JAMA - Journal of the American Medical Association, 2018, 320, 792.	3.8	189
195	The association between socioeconomic status and depression among older adults in Finland, Poland and Spain: A comparative cross-sectional study of distinct measures and pathways. Journal of Affective Disorders, 2018, 241, 311-318.	2.0	73
196	The burden of disease in Greece, health loss, risk factors, and health financing, 2000–16: an analysis of the Global Burden of Disease Study 2016. Lancet Public Health, The, 2018, 3, e395-e406.	4.7	37
197	Is Parental Longevity Associated With the Cardiovascular Risk and the Successful Aging of Their Offspring? Results From the Multinational MEDIS Study. Angiology, 2017, 68, 124-131.	0.8	2
198	The association between childhood adversities and subsequent first onset of psychotic experiences: a cross-national analysis of 23 998 respondents from 17 countries. Psychological Medicine, 2017, 47, 1230-1245.	2.7	108

#	Article	IF	Citations
199	Eficacia, eficiencia y efectividad en el tratamiento multidimensional de la esquizofrenia: proyecto Rethinking. Revista De PsiquiatrÃa Y Salud Mental, 2017, 10, 4-20.	1.0	18
200	Association of <i>DSM-IV</i> Posttraumatic Stress Disorder With Traumatic Experience Type and History in the World Health Organization World Mental Health Surveys. JAMA Psychiatry, 2017, 74, 270.	6.0	179
201	Loneliness and depression in the elderly: the role of social network. Social Psychiatry and Psychiatric Epidemiology, 2017, 52, 381-390.	1.6	274
202	Real-world Effectiveness of Antipsychotics for the Treatment of Negative Symptoms in Patients with Schizophrenia with Predominantly Negative Symptoms. Pharmacopsychiatry, 2017, 50, 56-63.	1.7	15
203	Trajectories of Verbal Episodic Memory in Middleâ€Aged and Older Adults: Evidence from the English Longitudinal Study of Ageing. Journal of the American Geriatrics Society, 2017, 65, 1274-1281.	1.3	51
204	Advanced analytical methodologies for measuring healthy ageing and its determinants, using factor analysis and machine learning techniques: the ATHLOS project. Scientific Reports, 2017, 7, 43955.	1.6	77
205	The Impact of Depression on the Development of Mild Cognitive Impairment over 3 Years of Follow-Up: A Population-Based Study. Dementia and Geriatric Cognitive Disorders, 2017, 43, 155-169.	0.7	29
206	Gender-related differences in the multi-pathway effect of social determinants on quality of life in older ageâ€"the COURAGE in Europe project. Quality of Life Research, 2017, 26, 1865-1878.	1.5	34
207	The protective properties of Act-Belong-Commit indicators against incident depression, anxiety, and cognitive impairment among older Irish adults: Findings from a prospective community-based study. Experimental Gerontology, 2017, 91, 79-87.	1.2	35
208	n-3 Fatty acids, Mediterranean diet and cognitive function in normal aging: A systematic review. Experimental Gerontology, 2017, 91, 39-50.	1.2	49
209	Dietary inflammatory potential is linked to cardiovascular disease risk burden in the US adult population. International Journal of Cardiology, 2017, 240, 409-413.	0.8	34
210	Healthcare Access and Quality Index based on mortality from causes amenable to personal health care in 195 countries and territories, 1990–2015: a novel analysis from the Global Burden of Disease Study 2015. Lancet, The, 2017, 390, 231-266.	6.3	480
211	The association between psychotic experiences and disability: results from the <scp>WHO</scp> World Mental Health Surveys. Acta Psychiatrica Scandinavica, 2017, 136, 74-84.	2.2	58
212	A novel approach to measuring response and remission in schizophrenia in clinical trials. Schizophrenia Research, 2017, 190, 123-128.	1.1	6
213	Posttraumatic stress disorder in the World Mental Health Surveys. Psychological Medicine, 2017, 47, 2260-2274.	2.7	675
214	Cross-sectional Comparison of the Epidemiology of <i>DSM-5</i> Generalized Anxiety Disorder Across the Globe. JAMA Psychiatry, 2017, 74, 465.	6.0	274
215	What Drives Country Differences in Cost of Alzheimer's Disease? An Explanation from Resource Use in the GERAS Study. Journal of Alzheimer's Disease, 2017, 57, 797-812.	1.2	23
216	Child death and maternal psychosis-like experiences in 44 low- and middle-income countries: The role of depression. Schizophrenia Research, 2017, 183, 41-46.	1.1	6

#	Article	IF	Citations
217	Undertreatment of people with major depressive disorder in 21 countries. British Journal of Psychiatry, 2017, 210, 119-124.	1.7	664
218	Posttraumatic stress disorder associated with unexpected death of a loved one: Cross-national findings from the world mental health surveys. Depression and Anxiety, 2017, 34, 315-326.	2.0	33
219	Post-traumatic stress disorder associated with natural and human-made disasters in the World Mental Health Surveys. Psychological Medicine, 2017, 47, 227-241.	2.7	110
220	Epidemiology of depression with psychotic experiences and its association with chronic physical conditions in 47 low- and middle-income countries. Psychological Medicine, 2017, 47, 531-542.	2.7	32
221	Exploring the path between depression, anxiety and 10-year cardiovascular disease incidence, among apparently healthy Greek middle-aged adults: The ATTICA study. Maturitas, 2017, 106, 73-79.	1.0	19
222	Trauma and psychotic experiences: transnational data from the World Mental Health Survey. British Journal of Psychiatry, 2017, 211, 373-380.	1.7	82
223	All-cause mortality and multimorbidity in older adults: The role of social support and loneliness. Experimental Gerontology, 2017, 99, 120-126.	1.2	67
224	Estimating treatment coverage for people with substance use disorders: an analysis of data from the World Mental Health Surveys. World Psychiatry, 2017, 16, 299-307.	4.8	160
225	Association Between Psychotic Experiences and Subsequent Suicidal Thoughts and Behaviors. JAMA Psychiatry, 2017, 74, 1136.	6.0	7 5
226	Global, regional, and national incidence, prevalence, and years lived with disability for 328 diseases and injuries for 195 countries, 1990–2016: a systematic analysis for the Global Burden of Disease Study 2016. Lancet, The, 2017, 390, 1211-1259.	6.3	5,578
227	Development and validation of the Questionnaire of Stressful Life Events (QSLE). Journal of Psychiatric Research, 2017, 95, 213-223.	1.5	9
228	Disease Progression in Mild Dementia due to Alzheimer Disease in an 18-Month Observational Study (GERAS): The Impact on Costs and Caregiver Outcomes. Dementia and Geriatric Cognitive Disorders Extra, 2017, 7, 87-100.	0.6	36
229	National funding for mental health research in Finland, France, Spain and the United Kingdom. European Neuropsychopharmacology, 2017, 27, 892-899.	0.3	13
230	Trauma and PTSD in the WHO World Mental Health Surveys. Högre Utbildning, 2017, 8, 1353383.	1.4	801
231	The role of socio-economic status and neighborhood social capital on loneliness among older adults: evidence from the Sant Boi Aging Study. Social Psychiatry and Psychiatric Epidemiology, 2017, 52, 1237-1246.	1.6	29
232	How useful is the EQ-5D in assessing the impact of caring for people with Alzheimer's disease?. Health and Quality of Life Outcomes, 2017, 15, 16.	1.0	27
233	Cognitive reserve is associated with quality of life: A population-based study. Experimental Gerontology, 2017, 87, 67-73.	1.2	31
234	Health conditions and role limitation in three European Regions: a public-health perspective. Gaceta Sanitaria, 2017, 31, 2-10.	0.6	6

#	Article	IF	Citations
235	Brain alterations in low-frequency fluctuations across multiple bands in obsessive compulsive disorder. Brain Imaging and Behavior, 2017, 11, 1690-1706.	1.1	25
236	Factors associated with and impact of pain persistence in Asian patients with depression: a 3-month, prospective observational study. International Journal of Psychiatry in Clinical Practice, 2017, 21, 29-35.	1.2	3
237	The descriptive epidemiology of DSM-IV Adult ADHD in the World Health Organization World Mental Health Surveys. ADHD Attention Deficit and Hyperactivity Disorders, 2017, 9, 47-65.	1.7	507
238	Brain structure in women at risk of postpartum psychosis: an MRI study. Translational Psychiatry, 2017, 7, 1286.	2.4	26
239	Factors associated with failure to achieve remission and with relapse after remission in patients with major depressive disorder in the PERFORM study. Neuropsychiatric Disease and Treatment, 2017, Volume 13, 2151-2165.	1.0	44
240	The Burden of Mental Disorders in the Eastern Mediterranean Region, 1990-2013. PLoS ONE, 2017, 12, e0169575.	1.1	102
241	Latent class analysis of multimorbidity patterns and associated outcomes in Spanish older adults: a prospective cohort study. BMC Geriatrics, 2017, 17, 186.	1.1	82
242	The cross-national epidemiology of social anxiety disorder: Data from the World Mental Health Survey Initiative. BMC Medicine, 2017, 15, 143.	2.3	258
243	Depression comorbid with tuberculosis and its impact on health status: cross-sectional analysis of community-based data from 48 low- and middle-income countries. BMC Medicine, 2017, 15, 209.	2.3	7 9
244	Association of experienced and evaluative well-being with health in nine countries with different income levels: a cross-sectional study. Globalization and Health, 2017, 13, 65.	2.4	20
245	Comparison of clinical outcomes with orodispersible versus standard oral olanzapine tablets in nonadherent patients with schizophrenia or bipolar disorder. Patient Preference and Adherence, 2017, Volume 11, 1019-1025.	0.8	11
246	Recovery in patients with major depressive disorder (MDD): results of a 6-month, multinational, observational study. Patient Preference and Adherence, 2017, Volume 11, 1859-1868.	0.8	20
247	The use of random-effects models to identify health care center-related characteristics modifying the effect of antipsychotic drugs. Clinical Epidemiology, 2017, Volume 9, 689-698.	1.5	0
248	Functional Impairment and Painful Physical Symptoms in Patients with Major Depressive Disorder Treated with Antidepressants: Real-World Evidence from the Middle East. Clinical Practice and Epidemiology in Mental Health, 2017, 13, 145-155.	0.6	2
249	Determinants of Quality of Life in Ageing Populations: Results from a Cross-Sectional Study in Finland, Poland and Spain. PLoS ONE, 2016, 11, e0159293.	1.1	64
250	Sex differences in the course of schizophrenia across diverse regions of the world. Neuropsychiatric Disease and Treatment, 2016, Volume 12, 2927-2939.	1.0	30
251	Impact of anxiety symptoms on outcomes of depression: an observational study in Asian patients. Neuropsychiatric Disease and Treatment, 2016, 12, 795.	1.0	2
252	Functioning in patients with major depression treated with duloxetine or a selective serotonin reuptake inhibitor in East Asia. Neuropsychiatric Disease and Treatment, 2016, 12, 383.	1.0	3

#	Article	IF	CITATIONS
253	Clinical features, comorbidity, and cognitive impairment in elderly bipolar patients. Neuropsychiatric Disease and Treatment, 2016, 12, 1203.	1.0	33
254	Positive Affect Is Inversely Associated with Mortality in Individuals without Depression. Frontiers in Psychology, 2016, 7, 1040.	1.1	13
255	Which Are the Most Burdensome Functioning Areas in Depression? A Cross-National Study. Frontiers in Psychology, 2016, 7, 1342.	1.1	16
256	Psychotic symptoms and smoking in 44 countries. Acta Psychiatrica Scandinavica, 2016, 133, 497-505.	2.2	32
257	Mild cognitive impairment in a Spanish representative sample: prevalence and associated factors. International Journal of Geriatric Psychiatry, 2016, 31, 858-867.	1.3	63
258	Social relationships, loneliness, and mental health among older men and women in Ireland: A prospective community-based study. Journal of Affective Disorders, 2016, 204, 59-69.	2.0	117
259	Post-traumatic stress disorder associated with life-threatening motor vehicle collisions in the WHO World Mental Health Surveys. BMC Psychiatry, 2016, 16, 257.	1.1	34
260	Weight Perception, Satisfaction, Control, and Low Energy Dietary Reporting in the US Adult Population: Results from the National Health and Nutrition Examination Survey 2007-2012. Journal of the Academy of Nutrition and Dietetics, 2016, 116, 579-589.	0.4	15
261	Identifying factors of activities of daily living important for cost and caregiver outcomes in Alzheimer's disease. International Psychogeriatrics, 2016, 28, 247-259.	0.6	35
262	Negative perceptions of ageing predict the onset and persistence of depression and anxiety: Findings from a prospective analysis of the Irish Longitudinal Study on Ageing (TILDA). Journal of Affective Disorders, 2016, 199, 132-138.	2.0	88
263	Subclinical psychosis and pain in an English national sample: The role of common mental disorders. Schizophrenia Research, 2016, 175, 209-215.	1.1	12
264	Study of positive and negative feedback sensitivity in psychosis using the Wisconsin Card Sorting Test. Comprehensive Psychiatry, 2016, 68, 119-128.	1.5	16
265	Psychotic-like experiences and disordered eating in the English general population. Psychiatry Research, 2016, 241, 26-34.	1.7	16
266	Spanish validation of the social stigma scale: Community Attitudes towards Mental Illness. Revista De PsiquiatrÃa Y Salud Mental (English Edition), 2016, 9, 150-157.	0.2	15
267	Factors associated with skeletal muscle mass, sarcopenia, and sarcopenic obesity in older adults: a multiâ€continent study. Journal of Cachexia, Sarcopenia and Muscle, 2016, 7, 312-321.	2.9	182
268	Incidence of stressful life events and influence of sociodemographic and clinical variables on the onset of first-episode psychosis. Psychiatry Research, 2016, 245, 108-115.	1.7	18
269	Mental disorders among college students in the World Health Organization World Mental Health Surveys. Psychological Medicine, 2016, 46, 2955-2970.	2.7	866
270	Estimates of global, regional, and national incidence, prevalence, and mortality of HIV, 1980–2015: the Global Burden of Disease Study 2015. Lancet HIV,the, 2016, 3, e361-e387.	2.1	461

#	Article	IF	Citations
271	Population prevalence of edentulism and its association with depression and self-rated health. Scientific Reports, 2016, 6, 37083.	1.6	89
272	The role of socio-economic status in depression: results from the COURAGE (aging survey in Europe). BMC Public Health, 2016, 16, 1098.	1.2	278
273	Cross-national epidemiology of panic disorder and panic attacks in the world mental health surveys. Depression and Anxiety, 2016, 33, 1155-1177.	2.0	144
274	Risk and functional significance of psychotic experiences among individuals with depression in 44 low- and middle-income countries. Psychological Medicine, 2016, 46, 2655-2665.	2.7	45
275	How to deal with missing longitudinal data in cost of illness analysis in Alzheimer's disease—suggestions from the GERAS observational study. BMC Medical Research Methodology, 2016, 16, 83.	1.4	13
276	Exploring the excess mortality due to depressive symptoms in a community-based sample: The role of Alzheimer's Disease. Journal of Affective Disorders, 2016, 202, 163-170.	2.0	22
277	Specificity proteins 1 and 4, hippocampal volume and first-episode psychosis. British Journal of Psychiatry, 2016, 208, 591-592.	1.7	2
278	Altered CSNK1E, FABP4 and NEFH protein levels in the dorsolateral prefrontal cortex in schizophrenia. Schizophrenia Research, 2016, 177, 88-97.	1.1	26
279	Direct and indirect costs and resource use in dementia care: A cross-sectional study in patients living at home. International Journal of Nursing Studies, 2016, 55, 39-49.	2.5	26
280	The Bidirectional Associations Between Psychotic Experiences and DSM-IV Mental Disorders. American Journal of Psychiatry, 2016, 173, 997-1006.	4.0	176
281	Anti-NMDAR antibodies in new-onset psychosis. Positive results in an HIV-infected patient. Brain, Behavior, and Immunity, 2016, 56, 56-60.	2.0	22
282	Associations between DSM-IV mental disorders and subsequent onset of arthritis. Journal of Psychosomatic Research, 2016, 82, 11-16.	1.2	6
283	The joint association of depression and cognitive function with severe disability among community-dwelling older adults in Finland, Poland and Spain. Experimental Gerontology, 2016, 76, 39-45.	1.2	9
284	Mild cognitive impairment is associated with falls among older adults: Findings from the Irish Longitudinal Study on Ageing (TILDA). Experimental Gerontology, 2016, 75, 42-47.	1.2	46
285	Validación al castellano de la escala de estigma social: Community Attitudes towards Mental Illness en población adolescente. Revista De PsiquiatrÃa Y Salud Mental, 2016, 9, 150-157.	1.0	24
286	Global Multimorbidity Patterns: A Cross-Sectional, Population-Based, Multi-Country Study. Journals of Gerontology - Series A Biological Sciences and Medical Sciences, 2016, 71, 205-214.	1.7	404
287	Measurement Invariance of the Day Reconstruction Method: Results from the COURAGE in Europe Project. Journal of Happiness Studies, 2016, 17, 1769-1787.	1.9	8
288	Realâ€world outcomes in patients with depression treated with duloxetine or a selective serotonin reuptake inhibitor in <scp>E</scp> ast <scp>A</scp> sia. Asia-Pacific Psychiatry, 2016, 8, 51-59.	1.2	12

#	Article	IF	Citations
289	Loneliness, Social Networks, and Health: A Cross-Sectional Study in Three Countries. PLoS ONE, 2016, 11, e0145264.	1.1	121
290	Personalidad internalizante y externalizante y efectos subjetivos en una muestra de adolescentes consumidores de cánnabis. Revista De Psicologia De La Salud, 2016, 28, 231.	0.2	1
291	Degree of cognitive impairment and mortality: a 17-year follow-up in a community study. Epidemiology and Psychiatric Sciences, 2015, 24, 503-511.	1.8	16
292	Comparing Resource Use In Alzheimer's Disease Across Three European Countries - 18-Month Results of The Geras Study. Value in Health, 2015, 18, A759-A760.	0.1	4
293	How Useful are Eq-5d And Zbi In Assessing The Impact of Caring for Alzheimer's Disease Patients?. Value in Health, 2015, 18, A760.	0.1	1
294	Visual encoding impairment in patients with schizophrenia: Contribution of reduced working memory span, decreased processing speed, and affective symptoms Neuropsychology, 2015, 29, 17-24.	1.0	15
295	Association of auditory-verbal and visual hallucinations with impaired and improved recognition of colored pictures Neuropsychology, 2015, 29, 667-674.	1.0	8
296	Which somatic symptoms are associated with an unfavorable course in <scp>C</scp> hinese patients with major depressive disorder?. Asia-Pacific Psychiatry, 2015, 7, 427-435.	1.2	17
297	Mild cognitive impairment diagnosed with the new <scp>DSM</scp> â€5 criteria: prevalence and associations with nonâ€cognitive psychopathology. Acta Psychiatrica Scandinavica, 2015, 131, 29-39.	2.2	48
298	Antidepressant medication treatment patterns in Asian patients with major depressive disorder. Patient Preference and Adherence, 2015, 9, 421.	0.8	13
299	Predictors of remission in the treatment of major depressive disorder: real-world evidence from a 6-month prospective observational study. Neuropsychiatric Disease and Treatment, 2015, 11, 197.	1.0	17
300	Health-related quality of life in patients with depression treated with duloxetine or a selective serotonin reuptake inhibitor in a naturalistic outpatient setting. Patient Preference and Adherence, 2015, 9, 1481.	0.8	2
301	Recreational drug use among individuals living with HIV in Europe: review of the prevalence, comparison with the general population and HIV guidelines recommendations. Frontiers in Microbiology, 2015, 6, 690.	1.5	29
302	Phosphorylation of Transcription Factor Specificity Protein 4 ls Increased in Peripheral Blood Mononuclear Cells of First-Episode Psychosis. PLoS ONE, 2015, 10, e0125115.	1.1	4
303	Impact of painful physical symptoms on depression outcomes in elderly Asian patients. International Psychogeriatrics, 2015, 27, 305-312.	0.6	7
304	Research priorities for public mental health in Europe: recommendations of the ROAMER project. European Journal of Public Health, 2015, 25, 249-254.	0.1	113
305	Country-level and individual correlates of overweight and obesity among primary school children: a cross-sectional study in seven European countries. BMC Public Health, 2015, 15, 475.	1.2	64
306	Global, regional, and national incidence, prevalence, and years lived with disability for 301 acute and chronic diseases and injuries in 188 countries, 1990–2013: a systematic analysis for the Global Burden of Disease Study 2013. Lancet, The, 2015, 386, 743-800.	6.3	4,951

#	Article	IF	Citations
307	Psychotic Experiences in the General Population. JAMA Psychiatry, 2015, 72, 697.	6.0	387
308	The state of the art on European well-being research within the area of mental health. International Journal of Clinical and Health Psychology, 2015, 15, 171-179.	2.7	19
309	A dimensional comparison between delusional disorder, schizophrenia and schizoaffective disorder. Schizophrenia Research, 2015, 169, 248-254.	1.1	27
310	The association between obesity and severe disability among adults aged 50 or over in nine high-income, middle-income and low-income countries: a cross-sectional study. BMJ Open, 2015, 5, e007313-e007313.	0.8	42
311	The Spanish Burden of Disease 2010: Neurological, mental and substance use disorders. Revista De PsiquiatrÃa Y Salud Mental (English Edition), 2015, 8, 207-217.	0.2	5
312	Desarrollo profesional en investigación traslacional en neurociencias y salud mental: educación y formación dentro del Centro de Investigación Biomédica en Red en Salud Mental. Revista De PsiquiatrÃa Y Salud Mental, 2015, 8, 65-74.	1.0	6
313	The association between social relationships and depression: A systematic review. Journal of Affective Disorders, 2015, 175, 53-65.	2.0	552
314	Cannabis and bipolar disorder: does quitting cannabis use during manic/mixed episode improve clinical/functional outcomes?. Acta Psychiatrica Scandinavica, 2015, 131, 100-110.	2.2	40
315	The association between obesity and back pain in nine countries: a cross-sectional study. BMC Public Health, 2015, 15, 123.	1.2	57
316	The role of muscle mass and body fat on disability among older adults: A cross-national analysis. Experimental Gerontology, 2015, 69, 27-35.	1.2	62
317	Suicidality and its relationship with depression, alcohol disorders and childhood experiences of violence: Results from the ESEMeD study. Journal of Affective Disorders, 2015, 175, 168-174.	2.0	30
318	Gender Differences in Psychotropic Use Across Europe: Results From a Large Cross-Sectional, Population-Based Study. European Psychiatry, 2015, 30, 778-788.	0.1	47
319	Associations between DSM-IV mental disorders and subsequent non-fatal, self-reported stroke. Journal of Psychosomatic Research, 2015, 79, 130-136.	1.2	17
320	The association of relationship quality and social networks with depression, anxiety, and suicidal ideation among older married adults: Findings from a cross-sectional analysis of the Irish Longitudinal Study on Ageing (TILDA). Journal of Affective Disorders, 2015, 179, 134-141.	2.0	74
321	Diabetes mellitus and its association with central obesity and disability among older adults: A global perspective. Experimental Gerontology, 2015, 64, 70-77.	1.2	56
322	Association between traumatic events and post-traumatic stress disorder: results from the ESEMeD-Spain study. Epidemiology and Psychiatric Sciences, 2015, 24, 172-183.	1.8	35
323	La carga de la enfermedad en España 2010: trastornos neurológicos, mentales y re: trastornos neurológicos, mentales y relacionados con el consumo de sustancias. Revista De PsiquiatrÃa Y Salud Mental, 2015, 8, 207-217.	1.0	17
324	Determinants of societal costs in Alzheimer's disease: GERASÂstudyÂbaseline results. Alzheimer's and Dementia, 2015, 11, 933-945.	0.4	62

#	Article	IF	Citations
325	Semaphorin and plexin gene expression is altered in the prefrontal cortex of schizophrenia patients with and without auditory hallucinations. Psychiatry Research, 2015, 229, 850-857.	1.7	31
326	Understanding the impact of persistent symptoms in schizophrenia: Cross-sectional findings from the Pattern study. Schizophrenia Research, 2015, 169, 234-240.	1.1	29
327	Mental health research priorities for Europe. Lancet Psychiatry, the, 2015, 2, 1036-1042.	3.7	158
328	The Role of Energy Balance in Successful Aging Among Elderly Individuals. Journal of Aging and Health, 2015, 27, 1375-1391.	0.9	11
329	Relationship of insight with medication adherence and the impact on outcomes in patients with schizophrenia and bipolar disorder: results from a 1-year European outpatient observational study. BMC Psychiatry, 2015, 15, 189.	1.1	82
330	Impact of Negative Symptoms on Quality of Life in Patients with Schizophrenia. Value in Health, 2015, 18, A836-A837.	0.1	8
331	Development and validation of the Self-Stigma Questionnaire (SSQ) for people with schizophrenia and its relation to social functioning. Comprehensive Psychiatry, 2015, 62, 93-99.	1.5	13
332	Chinese patients with major depression: Do concomitant pain symptoms affect quality of life independently of severity of depression?. International Journal of Psychiatry in Clinical Practice, 2015, 19, 174-181.	1.2	5
333	Global, regional, and national disability-adjusted life years (DALYs) for 306 diseases and injuries and healthy life expectancy (HALE) for 188 countries, 1990–2013: quantifying the epidemiological transition. Lancet, The, 2015, 386, 2145-2191.	6.3	1,544
334	Global, regional, and national comparative risk assessment of 79 behavioural, environmental and occupational, and metabolic risks or clusters of risks in 188 countries, 1990–2013: a systematic analysis for the Global Burden of Disease Study 2013. Lancet, The, 2015, 386, 2287-2323.	6.3	2,184
335	Subclinical psychosis and suicidal behavior in England: Findings from the 2007 Adult Psychiatric Morbidity Survey. Schizophrenia Research, 2015, 168, 62-67.	1.1	42
336	Social network typologies and mortality risk among older people in China, India, and Latin America: A 10/66 Dementia Research Group population-based cohort study. Social Science and Medicine, 2015, 147, 134-143.	1.8	54
337	The Associations Between Preexisting Mental Disorders andÂSubsequent Onset of Chronic Headaches: A Worldwide Epidemiologic Perspective. Journal of Pain, 2015, 16, 42-52.	0.7	29
338	Global, regional, and national age–sex specific all-cause and cause-specific mortality for 240 causes of death, 1990–2013: a systematic analysis for the Global Burden of Disease Study 2013. Lancet, The, 2015, 385, 117-171.	6.3	5,847
339	Determinants of the components of arterial pressure among older adults – The role of anthropometric and clinical factors: A multi-continent study. Atherosclerosis, 2015, 238, 240-249.	0.4	7
340	Decrease in olfactory and taste receptor expression in the dorsolateral prefrontal cortex in chronic schizophrenia. Journal of Psychiatric Research, 2015, 60, 109-116.	1.5	57
341	Proportion of patients without mental disorders being treated in mental health services worldwide. British Journal of Psychiatry, 2015, 206, 101-109.	1.7	27
342	Is cognitive impairment associated with suicidality? A population-based study. European Neuropsychopharmacology, 2015, 25, 203-213.	0.3	32

#	Article	IF	Citations
343	Psychotic-Like Experiences and Nonsuidical Self-Injury in England: Results from a National Survey. PLoS ONE, 2015, 10, e0145533.	1.1	22
344	Should Unexplained Painful Physical Symptoms be Considered within the Spectrum of Depressive Symptoms?. Clinical Practice and Epidemiology in Mental Health, 2015, 11, 130-139.	0.6	4
345	Long-term functional improvements in the 2-year treatment of schizophrenia outpatients with olanzapine long-acting injection. Neuropsychiatric Disease and Treatment, 2014, 10, 1125.	1.0	5
346	Multimorbidity Patterns in a National Representative Sample of the Spanish Adult Population. PLoS ONE, 2014, 9, e84794.	1.1	73
347	Impact of Multimorbidity on Disability and Quality of Life in the Spanish Older Population. PLoS ONE, 2014, 9, e111498.	1.1	122
348	Chronic Conditions and Sleep Problems among Adults Aged 50 years or over in Nine Countries: A Multi-Country Study. PLoS ONE, 2014, 9, e114742.	1.1	163
349	Horizon 2020 Priorities in Clinical Mental Health Research: Results of a Consensus-Based ROAMER Expert Survey. International Journal of Environmental Research and Public Health, 2014, 11, 10915-10939.	1.2	20
350	Towards Horizon 2020: challenges and advances for clinical mental health research & mp; ndash; outcome of an expert survey. Neuropsychiatric Disease and Treatment, 2014, 10, 1057.	1.0	20
351	Built Environment and Elderly Population Health: A Comprehensive Literature Review. Clinical Practice and Epidemiology in Mental Health, 2014, 10, 103-115.	0.6	106
352	Analysis of burden in caregivers of people with Alzheimer's disease using self-report and supervision hours. Journal of Nutrition, Health and Aging, 2014, 18, 677-684.	1.5	76
353	Are there gender differences in service use for mental disorders across countries in the European Union? Results from the EU-World Mental Health survey. Journal of Epidemiology and Community Health, 2014, 68, 649-656.	2.0	57
354	Factors associated with active aging in Finland, Poland, and Spain. International Psychogeriatrics, 2014, 26, 1363-1375.	0.6	30
355	Health and happiness: cross-sectional household surveys in Finland, Poland and Spain. Bulletin of the World Health Organization, 2014, 92, 716-725.	1.5	54
356	The need for a behavioural science focus in research on mental health and mental disorders. International Journal of Methods in Psychiatric Research, 2014, 23, 28-40.	1.1	38
357	How well can post-traumatic stress disorder be predicted from pre-trauma risk factors? An exploratory study in the WHO World Mental Health Surveys. World Psychiatry, 2014, 13, 265-274.	4.8	194
358	Determinants of Health and Disability in Ageing Population: The COURAGE in Europe Project (Collaborative Research on Ageing in Europe). Clinical Psychology and Psychotherapy, 2014, 21, 193-198.	1.4	70
359	The Validity of the Instrument to Evaluate Social Network in the Ageing Population: The Collaborative Research on Ageing in Europe Social Network Index. Clinical Psychology and Psychotherapy, 2014, 21, 227-241.	1.4	19
360	ROAMER: roadmap for mental health research in Europe. International Journal of Methods in Psychiatric Research, 2014, 23, 1-14.	1,1	109

#	Article	IF	CITATIONS
961	Factors Associated With Components of Arterial Pressure Among Older Individuals (the) Tj ETQq1 1 0.784314 rgB		
361	Consumption. Journal of Clinical Hypertension, 2014, 16, 645-651.	1.0	14
362	A comparative analysis of role attainment and impairment in binge-eating disorder and bulimia nervosa: results from the WHO World Mental Health Surveys. Epidemiology and Psychiatric Sciences, 2014, 23, 27-41.	1.8	46
363	Patient characteristics associated with treatment initiation among paediatric patients with Attention-Deficit/Hyperactivity Disorder symptoms in a naturalistic setting in Central Europe and East Asia. BMC Psychiatry, 2014, 14, 304.	1.1	6
364	The burden of disease in Spain: results from the global burden of disease study 2010. BMC Medicine, 2014, 12, 236.	2.3	27
365	Mood and anxiety disorders across the adult lifespan: a European perspective. Psychological Medicine, 2014, 44, 707-722.	2.7	13
366	Associations between mental disorders and subsequent onset of hypertension. General Hospital Psychiatry, 2014, 36, 142-149.	1.2	102
367	Evaluation of Affect in Mexico and Spain: Psychometric Properties and Usefulness of an Abbreviated Version of the Day Reconstruction Method. Journal of Happiness Studies, 2014, 15, 915-935.	1.9	7
368	Associations between DSM-IV mental disorders and diabetes mellitus: a role for impulse control disorders and depression. Diabetologia, 2014, 57, 699-709.	2.9	69
369	Health-related quality of life in the Cambridge City over-75s Cohort (CC75C): development of a dementia-specific scale and descriptive analyses. BMC Geriatrics, 2014, 14, 18.	1.1	5
370	CUMULATIVE TRAUMAS AND RISK THRESHOLDS: 12-MONTH PTSD IN THE WORLD MENTAL HEALTH (WMH) SURVEYS. Depression and Anxiety, 2014, 31, 130-142.	2.0	329
371	Stratified medicine for mental disorders. European Neuropsychopharmacology, 2014, 24, 5-50.	0.3	152
372	Alcohol abuse in developed and developing countries in the World Mental Health Surveys: Socially defined consequences or psychiatric disorder?. American Journal on Addictions, 2014, 23, 145-155.	1.3	15
373	Description of Agitation and Crisis Interventions in a Psychiatric Hospital in Spain: a Qualitative Study. Value in Health, 2014, 17, A463.	0.1	O
374	Use of Services and Cost of Agitation and Containment in Psychiatric Hospitals: a Systematic Review. Value in Health, 2014, 17, A460.	0.1	0
375	Predictors Pf Remission in the Treatment of Depression in the Middle East: Real-World Evidence From A 6-Month Prospective Observational Study. Value in Health, 2014, 17, A454.	0.1	O
376	Comparative Effectiveness in Terms of Treatment Discontinuation of Orodispersable Versus. Standard Oral Olanzapine Tablets in Non-Adherent Patients: Results from a 1-Year European Outpatient Observational Study. Value in Health, 2014, 17, A766.	0.1	1
377	Symtomatic Factors in Patients With Major Depressive Disorder (MDD): Results from an Observational Study. Value in Health, 2014, 17, A543.	0.1	O
378	Relationship of Insight with Medication Adherence and the Impact on Outcomes in Patients with Schizophrenia and Bipolar Disorder: Results From A 1-Year European Outpatient Observational Study. Value in Health, 2014, 17, A455.	0.1	2

#	Article	IF	CITATIONS
379	Association between mental disorders and subsequent adult onset asthma. Journal of Psychiatric Research, 2014, 59, 179-188.	1.5	58
380	The flavonoid quercetin induces acute vasodilator effects in healthy volunteers: Correlation with beta-glucuronidase activity. Pharmacological Research, 2014, 89, 11-18.	3.1	73
381	Symptomatic remission and patient quality of life in an observational study of schizophrenia: Is there a relationship?. Psychiatry Research, 2014, 220, 163-169.	1.7	26
382	Visual impairment and multimorbidity in a representative sample of the Spanish population. BMC Public Health, 2014, 14, 815.	1.2	38
383	Global, regional, and national incidence and mortality for HIV, tuberculosis, and malaria during 1990–2013: a systematic analysis for the Global Burden of Disease Study 2013. Lancet, The, 2014, 384, 1005-1070.	6.3	786
384	Increased SP4 and SP1 transcription factor expression in the postmortem hippocampus of chronic schizophrenia. Journal of Psychiatric Research, 2014, 58, 189-196.	1.5	23
385	Successful aging, dietary habits and health status of elderly individuals: A k-dimensional approach within the multi-national MEDIS study. Experimental Gerontology, 2014, 60, 57-63.	1.2	70
386	Partial Likelihood Estimation of IRT Models with Censored Lifetime Data: An Application to Mental Disorders in the ESEMeD Surveys. Psychometrika, 2014, 79, 470-488.	1.2	2
387	Relation between jumping to conclusions and cognitive functioning in people with schizophrenia in contrast with healthy participants. Schizophrenia Research, 2014, 159, 211-217.	1.1	34
388	Replication of previous genome-wide association studies of psychiatric diseases in a large schizophrenia case–control sample from Spain. Schizophrenia Research, 2014, 159, 107-113.	1.1	36
389	The state of the art in European research on reducing social exclusion and stigma related to mental health: A systematic mapping of the literature. European Psychiatry, 2014, 29, 381-389.	0.1	110
390	Decreased processing speed might account for working memory span deficit in schizophrenia, and might mediate the associations between working memory span and clinical symptoms. European Psychiatry, 2014, 29, 473-478.	0.1	10
391	Factors associated with suicidal ideation and attempts in Spain for different age groups. Prevalence before and after the onset of the economic crisis. Journal of Affective Disorders, 2014, 163, 1-9.	2.0	81
392	How to deal with missing longitudinal data in cost of illness models in alzheimer's disease – suggestions from the geras study results. Value in Health, 2014, 17, A185.	0.1	0
393	Caregiver Burden in Alzheimer's Disease: Differential Associations in Adult-Child and Spousal Caregivers in the GERAS Observational Study. Dementia and Geriatric Cognitive Disorders Extra, 2014, 4, 51-64.	0.6	77
394	Empirically driven definitions of "good,―"moderate,―and "poor―levels of functioning in the treatment of schizophrenia. Quality of Life Research, 2013, 22, 2085-2094.	1.5	17
395	Association of childhood adversities with the first onset of mental disorders in Spain: results from the ESEMeD project. Social Psychiatry and Psychiatric Epidemiology, 2013, 48, 371-384.	1.6	12
396	Empirical redefinition of delusional disorder and its phenomenology: the DELIREMP study. Comprehensive Psychiatry, 2013, 54, 243-255.	1.5	34

#	Article	IF	Citations
397	Three-factor model of premorbid adjustment in a sample with chronic schizophrenia and first-episode psychosis. Schizophrenia Research, 2013, 151, 252-258.	1.1	17
398	Clinical Outcomes of Patients with Major Depressive Disorder Treated with Either Duloxetine or Selective Serotonin Reuptake Inhibitors in Mexico. Value in Health, 2013, 16, A694.	0.1	0
399	Subjective well-being in schizophrenia: A randomised controlled open-label 12-month non-inferiority study comparing quetiapine XR with risperidone (RECOVER). European Neuropsychopharmacology, 2013, 23, 1257-1269.	0.3	9
400	The Burden of Treatment Change in Major Depressive Disorder: Comparison of Switch Versus Non-Switch Patients in the Perform Study. Value in Health, 2013, 16, A543.	0.1	1
401	Patient-Reported Cognitive Dysfunction Negatively Impacts Functioning in Patients with Major Depressive Disorder – Preliminary Findings from the Perform Study. Value in Health, 2013, 16, A543-A544.	0.1	6
402	The role of negative symptoms in the context of cognitive remediation for schizophrenia. Schizophrenia Research, 2013, 150, 58-63.	1.1	28
403	Functional similarity of facial emotion processing between people with a first episode of psychosis and healthy subjects. Schizophrenia Research, 2013, 149, 35-41.	1.1	12
404	Cognitive profiles of three clusters of patients with a first-episode psychosis. Schizophrenia Research, 2013, 150, 151-156.	1.1	16
405	The Prevalence and Correlates of Binge Eating Disorder in the World Health Organization World Mental Health Surveys. Biological Psychiatry, 2013, 73, 904-914.	0.7	1,025
406	The Mental Component of the Short-Form 12 Health Survey (SF-12) as a Measure of Depressive Disorders in the General Population: Results with Three Alternative Scoring Methods. Value in Health, 2013, 16, 564-573.	0.1	187
407	Predictors of psychiatric hospitalization during 6 months of maintenance treatment with olanzapine long-acting injection: post hoc analysis of a randomized, double-blind study. BMC Psychiatry, 2013, 13, 224.	1.1	6
408	Diagnosis delay in first episodes of major depression: A study of primary care patients in Spain. Journal of Affective Disorders, 2013, 150, 1247-1250.	2.0	20
409	Use of general practitioners versus mental health professionals in six European countries: the decisive role of the organization of mental health-care systems. Social Psychiatry and Psychiatric Epidemiology, 2013, 48, 137-149.	1.6	44
410	Social inequalities in mental health: results from the EU contribution to the World Mental Health Surveys Initiative. Social Psychiatry and Psychiatric Epidemiology, 2013, 48, 173-181.	1.6	53
411	Dissociation in Posttraumatic Stress Disorder: Evidence from the World Mental Health Surveys. Biological Psychiatry, 2013, 73, 302-312.	0.7	251
412	Which somatic symptoms are associated with an unfavorable course in Asian patients with major depressive disorder?. Journal of Affective Disorders, 2013, 149, 182-188.	2.0	71
413	Spanish validation of the Premorbid Adjustment Scale (PAS-S). Comprehensive Psychiatry, 2013, 54, 187-194.	1.5	11
414	Analysis of Sp transcription factors in the postmortem brain of chronic schizophrenia: A pilot study of relationship to negative symptoms. Journal of Psychiatric Research, 2013, 47, 926-934.	1.5	39

#	Article	IF	Citations
415	Usefulness of 2 Questions About Age and Year of Birth in the Case-Finding ofÂDementia. Journal of the American Medical Directors Association, 2013, 14, 627.e7-627.e12.	1.2	4
416	P.3.a.009 SP1 and SP4 transcription factors are increased in the postmortem hippocampus of subjects with chronic schizophrenia. European Neuropsychopharmacology, 2013, 23, S426-S427.	0.3	0
417	Factors Associated with Pain Persistence in Patient with Depression During a 3 –Month Follow-Up Period. Value in Health, 2013, 16, A544.	0.1	0
418	Reduced expression of SP1 and SP4 transcription factors in peripheral blood mononuclear cells in first-episode psychosis. Journal of Psychiatric Research, 2013, 47, 1608-1614.	1.5	27
419	Cognitive correlates of verbal memory and verbal fluency in schizophrenia, and differential effects of various clinical symptoms between male and female patients. Schizophrenia Research, 2013, 147, 81-85.	1.1	28
420	Abnormal functioning of the semantic network in schizophrenia patients with thought disorganization. An exemplar production task. Psychiatry Research, 2013, 205, 1-6.	1.7	10
421	Effects of common mental disorders and physical conditions on role functioning in Spain. Gaceta Sanitaria, 2013, 27, 480-486.	0.6	5
422	The GERAS Study: A Prospective Observational Study of Costs and Resource Use in Community Dwellers with Alzheimer's Disease in Three European Countries – Study Design and Baseline Findings. Journal of Alzheimer's Disease, 2013, 36, 385-399.	1,2	129
423	Validation of an instrument to evaluate quality of life in the aging population: WHOQOL-AGE. Health and Quality of Life Outcomes, 2013, 11, 177.	1.0	89
424	Cross-national differences in the prevalence and correlates of burden among older family caregivers in the World Health Organization World Mental Health (WMH) Surveys. Psychological Medicine, 2013, 43, 865-879.	2.7	68
425	Changes in adherence and treatment costs following initiation of oral or depot typical antipsychotics among previously nonâ€adherent patients with schizophrenia. Human Psychopharmacology, 2013, 28, 438-446.	0.7	5
426	IS ADHD IN CHILDHOOD ASSOCIATED WITH LIFETIME HOARDING SYMPTOMS? AN EPIDEMIOLOGICAL STUDY. Depression and Anxiety, 2013, 30, 741-748.	2.0	41
427	Gender differences in mental health expectancies in early- and midlife in six European countries. British Journal of Psychiatry, 2013, 202, 294-300.	1.7	13
428	Health-related quality-of-life instruments for Alzheimer's disease and mixed dementia. International Psychogeriatrics, 2013, 25, 691-706.	0.6	53
429	Factors Predicting Hostility in Outpatients With Schizophrenia. Journal of Nervous and Mental Disease, 2013, 201, 464-470.	0.5	8
430	Family burden related to mental and physical disorders in the world: results from the WHO World Mental Health (WMH) surveys. Revista Brasileira De Psiquiatria, 2013, 35, 115-125.	0.9	76
431	Do concomitant pain symptoms in patients with major depression affect quality of life even when taking into account baseline depression severity?. Patient Preference and Adherence, 2013, 7, 463.	0.8	13
432	Predictors and consequences of adherence to the treatment of pediatric patients with attention-deficit/hyperactivity disorder in Central Europe and East Asia. Patient Preference and Adherence, 2013, 7, 987.	0.8	27

#	Article	IF	Citations
433	Multi-Country Evaluation of Affective Experience: Validation of an Abbreviated Version of the Day Reconstruction Method in Seven Countries. PLoS ONE, 2013, 8, e61534.	1.1	31
434	Disability Mediates the Impact of Common Conditions on Perceived Health. PLoS ONE, 2013, 8, e65858.	1.1	27
435	Deficits in Executive and Memory Processes in Delusional Disorder: A Case-Control Study. PLoS ONE, 2013, 8, e67341.	1.1	26
436	Psychotropic prescribing in Catalonia: results from an epidemiological study. Family Practice, 2012, 29, 154-162.	0.8	21
437	Role of common mental and physical disorders in partial disability around the world. British Journal of Psychiatry, 2012, 200, 454-461.	1.7	77
438	Comparison of treatment discontinuation and hospitalization among nonadherent patients initiating depot or oral typical antipsychotic medications. International Clinical Psychopharmacology, 2012, 27, 275-282.	0.9	14
439	Parent psychopathology and offspring mental disorders: Results from the WHO World Mental Health Surveys. British Journal of Psychiatry, 2012, 200, 290-299.	1.7	201
440	Antipsychotic Monotherapy and Polypharmacy in the Treatment of Outpatients With Schizophrenia in the European Schizophrenia Outpatient Health Outcomes Study. Journal of Nervous and Mental Disease, 2012, 200, 637-643.	0.5	9
441	Adjustment disorders in primary care: prevalence, recognition and use of services. British Journal of Psychiatry, 2012, 201, 137-142.	1.7	73
442	REPYFLEC cognitive remediation group training in schizophrenia. Schizophrenia Research, 2012, 142, 137-144.	1.1	26
443	Disability-adjusted life years (DALYs) for 291 diseases and injuries in 21 regions, 1990–2010: a systematic analysis for the Global Burden of Disease Study 2010. Lancet, The, 2012, 380, 2197-2223.	6.3	7,061
444	Years lived with disability (YLDs) for 1160 sequelae of 289 diseases and injuries 1990–2010: a systematic analysis for the Global Burden of Disease Study 2010. Lancet, The, 2012, 380, 2163-2196.	6.3	6,376
445	Association between annual earnings and mental disorders in Spain: individual and societal-level estimates. Social Psychiatry and Psychiatric Epidemiology, 2012, 47, 1717-1725.	1.6	6
446	Development of the insight scale for affective disorders (ISAD): Modification from the scale to assess unawareness of mental disorder. Journal of Affective Disorders, 2012, 142, 65-71.	2.0	30
447	PND48 Health State Values in Alzheimer'S Disease – Baseline Results From a Large Prospective Observational Study in Three European Countries (GERAS). Value in Health, 2012, 15, A554.	0.1	0
448	PHS50 Health Care Resource Use and Cost in Alzheimer's Disease in Three European Countries – Baseline Findings of the GERAS Study. Value in Health, 2012, 15, A527.	0.1	2
449	An updated global picture of cigarette smoking persistence among adults. Journal of Epidemiology and Global Health, 2012, 2, 135.	1.1	11
450	Comorbidity of major depressive episode and chronic physical conditions in Spain, a country with low prevalence of depression. General Hospital Psychiatry, 2012, 34, 510-517.	1.2	20

#	Article	IF	CITATIONS
451	Clinical consequences of switching from olanzapine to risperidone and vice versa in outpatients with schizophrenia: 36-month results from the worldwide schizophrenia outpatients health outcomes (W-SOHO) study. BMC Psychiatry, 2012, 12, 218.	1.1	8
452	Risk factors for suicide behaviors in the observational schizophrenia outpatient health outcomes (SOHO) study. BMC Psychiatry, 2012, 12, 83.	1.1	19
453	Schizophrenia Outpatient Health Outcomes study: twelve-month findings. Journal of Pragmatic and Observational Research, 2012, 3, 27.	1.1	4
454	Recognition of anxiety disorders by the general practitioner: results from the DASMAP Study. General Hospital Psychiatry, 2012, 34, 227-233.	1.2	30
455	Regional differences in treatment response and three year course of schizophrenia across the world. Journal of Psychiatric Research, 2012, 46, 856-864.	1.5	16
456	Clinical and Cognitive Correlates of Psychiatric Comorbidity in Delusional Disorder Outpatients. Australian and New Zealand Journal of Psychiatry, 2011, 45, 416-425.	1.3	30
457	Cross-National Comparisons of Sex Differences in Opportunities to Use Alcohol or Drugs, and the Transitions to Use. Substance Use and Misuse, 2011, 46, 1169-1178.	0.7	28
458	Clinical and economic consequences of medication non-adherence in the treatment of patients with a manic/mixed episode of bipolar disorder: Results from the European Mania in Bipolar Longitudinal Evaluation of Medication (EMBLEM) Study. Psychiatry Research, 2011, 190, 110-114.	1.7	96
459	Effectiveness of olanzapine monotherapy and olanzapine combination treatment in the long term following acute mania — Results of a two year observational study in bipolar disorder (EMBLEM). Journal of Affective Disorders, 2011, 131, 320-329.	2.0	16
460	2-year course of bipolar disorder type I patients in outpatient care: Factors associated with remission and functional recovery. European Neuropsychopharmacology, 2011, 21, 287-293.	0.3	60
461	Impact of depressive symptoms on subjective well-being: the importance of patient-reported outcomes in schizophrenia. Patient Preference and Adherence, 2011, 5, 471.	0.8	13
462	Estudio cualitativo sobre la autopercepción del estigma social en personas con esquizofrenia. Revista De La Asociación Española De NeuropsiquiatrÃa, 2011, 31, 477-489.	0.1	13
463	Factors Associated With Use of Psychiatrists and Nonpsychiatrist Providers by ESEMeD Respondents in Six European Countries. Psychiatric Services, 2011, 62, 143-151.	1.1	20
464	Differential 3-Year Effects of First-Versus Second-Generation Antipsychotics on Subjective Well-Being in Schizophrenia Using Marginal Structural Models. Journal of Clinical Psychopharmacology, 2011, 31, 226-230.	0.7	13
465	Cross-national clinical and functional remission rates: Worldwide Schizophrenia Outpatient Health Outcomes (W-SOHO) study. British Journal of Psychiatry, 2011, 199, 194-201.	1.7	148
466	Premarital mental disorders and physical violence in marriage: cross-national study of married couples. British Journal of Psychiatry, 2011, 199, 330-337.	1.7	32
467	The transcription factor SP4 is reduced in postmortem cerebellum of bipolar disorder subjects: control by depolarization and lithium. Bipolar Disorders, 2011, 13, 474-485.	1.1	47
468	Days out of role due to common physical and mental conditions: results from the WHO World Mental Health surveys. Molecular Psychiatry, 2011, 16, 1234-1246.	4.1	360

#	Article	IF	Citations
469	Prevalence and correlates of respiratory and non-respiratory panic attacks in the general population. Journal of Affective Disorders, 2011, 131, 330-338.	2.0	8
470	High and low suicidality in Europe: A fine-grained comparison of France and Spain within the ESEMeD surveys. Journal of Affective Disorders, 2011, 133, 247-256.	2.0	36
471	Complex Interaction Between Symptoms, Social Factors, and Gender in Social Functioning in a Community-Dwelling Sample of Schizophrenia. Psychiatric Quarterly, 2011, 82, 261-274.	1.1	37
472	Gender differences in outcomes of acute mania: a 12-month follow-up study. Archives of Women's Mental Health, 2011, 14, 107-113.	1.2	9
473	Dropping out of mental health treatment among patients with depression and anxiety by type of provider: results of the European Study of the Epidemiology of Mental Disorders. Social Psychiatry and Psychiatric Epidemiology, 2011, 46, 273-280.	1.6	19
474	Measuring use of services for mental health problems in epidemiological surveys. International Journal of Methods in Psychiatric Research, 2011, 20, 182-191.	1.1	8
475	Use of Primary and Specialized Mental Health Care for a Major Depressive Episode in Spain by ESEMeD Respondents. Psychiatric Services, 2011, 62, 152-161.	1.1	40
476	Development of Lifetime Comorbidity in the World Health Organization World Mental Health Surveys. Archives of General Psychiatry, 2011, 68, 90.	13.8	410
477	Including information about co-morbidity in estimates of disease burden: results from the World Health Organization World Mental Health Surveys. Psychological Medicine, 2011, 41, 873-886.	2.7	49
478	Association of Childhood Adversities and Early-Onset Mental Disorders With Adult-Onset Chronic Physical Conditions. Archives of General Psychiatry, 2011, 68, 838.	13.8	281
479	Comparison of olanzapine and risperidone in the EMBLEM Study: translation of randomized controlled trial findings into clinical practice. International Clinical Psychopharmacology, 2010, 25, 257-263.	0.9	4
480	Incidence of Extrapyramidal Symptoms and Tardive Dyskinesia in Schizophrenia. Journal of Clinical Psychopharmacology, 2010, 30, 531-540.	0.7	68
481	Chronic Physical Conditions and Their Association With First Onset of Suicidal Behavior in the World Mental Health Surveys. Psychosomatic Medicine, 2010, 72, 712-719.	1.3	178
482	Burden of chronic physical conditions and mental disorders in primary care. British Journal of Psychiatry, 2010, 196, 302-309.	1.7	63
483	Childhood adversities as risk factors for onset and persistence of suicidal behaviour. British Journal of Psychiatry, 2010, 197, 20-27.	1.7	278
484	Associations of serious mental illness with earnings: results from the WHO World Mental Health surveys. British Journal of Psychiatry, 2010, 197, 114-121.	1.7	156
485	Seeking Help for Mental Health Problems outside the Conventional Health Care System: Results from the European Study of the Epidemiology of Mental Disorders (ESEMeD). Canadian Journal of Psychiatry, 2010, 55, 586-597.	0.9	10
486	Prevalence of mental disorders in primary care: results from the diagnosis and treatment of mental disorders in primary care study (DASMAP). Social Psychiatry and Psychiatric Epidemiology, 2010, 45, 201-210.	1.6	137

#	Article	IF	CITATIONS
487	Utility of the twelve-item World Health Organization Disability Assessment Schedule II (WHO-DAS II) for discriminating depression "caseness―and severity in Spanish primary care patients. Quality of Life Research, 2010, 19, 97-101.	1.5	30
488	The 12-item World Health Organization Disability Assessment Schedule II (WHO-DAS II): a nonparametric item response analysis. BMC Medical Research Methodology, 2010, 10, 45.	1.4	67
489	Is major depression adequately diagnosed and treated by general practitioners? Results from an epidemiological study. General Hospital Psychiatry, 2010, 32, 201-209.	1.2	81
490	Epidemiology of major depressive episode in a southern European country: Results from the ESEMeD-Spain project. Journal of Affective Disorders, 2010, 120, 76-85.	2.0	98
491	Psychometric properties of the twelve item World Health Organization Disability Assessment Schedule II (WHO-DAS II) in Spanish primary care patients with a first major depressive episode. Journal of Affective Disorders, 2010, 121, 52-58.	2.0	78
492	Obsessive–compulsive symptom dimensions in the general population: Results from an epidemiological study in six European countries. Journal of Affective Disorders, 2010, 124, 291-299.	2.0	105
493	Age differences in the prevalence and co-morbidity of DSM-IV major depressive episodes: results from the WHO World Mental Health Survey Initiative. Depression and Anxiety, 2010, 27, 351-364.	2.0	363
494	Factor structure, internal consistency and construct validity of the Sheehan Disability Scale in a Spanish primary care sample. Journal of Evaluation in Clinical Practice, 2010, 16, 895-901.	0.9	54
495	Frequent attendance in primary care: comparison and implications of different definitions. British Journal of General Practice, 2010, 60, e49-e55.	0.7	45
496	Childhood adversities and adult psychopathology in the WHO World Mental Health Surveys. British Journal of Psychiatry, 2010, 197, 378-385.	1.7	1,854
497	Early Cannabis Use and Estimated Risk of Later Onset of Depression Spells: Epidemiologic Evidence From the Population-based World Health Organization World Mental Health Survey Initiative. American Journal of Epidemiology, 2010, 172, 149-159.	1.6	64
498	Gender and the relationship between marital status and first onset of mood, anxiety and substance use disorders. Psychological Medicine, 2010, 40, 1495-1505.	2.7	133
499	Assessment of Medication Adherence in a Cohort of Patients with Bipolar Disorder. Pharmacopsychiatry, 2010, 43, 263-270.	1.7	54
500	Health status of the advanced elderly in six european countries: results from a representative survey using EQ-5D and SF-12. Health and Quality of Life Outcomes, 2010, 8, 143.	1.0	98
501	Predictors and clinical consequences of non-adherence with antipsychotic medication in the outpatient treatment of schizophrenia. Psychiatry Research, 2010, 176, 109-113.	1.7	359
502	Gender differences in delusional disorder: Evidence from an outpatient sample. Psychiatry Research, 2010, 177, 235-239.	1.7	32
503	Work impairment in bipolar disorder patients – results from a two-year observational study (EMBLEM). European Psychiatry, 2010, 25, 338-344.	0.1	29
504	The Cost of Relapse for Patients with a Manic/Mixed Episode of Bipolar Disorder in the EMBLEM Study. Pharmacoeconomics, 2010, 28, 555-566.	1.7	22

#	Article	IF	Citations
505	Why Do Clinicians Maintain Antidepressants in Some Patients With Acute Mania?. Journal of Clinical Psychiatry, 2010, 71, 1000-1006.	1.1	43
506	Gender differences in incipient psychosis. European Journal of Psychiatry, 2010, 24, .	0.7	8
507	Implications of modifying the duration requirement of generalized anxiety disorder in developed and developing countries. Psychological Medicine, 2009, 39, 1163.	2.7	41
508	Cross-National Associations Between Gender and Mental Disorders in the World Health Organization World Mental Health Surveys. Archives of General Psychiatry, 2009, 66, 785.	13.8	1,018
509	Translation of Randomised Controlled Trial Findings into Clinical Practice: Comparison of Olanzapine and Valproate in the EMBLEM Study. Pharmacopsychiatry, 2009, 42, 145-152.	1.7	9
510	Reasons and Outcomes of Olanzapine Dose Adjustments in the Outpatient Treatment of Schizophrenia. Pharmacopsychiatry, 2009, 42, 135-140.	1.7	9
511	Dog-Assisted Therapy in the Treatment of Chronic Schizophrenia Inpatients. Anthrozoos, 2009, 22, 149-159.	0.7	44
512	Cross-National Analysis of the Associations among Mental Disorders and Suicidal Behavior: Findings from the WHO World Mental Health Surveys. PLoS Medicine, 2009, 6, e1000123.	3.9	592
513	Antipsychotic treatment discontinuation in previously untreated patients with schizophrenia: 36-month results from the SOHO study. Journal of Psychiatric Research, 2009, 43, 265-273.	1.5	36
514	The epidemiology of eating disorders in six European countries: Results of the ESEMeD-WMH project. Journal of Psychiatric Research, 2009, 43, 1125-1132.	1.5	373
515	Suicide ideation across reproductive life cycle of women Results from a European epidemiological study. Journal of Affective Disorders, 2009, 116, 144-147.	2.0	31
516	Pain and depression in older people: Comorbidity and patterns of help seeking. Journal of Affective Disorders, 2009, 117, 193-196.	2.0	43
517	Perceived stigma among individuals with common mental disorders. Journal of Affective Disorders, 2009, 118, 180-186.	2.0	152
518	Predictors of switching from mania to depression in a large observational study across Europe (EMBLEM). Journal of Affective Disorders, 2009, 118, 118-123.	2.0	33
519	Impact of mental disorders and chronic physical conditions in health-related quality of life among primary care patients: results from an epidemiological study. Quality of Life Research, 2009, 18, 1011-1018.	1.5	21
520	Fluoxetine and imipramine: are there differences in costâ€utility for depression in primary care?. Journal of Evaluation in Clinical Practice, 2009, 15, 195-203.	0.9	9
521	Worldwide-Schizophrenia Outpatient Health Outcomes (W-SOHO): baseline characteristics of pan-regional observational data from more than 17,000 patients*. International Journal of Clinical Practice, 2009, 63, 1578-1588.	0.8	63
522	Comparison of Different Valuation Methods for Population Health Status Measured by the EQ-5D in Three European Countries. Value in Health, 2009, 12, 750-758.	0.1	78

#	Article	IF	CITATIONS
523	Adaptation and validation of the quality-of-life scale: Satisfaction with Life Domains Scale by Baker and Intagliata. Comprehensive Psychiatry, 2009, 50, 76-80.	1.5	17
524	Is it always the same? Variability of depressive symptoms across six European countries. Psychiatry Research, 2009, 168, 137-144.	1.7	19
525	Observational study designs for bipolar disorder â€" What can they tell us about treatment in acute mania?. Progress in Neuro-Psychopharmacology and Biological Psychiatry, 2009, 33, 715-721.	2.5	13
526	The cost of relapse in patients with schizophrenia in the European SOHO (Schizophrenia Outpatient) Tj ETQq0 0 0 835-841.	0 rgBT /Ov 2.5	erlock 10 Tf 63
527	Childhood Predictors of Adult Attention-Deficit/Hyperactivity Disorder: Results from the World Health Organization World Mental Health Survey Initiative. Biological Psychiatry, 2009, 65, 46-54.	0.7	346
528	Recovery in the outpatient setting: 36-month results from the Schizophrenia Outpatients Health Outcomes (SOHO) study. Schizophrenia Research, 2009, 108, 223-230.	1.1	111
529	Religion and Spiritual Advice as Surrogate Care Provision for Mental Health Problems in the ESEMED Survey. European Psychiatry, 2009, 24, .	0.1	О
530	Evaluación telefónica de los sistemas de activación e inhibición conductual. Revista De PsiquiatrÃa Y Salud Mental, 2009, 2, 66-71.	1.0	6
531	Un estudio empÃrico de los correlatos psicosociales y clÃnicos del trastorno delirante: el estudio DELIREMP. Revista De PsiquiatrÃa Y Salud Mental, 2009, 2, 72-82.	1.0	12
532	Variabilidad en el gen COMT y modificación del riesgo de esquizofrenia conferido por consumo de cannabis. Revista De PsiquiatrÃa Y Salud Mental, 2009, 2, 89-94.	1.0	8
533	Evaluation of behavioural activation and inhibition systems by telephone. Revista De PsiquiatrÃa Y Salud Mental (English Edition), 2009, 2, 66-71.	0.2	1
534	An empirical study of psychosocial and clinical correlates of delusional disorder: the DELIREMP study. Revista De PsiquiatrÃa Y Salud Mental (English Edition), 2009, 2, 72-82.	0.2	4
535	Tolerability of outpatient antipsychotic treatment: 36-month results from the European Schizophrenia Outpatient Health Outcomes (SOHO) study. European Neuropsychopharmacology, 2009, 19, 542-550.	0.3	33
536	PMH25 CLINICAL AND ECONOMIC CONSEQUENCES OF MEDICATION NONADHERENCE IN THE TREATMENT OF PATIENTS WITH A MANIC/ MIXED EPISODE OF BIPOLAR DISORDER: RESULTS FROM THE EUROPEAN MANIA IN BIPOLAR LONGITUDINAL EVALUATION OF MEDICATION (EMBLEM) STUDY. Value in Health, 2009, 12, A355.	0.1	0
537	Comparison of Population Health Status in Six European Countries. Medical Care, 2009, 47, 255-261.	1.1	191
538	Validity and Reliability of the Hamilton Depression Rating Scale (5 Items) for Manic and Mixed Bipolar Disorders. Journal of Nervous and Mental Disease, 2009, 197, 682-686.	0.5	22
539	Perceived Need for Mental Health Care and Service Use Among Adults in Western Europe: Results of the ESEMeD Project. Psychiatric Services, 2009, 60, 1051-1058.	1.1	68
540	Association of headache with childhood adversity and mental disorder: cross-national study. British Journal of Psychiatry, 2009, 194, 111-116.	1.7	33

#	Article	IF	CITATIONS
541	Mental–physical co-morbidity and its relationship with disability: results from the World Mental Health Surveys. Psychological Medicine, 2009, 39, 33-43.	2.7	239
542	Mental disorders and termination of education in high-income and low- and middle-income countries: epidemiological study. British Journal of Psychiatry, 2009, 194, 411-417.	1.7	129
543	Assessment of remission in schizophrenia with the CGI and CGIâ€SCH scales. Acta Psychiatrica Scandinavica, 2008, 117, 156-156.	2.2	3
544	Adverse parenting as a risk factor in the occurrence of anxiety disorders. Social Psychiatry and Psychiatric Epidemiology, 2008, 43, 266-272.	1.6	36
545	Do needs, symptoms or disability of outpatients with schizophrenia influence family burden?. Social Psychiatry and Psychiatric Epidemiology, 2008, 43, 612-618.	1.6	57
546	Main traumatic events in Europe: PTSD in the European study of the epidemiology of mental disorders survey. Journal of Traumatic Stress, 2008, 21, 455-462.	1.0	218
547	Rates and predictors of remission and recovery during 3 years in 392 neverâ€treated patients with schizophrenia. Acta Psychiatrica Scandinavica, 2008, 118, 220-229.	2.2	145
548	Association of perceived stigma and mood and anxiety disorders: results from the World Mental Health Surveys. Acta Psychiatrica Scandinavica, 2008, 118, 305-314.	2.2	182
549	Obesity and mental disorders in the general population: results from the world mental health surveys. International Journal of Obesity, 2008, 32, 192-200.	1.6	348
550	Development of EMPRO: A Tool for the Standardized Assessment of Patient-Reported Outcome Measures. Value in Health, 2008, 11 , 700-708.	0.1	166
551	Comorbid painful physical symptoms and anxiety: Prevalence, work loss and help-seeking. Journal of Affective Disorders, 2008, 109, 264-272.	2.0	23
552	Clinical factors influencing the prescription of antidepressants and benzodiazepines:. Journal of Affective Disorders, 2008, 110, 84-93.	2.0	128
553	Rapid-cycling bipolar I disorder: Course and treatment outcome of a large sample across Europe. Journal of Psychiatric Research, 2008, 42, 1068-1075.	1.5	56
554	Mental disorders among persons with diabetesâ€"Results from the World Mental Health Surveys. Journal of Psychosomatic Research, 2008, 65, 571-580.	1.2	154
555	Marginal structural models might overcome confounding when analyzing multiple treatment effects in observational studies. Journal of Clinical Epidemiology, 2008, 61, 525-530.	2.4	33
556	The relation between multiple pains and mental disorders: Results from the World Mental Health Surveys. Pain, 2008, 135, 82-91.	2.0	260
557	Predictors of the course of illness in outpatients with schizophrenia: A prospective three year study. Progress in Neuro-Psychopharmacology and Biological Psychiatry, 2008, 32, 1287-1292.	2.5	39
558	The Schizophrenia Outpatient Health Outcomes (SOHO) study: 3-year results of antipsychotic treatment discontinuation and related clinical factors in Spain. European Psychiatry, 2008, 23, 1-7.	0.1	28

#	Article	IF	CITATIONS
559	A descriptive case-register study of delusional disorder. European Psychiatry, 2008, 23, 125-133.	0.1	54
560	Previously untreated patients with schizophrenia: The nnt for all causes of treatment discontinuation and the nnh for weight gain. European Psychiatry, 2008, 23, S166.	0.1	0
561	Stability and treatment outcome of distinct classes of mania. European Psychiatry, 2008, 23, 360-367.	0.1	18
562	Remission in the outpatient care of schizophrenia: 3-Year results from the Schizophrenia Outpatients Health Outcomes (SOHO) Study in France. European Psychiatry, 2008, 23, 491-496.	0.1	21
563	Common Chronic Pain Conditions in Developed and Developing Countries: Gender and Age Differences and Comorbidity With Depression-Anxiety Disorders. Journal of Pain, 2008, 9, 883-891.	0.7	918
564	What is the relative importance of self reported psychotic symptoms in epidemiological studies? Results from the ESEMeDâ€"Catalonia Study. Schizophrenia Research, 2008, 102, 261-269.	1.1	39
565	Cost-Utility Analysis of Treatment with Olanzapine Compared with Other Antipsychotic Treatments in Patients with Schizophrenia in the Pan-European SOHO Study. Pharmacoeconomics, 2008, 26, 341-358.	1.7	25
566	Chronic mania revisited: Factors associated with treatment non-response during prospective follow-up of a large European cohort (EMBLEM). World Journal of Biological Psychiatry, 2008, 9, 313-320.	1.3	22
567	The prevalence and effects of adult attention-deficit/hyperactivity disorder (ADHD) on the performance of workers: results from the WHO World Mental Health Survey Initiative. Occupational and Environmental Medicine, 2008, 65, 835-842.	1.3	262
568	Cross-national prevalence and risk factors for suicidal ideation, plans and attempts. British Journal of Psychiatry, 2008, 192, 98-105.	1.7	1,993
569	Overview of the findings from the European SOHO study. Expert Review of Neurotherapeutics, 2008, 8, 873-880.	1.4	18
570	Determinants of Voluntary vs. Involuntary Admission in Bipolar Disorder and the Impact of Adherence. Pharmacopsychiatry, 2008, 41, 29-36.	1.7	24
571	Mental disorders among persons with arthritis: results from the World Mental Health Surveys. Psychological Medicine, 2008, 38, 1639-1650.	2.7	81
572	The Relation between Body Mass Index, Mental Health, and Functional Disability: A European Population Perspective. Canadian Journal of Psychiatry, 2008, 53, 679-688.	0.9	47
573	Disability and treatment of specific mental and physical disorders across the world. British Journal of Psychiatry, 2008, 192, 368-375.	1.7	326
574	Childhood Adversity, Early-Onset Depressive/Anxiety Disorders, and Adult-Onset Asthma. Psychosomatic Medicine, 2008, 70, 1035-1043.	1.3	91
575	Adequacy of Antidepressant Treatment in Spanish Primary Care: A Naturalistic Six-Month Follow-Up Study. Psychiatric Services, 2008, 59, 78-83.	1.1	27
576	Clinical consequences of switching antipsychotic drugs in outpatients with schizophrenia: 36-month results from the European Schizophrenia Outpatient Health Outcomes study. International Clinical Psychopharmacology, 2008, 23, 203-208.	0.9	4

#	Article	IF	Citations
577	Estudio SOHO (Variables Relacionadas con la Salud en Pacientes Ambulatorios con Esquizofrenia): resultados a los 3 años de la suspensión del tratamiento antipsicótico y factores clĀnicos relacionados con ella en España. European Psychiatry (Ed Española), 2008, 15, 137-144.	0.0	0
578	Estudio descriptivo de un registro de casos de trastorno delirante. European Psychiatry (Ed) Tj ETQq0 0 0 rgBT /C	Overlock 1	0 Tf 50 702 T
579	Cross-national prevalence and correlates of adult attention-deficit hyperactivity disorder. British Journal of Psychiatry, 2007, 190, 402-409.	1.7	1,118
580	Population level of unmet need for mental healthcare in Europe. British Journal of Psychiatry, 2007, 190, 299-306.	1.7	222
581	Parental Bonding and Suicidality in Adulthood. Australian and New Zealand Journal of Psychiatry, 2007, 41, 66-73.	1.3	36
582	Sexual Intercourse Frequency in Men Presenting for Treatment of Erectile Dysfunction: Results from the Pan-European Erectile Dysfunction Observational Study. Journal of Sex and Marital Therapy, 2007, 33, 3-18.	1.0	11
583	Treatment adequacy for anxiety and depressive disorders in six European countries. British Journal of Psychiatry, 2007, 190, 172-173.	1.7	147
584	Differences in Lifetime Use of Services for Mental Health Problems in Six European Countries. Psychiatric Services, 2007, 58, 213-220.	1.1	103
585	Gender differences in response to antipsychotic treatment in outpatients with schizophrenia. Psychiatry Research, 2007, 153, 225-231.	1.7	103
586	Mental disorders among persons with chronic back or neck pain: Results from the world mental health surveys. Pain, 2007, 129, 332-342.	2.0	471
587	Three-year antipsychotic effectiveness in the outpatient care of schizophrenia: Observational versus randomized studies results. European Neuropsychopharmacology, 2007, 17, 235-244.	0.3	106
588	Number needed to treat (NNT) for all-cause medication discontinuation in catie compared to the schizophrenia outpatients health outcomes (SOHO) study. European Psychiatry, 2007, 22, S130.	0.1	0
589	Use of mental health services for anxiety, mood, and substance disorders in 17 countries in the WHO world mental health surveys. Lancet, The, 2007, 370, 841-850.	6.3	1,271
590	Assessment of remission in schizophrenia with the CGI and CGI-SCH scales. Acta Psychiatrica Scandinavica, 2007, 115, 163-164.	2.2	23
591	Patterns of switching phosphodiesterase type 5 inhibitors in the treatment of erectile dysfunction: results from the Erectile Dysfunction Observational Study. International Journal of Clinical Practice, 2007, 61, 1850-1862.	0.8	28
592	Mental disorders among adults with asthma: results from the World Mental Health Survey. General Hospital Psychiatry, 2007, 29, 123-133.	1.2	251
593	Mental disorders among persons with heart disease â€" results from World Mental Health surveys. General Hospital Psychiatry, 2007, 29, 325-334.	1.2	204
594	Risk factors for suicidality in Europe: Results from the ESEMED study. Journal of Affective Disorders, 2007, 101, 27-34.	2.0	465

#	Article	IF	Citations
595	Depression–anxiety relationships with chronic physical conditions: Results from the World Mental Health surveys. Journal of Affective Disorders, 2007, 103, 113-120.	2.0	414
596	Therapeutic Effectiveness and Patient Satisfaction after 6 Months of Treatment with Tadalafil, Sildenafil, and Vardenafil: Results from the Erectile Dysfunction Observational Study (EDOS). European Urology, 2007, 51, 541-550.	0.9	54
597	Service utilization and costs of first-onset schizophrenia in two widely differing health service areas in North-East Spain. Acta Psychiatrica Scandinavica, 2007, 100, 335-343.	2.2	21
598	Symptomatic remission in previously untreated patients with schizophrenia: 2-year results from the SOHO study. Psychopharmacology, 2007, 191, 1015-1022.	1.5	33
599	The social, psychopathological and consumer context of rate of symptom improvement in acute mania. Social Psychiatry and Psychiatric Epidemiology, 2007, 42, 631-638.	1.6	9
600	Selective serotonin reuptake inhibitors: are there differences in effectiveness in primary care?. Primary Care and Community Psychiatry, 2007, 11, 113-120.	0.4	3
601	Social functioning in schizophrenia: what is the influence of gender?. European Journal of Psychiatry, 2007, 21, .	0.7	11
602	Delay and failure in treatment seeking after first onset of mental disorders in the World Health Organization's World Mental Health Survey Initiative. World Psychiatry, 2007, 6, 177-85.	4.8	351
603	Lifetime prevalence and age-of-onset distributions of mental disorders in the World Health Organization's World Mental Health Survey Initiative. World Psychiatry, 2007, 6, 168-76.	4.8	1,239
604	Validation in Spanish population of the family objective and subjective burden interview (ECFOS-II) for relatives of patients with schizophrenia. Actas Espanolas De Psiquiatria, 2007, 35, 372-81.	0.1	15
605	The SOHO (Schizophrenia Outpatient Health Outcome) Study. CNS Drugs, 2006, 20, 293-301.	2.7	74
606	Methodological Aspects in the Assessment of Treatment Effects in Observational Health Outcomes Studies. Applied Health Economics and Health Policy, 2006, 5, 11-25.	1.0	41
607	P.2.e.006 Outcomes of acute mania: 12 months results from the European mania in bipolar longitudinal evaluation of medication (EMBLEM) study. European Neuropsychopharmacology, 2006, 16, S349.	0.3	0
608	P.3.a.042 Clinical course, social functioning, quality of life & medication tolerability in previously untreated patients with schizophrenia: 36-month results from the European schizophrenia outpatient health outcomes (SOHO) study. European Neuropsychopharmacology, 2006, 16, S387-S388.	0.3	0
609	Antipsychotic type and correlates of antipsychotic treatment discontinuation in the outpatient treatment of schizophrenia. European Psychiatry, 2006, 21, 41-47.	0.1	41
610	0428 EFFECTIVENESS OF ANTIPSYCHOTIC TREATMENT IN NEVER TREATED PATIENTS WITH SCHIZOPHRENIA: 36-MONTH RESULTS FROM THE SCHIZOPHRENIA OUTPATIENTS HEALTH OUTCOMES (SOHO) STUDY. Schizophrenia Research, 2006, 86, S131-S132.	1.1	1
611	PMH35 THE COST OF RELAPSE IN PATIENTS WITH SCHIZOPHRENIA IN THE PAN-EUROPEAN SOHO (SCHIZOPHRENIA OUTPATIENT HEALTH OUTCOMES) STUDY. Value in Health, 2006, 9, A75.	0.1	0
612	Estimating the prevalence of schizophrenia in Spain using a disease model. Schizophrenia Research, 2006, 86, 194-201.	1.1	33

#	Article	IF	Citations
613	Neurocognitive performance and negative symptoms: Are they equal in explaining disability in schizophrenia outpatients?. Schizophrenia Research, 2006, 87, 246-253.	1.1	75
614	Four symptom dimensions in outpatients with schizophrenia. Comprehensive Psychiatry, 2006, 47, 384-388.	1.5	11
615	Methodological approach for assessing the cost-effectiveness of treatments using longitudinal observational data: The SOHO study. International Journal of Technology Assessment in Health Care, 2006, 22, 460-468.	0.2	10
616	Remission and Relapse in the Outpatient Care of Schizophrenia. Journal of Clinical Psychopharmacology, 2006, 26, 571-578.	0.7	112
617	Evidence for three distinct classes of 'typical', 'psychotic' and 'dual' mania: results from the EMBLEM study. Acta Psychiatrica Scandinavica, 2006, 113, 112-120.	2.2	75
618	Treatmentâ€Seeking Behavior of Erectile Dysfunction Patients in Europe: Results of the Erectile Dysfunction Observational Study. Journal of Sexual Medicine, 2006, 3, 530-540.	0.3	35
619	A framework for evidence-based mental health care and policy. Acta Psychiatrica Scandinavica, 2006, 114, 5-11.	2.2	24
620	Assessment of service use patterns in out-patients with schizophrenia: a Spanish study. Acta Psychiatrica Scandinavica, 2006, 114, 12-18.	2.2	13
621	Cost-effectiveness of different clinical interventions for reducing the burden of schizophrenia in Spain. Acta Psychiatrica Scandinavica, 2006, 114, 29-38.	2.2	29
622	Cost?utility of selective serotonin reuptake inhibitors for depression in primary care in Catalonia. Acta Psychiatrica Scandinavica, 2006, 114, 39-47.	2.2	21
623	Effectiveness and cost-effectiveness of antidepressant treatment in primary health care: A six-month randomised study comparing fluoxetine to imipramine. Journal of Affective Disorders, 2006, 91, 153-163.	2.0	18
624	Gender differences in response to antidepressant treatment prescribed in primary care. Does menopause make a difference?. Journal of Affective Disorders, 2006, 93, 53-60.	2.0	79
625	Treatment adequacy of anxiety and depressive disorders: Primary versus specialised care in Spain. Journal of Affective Disorders, 2006, 96, 9-20.	2.0	60
626	Concordance of the Composite International Diagnostic Interview Version 3.0 (CIDI 3.0) with standardized clinical assessments in the WHO World Mental Health Surveys. International Journal of Methods in Psychiatric Research, 2006, 15, 167-180.	1.1	949
627	Open-label olanzapine treatment in bipolar I disorder: clinical and work functional outcomes. Current Medical Research and Opinion, 2006, 22, 961-966.	0.9	13
628	The World Health Organization World Mental Health Survey Initiative. Epidemiologia E Psichiatria Sociale, 2006, 15, 161-166.	1.0	81
629	Financing Mental Health Care in Spain: Context and critical issues. European Journal of Psychiatry, 2006, 20, .	0.7	22
630	Influence of age at onset on social functioning in outpatients with schizophrenia. European Journal of Psychiatry, 2006, 20, .	0.7	19

#	Article	IF	CITATIONS
631	Pharmacological treatment and other predictors of treatment outcomes in previously untreated patients with schizophrenia: results from the European Schizophrenia Outpatient Health Outcomes (SOHO) study. International Clinical Psychopharmacology, 2005, 20, 199-205.	0.9	44
632	Assessing depression in primary care with the PHQ-9: Can it be carried out over the telephone?. Journal of General Internal Medicine, 2005, 20, 738-742.	1.3	298
633	Effectiveness of antipsychotic treatment for schizophrenia: 6-month results of the Pan-European Schizophrenia Outpatient Health Outcomes (SOHO) study. Acta Psychiatrica Scandinavica, 2005, 111, 220-231.	2.2	83
634	Olanzapine vs. other antipsychotics in actual out-patient settings: six months tolerability results from the European Schizophrenia Out-patient Health Outcomes study. Acta Psychiatrica Scandinavica, 2005, 111, 232-243.	2.2	65
635	PMH27 12-MONTH COST-EFFECTIVENESS ANALYSIS OF ORAL ANTIPSYCHOTIC TREATMENTS IN PATIENTS WITH SCHIZOPHRENIA IN THE PAN-EUROPEAN SOHO (SCHIZOPHRENIA OUTPATIENT HEALTH OUTCOMES) STUDY. Value in Health, 2005, 8, A204.	0.1	O
636	PMH31 12 MONTH COST-UTILITY ANALYSIS OF ORAL ANTIPSYCHOTIC TREATMENTS IN PATIENTS WITH SCHIZOPHRENIA IN THE PAN-EUROPEAN SOHO (SCHIZOPHRENIA OUTPATIENT HEALTH OUTCOMES) STUDY. Value in Health, 2005, 8, A205.	0.1	0
637	Use of concomitant medication with antipsychotic treatment in outpatients with schizophrenia: Results from the European Schizophrenia Outpatients Health Outcomes (SOHO) study. Progress in Neuro-Psychopharmacology and Biological Psychiatry, 2005, 29, 972-982.	2.5	28
638	Needs and its relation to symptom dimensions in a sample of outpatients with schizophrenia. Schizophrenia Research, 2005, 75, 129-134.	1.1	28
639	Prevalence, Severity, and Unmet Need for Treatment of Mental Disorders in the World Health Organization World Mental Health Surveys. JAMA - Journal of the American Medical Association, 2004, 291, 2581.	3.8	2,696
640	Sampling and methods of the European Study of the Epidemiology of Mental Disorders (ESEMeD) project. Acta Psychiatrica Scandinavica, 2004, 109, 8-20.	2.2	136
641	Prevalence of mental disorders in Europe: results from the European Study of the Epidemiology of Mental Disorders (ESEMeD) project. Acta Psychiatrica Scandinavica, 2004, 109, 21-27.	2.2	1,053
642	12-Month comorbidity patterns and associated factors in Europe: results from the European Study of the Epidemiology of Mental Disorders (ESEMeD) project. Acta Psychiatrica Scandinavica, 2004, 109, 28-37.	2.2	157
643	Disability and quality of life impact of mental disorders in Europe: results from the European Study of the Epidemiology of Mental Disorders (ESEMeD) project. Acta Psychiatrica Scandinavica, 2004, 109, 38-46.	2.2	285
644	Use of mental health services in Europe: results from the European Study of the Epidemiology of Mental Disorders (ESEMeD) project. Acta Psychiatrica Scandinavica, 2004, 109, 47-54.	2.2	333
645	Psychotropic drug utilization in Europe: results from the European Study of the Epidemiology of Mental Disorders (ESEMeD) project. Acta Psychiatrica Scandinavica, 2004, 109, 55-64.	2.2	143
646	Unit costs in international economic evaluations: resource costing of the Schizophrenia Outpatient Health Outcomes Study. Acta Psychiatrica Scandinavica, 2003, 107, 41-47.	2.2	6
647	The European Schizophrenia Outpatient Health Outcomes Study: baseline findings across country and treatment. Acta Psychiatrica Scandinavica, 2003, 107, 7-15.	2.2	44
648	The Clinical Global Impression-Schizophrenia scale: a simple instrument to measure the diversity of symptoms present in schizophrenia. Acta Psychiatrica Scandinavica, 2003, 107, 16-23.	2.2	508

#	Article	IF	Citations
649	The European Schizophrenia Outpatient Health Outcomes (SOHO) study: rationale, methods and recruitment. Acta Psychiatrica Scandinavica, 2003, 107, 222-232.	2.2	98
650	Clinical and functional status over the first three months of treatment: Results from the schizophrenia outpatient health outcomes (SOHO) study. Schizophrenia Research, 2003, 60, 282.	1.1	1
651	Met and Unmet Needs of Schizophrenia Patients in a Spanish Sample. Schizophrenia Bulletin, 2003, 29, 201-210.	2.3	77
652	Olanzapine versus haloperidol in the treatment of acute mania: clinical outcomes, health-related quality of life and work status. International Clinical Psychopharmacology, 2002, 17, 227-237.	0.9	55
653	Olanzapine versus haloperidol as initial treatment in acute mania: Prospective comparison of work functional outcomes. European Neuropsychopharmacology, 2002, 12, 307.	0.3	1
654	Statistical models for population heterogeneity in epidemiological longitudinal studies: Person centred latent class analysis to identify casemix groups in the schizophrenia outpatient health outcomes study. European Neuropsychopharmacology, 2002, 12, 314.	0.3	0
655	The treatment of schizophrenia across Europe: Between country baseline differences in the schizophrenia outpatient health outcomes (SOHO) study. European Neuropsychopharmacology, 2002, 12, 317.	0.3	0
656	The European Study of the Epidemiology of Mental Disorders (ESEMeD/MHEDEA 2000) project: rationale and methods. International Journal of Methods in Psychiatric Research, 2002, 11, 55-67.	1.1	85
657	Influence of gender on social outcome in schizophrenia. Acta Psychiatrica Scandinavica, 2002, 106, 337-342.	2.2	63
658	Assessment instruments: standardization of the European Service Mapping Schedule (ESMS) in Spain. Acta Psychiatrica Scandinavica, 2000, 102, 24-32.	2.2	8
659	Utilisation of mental health services and costs of patients with schizophrenia in three areas of Spain. British Journal of Psychiatry, 1998, 173, 334-340.	1.7	57
660	The Relationship Between Plasma Haloperidol Concentrations and Clinical Results. Archives of General Psychiatry, 1996, 53, 1167.	13.8	5
661	Predictability of rehospitalization for schizophrenia. European Archives of Psychiatry and Clinical Neuroscience, 1994, 244, 241-246.	1.8	26
662	Emotional Distress in Disaster Victims. Journal of Nervous and Mental Disease, 1993, 181, 388-393.	0.5	32
663	Long-term Course of Hospitalization for Schizophrenia: Part II. Change With Passage of Time. Schizophrenia Bulletin, 1992, 18, 229-241.	2.3	82
664	Methimazole treatment reduces cardiac hypertrophy and mortality without a concomitant reduction in blood pressure in established Goldblatt two-kidney one clip hypertension. Experientia, 1992, 48, 755-758.	1.2	4
665	Psychiatric disorders in primary health care clinics one year after a major latin american disaster. Stress and Health, 1991, 7, 25-32.	0.7	7
666	Attention-Deficit Hyperactivity Disorder. , 0, , 195-208.		0

#	Article	IF	CITATIONS
667	The Effects of Trauma Type, Timing, Accumulation, and Sequencing. , 0, , 153-162.		O
668	Analysis of burden in caregivers of people with Alzheimer's disease using self-report and supervision hours. Journal of Nutrition, Health and Aging, 0, , .	1.5	9
669	The Use of Bayesian Hierarchical Methods to Assess Selection and Observer Bias in Observational Health Outcome Studies. SSRN Electronic Journal, 0, , .	0.4	O
670	Launay–Slade Hallucination Scale-Extended: simplifying its interpretation. Psychosis, 0, , 1-10.	0.4	0
671	Psychotic experiences among informal caregivers: findings from 48 low- and middle-income countries. Social Psychiatry and Psychiatric Epidemiology, 0, , .	1.6	0