

Yusuke Suzuki

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4348338/publications.pdf>

Version: 2024-02-01

122
papers

3,957
citations

147801

31
h-index

144013

57
g-index

124
all docs

124
docs citations

124
times ranked

3050
citing authors

#	ARTICLE	IF	CITATIONS
1	Gross hematuria after SARS-CoV-2 vaccination: questionnaire survey in Japan. <i>Clinical and Experimental Nephrology</i> , 2022, 26, 316-322.	1.6	18
2	IgA Nephropathy with Gross Hematuria Following COVID-19 mRNA Vaccination. <i>Internal Medicine</i> , 2022, 61, 1033-1037.	0.7	16
3	Negative-pressure wound therapy is effective for peritoneal dialysis catheter exit-site management in the early postoperative period. <i>Scientific Reports</i> , 2022, 12, 70.	3.3	2
4	Safety, Tolerability, Pharmacokinetics, and Pharmacodynamics of VIS649 (Sibeprenlimab), an APRIL-Neutralizing IgG2 Monoclonal Antibody, in Healthy Volunteers. <i>Kidney International Reports</i> , 2022, 7, 993-1003.	0.8	18
5	Association of cardiac autonomic neuropathy assessed by heart rate response during exercise with intradialytic hypotension and mortality in hemodialysis patients. <i>Kidney International</i> , 2022, 101, 1054-1062.	5.2	2
6	Progranulin and Its Receptor Predict Kidney Function Decline in Patients With Type 2 Diabetes. <i>Frontiers in Endocrinology</i> , 2022, 13, 849457.	3.5	5
7	Coronary Artery Bypass Grafting in Patients with Chronic Kidney Disease: Chronic Kidney Disease Has an Independent Adverse Effect on the Long-Term Outcome of Coronary Artery Bypass Grafting. <i>BioMed Research International</i> , 2022, 2022, 1-14.	1.9	1
8	Application of the International IgA Nephropathy Prediction Tool one or two years post-biopsy. <i>Kidney International</i> , 2022, 102, 160-172.	5.2	25
9	FC051: Atacicept Reduces Serum ANTI-GD-IGA1 Levels in IgAN Patients. <i>Nephrology Dialysis Transplantation</i> , 2022, 37, .	0.7	1
10	Galactose-Deficient IgA1 as a Candidate Urinary Marker of IgA Nephropathy. <i>Journal of Clinical Medicine</i> , 2022, 11, 3173.	2.4	6
11	Predictors of early remission of proteinuria in adult patients with minimal change disease: a retrospective cohort study. <i>Scientific Reports</i> , 2022, 12, .	3.3	3
12	Fractional excretion of tumor necrosis factor receptor ¹ and 2 in patients with type ² diabetes and normal renal function. <i>Journal of Diabetes Investigation</i> , 2021, 12, 382-389.	2.4	9
13	Renal pathological analysis using galactose-deficient IgA1-specific monoclonal antibody is a strong tool for differentiation of primary IgA nephropathy from secondary IgA nephropathy. <i>CEN Case Reports</i> , 2021, 10, 17-22.	0.9	11
14	Continuous extracorporeal treatments in a dialysis patient with COVID-19. <i>CEN Case Reports</i> , 2021, 10, 172-177.	0.9	7
15	Impact of the number of steroid pulses in tonsillectomy combined with steroid pulse therapy: a nationwide retrospective study in Japan. <i>Clinical and Experimental Nephrology</i> , 2021, 25, 19-27.	1.6	4
16	LCC18, a benzamide-linked small molecule, ameliorates IgA nephropathy in mice. <i>Journal of Pathology</i> , 2021, 253, 427-441.	4.5	5
17	Ischemic Stroke Induces Rapid Renal Oxidative Stress and Lipometabolic Change. <i>Juntendo Medical Journal</i> , 2021, 67, 39-45.	0.1	0
18	A case of hereditary angioedema due to C1-inhibitor deficiency with recurrent abdominal pain diagnosed 40 years after the occurrence of the initial symptom. <i>Clinical Journal of Gastroenterology</i> , 2021, 14, 1175-1179.	0.8	2

#	ARTICLE	IF	CITATIONS
19	Galactose-deficient IgA1 and nephritis-associated plasmin receptors as markers for IgA-dominant infection-related glomerulonephritis. <i>Medicine (United States)</i> , 2021, 100, e24460.	1.0	6
20	The Phenotypic Difference of IgA Nephropathy and its Race/Gender-dependent Molecular Mechanisms. <i>Kidney360</i> , 2021, 2, 1339-1348.	2.1	15
21	Utility of remission criteria for the renal prognosis of IgA nephropathy. <i>Clinical and Experimental Nephrology</i> , 2021, 25, 988-995.	1.6	3
22	Association between social frailty as well as early physical dysfunction and exercise intolerance among older patients receiving hemodialysis. <i>Geriatrics and Gerontology International</i> , 2021, 21, 664-669.	1.5	8
23	Association Between Kidney Function Decline and Baseline TNFR Levels or Change Ratio in TNFR by Febuxostat Chiefly in Non-diabetic CKD Patients With Asymptomatic Hyperuricemia. <i>Frontiers in Medicine</i> , 2021, 8, 634932.	2.6	5
24	Are there animal models of IgA nephropathy?. <i>Seminars in Immunopathology</i> , 2021, 43, 639-648.	6.1	10
25	COVID-19-induced acute renal tubular injury associated with elevation of serum inflammatory cytokine. <i>Clinical and Experimental Nephrology</i> , 2021, 25, 1240-1246.	1.6	16
26	Nasal-associated lymphoid tissue is the major induction site for nephritogenic IgA in murine IgA nephropathy. <i>Kidney International</i> , 2021, 100, 364-376.	5.2	25
27	Executive summary of the KDIGO 2021 Guideline for the Management of Glomerular Diseases. <i>Kidney International</i> , 2021, 100, 753-779.	5.2	325
28	Differential organ-specific inflammatory response to progranulin in high-fat diet-fed mice. <i>Scientific Reports</i> , 2021, 11, 1194.	3.3	7
29	Quantifying Duration of Proteinuria Remission and Association with Clinical Outcome in IgA Nephropathy. <i>Journal of the American Society of Nephrology: JASN</i> , 2021, 32, 436-447.	6.1	34
30	Effect of blood volume change related to intensity of intradialytic aerobic exercise on hemodialysis adequacy: a pilot study. <i>International Urology and Nephrology</i> , 2021, , 1.	1.4	1
31	TLR9 activation induces aberrant IgA glycosylation via APRIL- and IL-6-mediated pathways in IgA nephropathy. <i>Kidney International</i> , 2020, 97, 340-349.	5.2	78
32	Efficacy of endoscopic sinus surgery for eosinophilic chronic rhinosinusitis with asthma. <i>Allergology International</i> , 2020, 69, 144-145.	3.3	6
33	Thrombotic microangiopathy in dasatinib-treated patients with chronic myeloid leukemia. <i>Journal of Onco-Nephrology</i> , 2020, 4, 41-45.	0.6	1
34	P0350THE DURATION OF PROTEINURIA REMISSION AND CLINICAL OUTCOMES IN IGA NEPHROPATHY. <i>Nephrology Dialysis Transplantation</i> , 2020, 35, .	0.7	0
35	P0505COST ANALYSIS OF SCREENING FOR IGA NEPHROPATHY USING NOVEL BIOMARKERS. <i>Nephrology Dialysis Transplantation</i> , 2020, 35, .	0.7	0
36	Protective effects of DPP-4 inhibitor on podocyte injury in glomerular diseases. <i>BMC Nephrology</i> , 2020, 21, 402.	1.8	11

#	ARTICLE	IF	CITATIONS
37	Rapid progression to end-stage renal disease in a child with IgA-dominant infection-related glomerulonephritis associated with parvovirus B19. <i>CEN Case Reports</i> , 2020, 9, 423-430.	0.9	5
38	Emergent initiation of dialysis is related to an increase in both mortality and medical costs. <i>Scientific Reports</i> , 2020, 10, 19638.	3.3	10
39	IgA Nephropathy Benefits from Compound K Treatment by Inhibiting NF- κ B/NLRP3 Inflammasome and Enhancing Autophagy and SIRT1. <i>Journal of Immunology</i> , 2020, 205, 202-212.	0.8	22
40	Circulating Tumor Necrosis Factor Receptors: A Potential Biomarker for the Progression of Diabetic Kidney Disease. <i>International Journal of Molecular Sciences</i> , 2020, 21, 1957.	4.1	34
41	Incidence of remission and relapse of proteinuria, end-stage kidney disease, mortality, and major outcomes in primary nephrotic syndrome: the Japan Nephrotic Syndrome Cohort Study (JNSCS). <i>Clinical and Experimental Nephrology</i> , 2020, 24, 526-540.	1.6	33
42	Crucial Role of AIM/CD5L in the Development of Glomerular Inflammation in IgA Nephropathy. <i>Journal of the American Society of Nephrology: JASN</i> , 2020, 31, 2013-2024.	6.1	29
43	Better remission rates in elderly Japanese patients with primary membranous nephropathy in nationwide real-world practice: The Japan Nephrotic Syndrome Cohort Study (JNSCS). <i>Clinical and Experimental Nephrology</i> , 2020, 24, 893-909.	1.6	6
44	Lipopolysaccharide-Deficient <i>Acinetobacter baumannii</i> Due to Colistin Resistance Is Killed by Neutrophil-Produced Lysozyme. <i>Frontiers in Microbiology</i> , 2020, 11, 573.	3.5	26
45	Improving treatment decisions using personalized risk assessment from the International IgA Nephropathy Prediction Tool. <i>Kidney International</i> , 2020, 98, 1009-1019.	5.2	35
46	17. Novel Therapeutic Approaches Based on Recent Advance in Elucidation of Pathogenesis of IgA Nephropathy. <i>The Journal of the Japanese Society of Internal Medicine</i> , 2020, 109, 1976-1982.	0.0	0
47	Multi-point analysis of airborne Japanese cedar (<i>Cryptomeria japonica</i> D. Don) pollen by Pollen Robo and the relationship between pollen count and the severity of symptoms. <i>Aerobiologia</i> , 2019, 35, 635-646.	1.7	3
48	Galactose-deficient IgA1 in skin and serum from patients with skin-limited and systemic IgA vasculitis. <i>Journal of the American Academy of Dermatology</i> , 2019, 81, 1078-1085.	1.2	15
49	208. GALACTOSE DEFICIENT IGA1 (GD-IGA1) IN SKIN AND SERUM FROM PATIENTS WITH SKIN-LIMITED AND SYSTEMIC IGA VASCULITIS. <i>Rheumatology</i> , 2019, 58, .	1.9	2
50	Association Between Tonsillectomy and Outcomes in Patients With Immunoglobulin A Nephropathy. <i>JAMA Network Open</i> , 2019, 2, e194772.	5.9	59
51	Evaluating a New International Risk-Prediction Tool in IgA Nephropathy. <i>JAMA Internal Medicine</i> , 2019, 179, 942.	5.1	266
52	A Proliferation Inducing Ligand (APRIL) targeted antibody is a safe and effective treatment of murine IgA nephropathy. <i>Kidney International</i> , 2019, 96, 104-116.	5.2	41
53	Exercise-Induced Acute Kidney Injury in a Police Officer with Hereditary Renal Hypouricemia. <i>Case Reports in Nephrology and Dialysis</i> , 2019, 9, 92-101.	0.6	10
54	A grading system that predicts the risk of dialysis induction in IgA nephropathy patients based on the combination of the clinical and histological severity. <i>Clinical and Experimental Nephrology</i> , 2019, 23, 16-25.	1.6	18

#	ARTICLE	IF	CITATIONS
55	Regional variations in immunosuppressive therapy in patients with primary nephrotic syndrome: the Japan nephrotic syndrome cohort study. <i>Clinical and Experimental Nephrology</i> , 2018, 22, 1266-1280.	1.6	21
56	IgA nephropathy and IgA vasculitis with nephritis have a shared feature involving galactose-deficient IgA1-oriented pathogenesis. <i>Kidney International</i> , 2018, 93, 700-705.	5.2	151
57	Enrichment of airborne Japanese cedar (<i>Cryptomeria japonica</i>) pollen in mountain ranges when passing through a front accompanying temperate low pressure. <i>Aerobiologia</i> , 2018, 34, 105-110.	1.7	6
58	Chronic intermittent hypoxia-mediated renal sympathetic nerve activation in hypertension and cardiovascular disease. <i>Scientific Reports</i> , 2018, 8, 17926.	3.3	28
59	Galactose-Deficient IgA1-Specific Antibody Recognizes GalNAc-Modified Unique Epitope on Hinge Region of IgA1. <i>Monoclonal Antibodies in Immunodiagnosis and Immunotherapy</i> , 2018, 37, 252-256.	1.6	9
60	The Xanthine Oxidase Inhibitor Febuxostat Suppresses the Progression of IgA Nephropathy, Possibly via Its Anti-Inflammatory and Anti-Fibrotic Effects in the gddY Mouse Model. <i>International Journal of Molecular Sciences</i> , 2018, 19, 3967.	4.1	10
61	Association between circulating tumor necrosis factor-related biomarkers and estimated glomerular filtration rate in type 2 diabetes. <i>Scientific Reports</i> , 2018, 8, 15302.	3.3	30
62	Murine Models of Human IgA Nephropathy. <i>Seminars in Nephrology</i> , 2018, 38, 513-520.	1.6	23
63	Suffocation due to Acute Airway Edema in a Patient with Hereditary Angioedema Highlighted the Need for Urgent Improvements in Treatment Availability in Japan. <i>Internal Medicine</i> , 2018, 57, 3193-3197.	0.7	11
64	Clinical predictive biomarkers for normoalbuminuric diabetic kidney disease. <i>Diabetes Research and Clinical Practice</i> , 2018, 141, 62-68.	2.8	29
65	Expression of Cathepsin L and Its Intrinsic Inhibitors in Glomeruli of Rats With Puromycin Aminonucleoside Nephrosis. <i>Journal of Histochemistry and Cytochemistry</i> , 2018, 66, 863-877.	2.5	6
66	Establishment of a novel mouse xenograft model of human uterine leiomyoma. <i>Scientific Reports</i> , 2018, 8, 8872.	3.3	9
67	Serum galactose-deficient-IgA1 and IgG autoantibodies correlate in patients with IgA nephropathy. <i>PLoS ONE</i> , 2018, 13, e0190967.	2.5	56
68	High doses of antipsychotic polypharmacy are related to an increase in serum levels of pentosidine in patients with schizophrenia. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2017, 76, 42-48.	4.8	8
69	Evaluation of Long-Term Combination Therapy With Peritoneal Dialysis and Hemodialysis. <i>Therapeutic Apheresis and Dialysis</i> , 2017, 21, 180-184.	0.9	19
70	Circulating TNF Receptors 1 and 2 Predict Mortality in Patients with End-stage Renal Disease Undergoing Dialysis. <i>Scientific Reports</i> , 2017, 7, 43520.	3.3	49
71	High serum soluble tumor necrosis factor receptor 1 predicts poor treatment response in acute-stage schizophrenia. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2017, 76, 145-154.	4.8	11
72	Toll-Like Receptor 9 Stimulation Induces Aberrant Expression of a Proliferation-Inducing Ligand by Tonsillar Germinal Center B Cells in IgA Nephropathy. <i>Journal of the American Society of Nephrology: JASN</i> , 2017, 28, 1227-1238.	6.1	91

#	ARTICLE	IF	CITATIONS
73	Inhibition of STAT3 Signaling Reduces IgA1 Autoantigen Production in IgA Nephropathy. <i>Kidney International Reports</i> , 2017, 2, 1194-1207.	0.8	49
74	Galactose-Deficient IgA1 as a Candidate Urinary Polypeptide Marker of IgA Nephropathy?. <i>Disease Markers</i> , 2016, 2016, 1-6.	1.3	32
75	Apoptosis inhibitor of macrophage protein enhances intraluminal debris clearance and ameliorates acute kidney injury in mice. <i>Nature Medicine</i> , 2016, 22, 183-193.	30.7	161
76	Beyond the Differences in Tonsillectomy in IgA Nephropathy: From Rationale To Indications in Patients. <i>PLoS ONE</i> , 2016, 11, 311-319.		0
77	Circulating TNF Receptors 1 and 2 Are Associated with the Severity of Renal Interstitial Fibrosis in IgA Nephropathy. <i>PLoS ONE</i> , 2015, 10, e0122212.	2.5	48
78	Pathogenic Role of a Proliferation-Inducing Ligand (APRIL) in Murine IgA Nephropathy. <i>PLoS ONE</i> , 2015, 10, e0137044.	2.5	24
79	FP304 ABERRANT APRIL EXPRESSION IN TONSILLAR GERMINAL CENTER B CELLS IN IGA NEPHROPATHY PATIENTS. <i>Nephrology Dialysis Transplantation</i> , 2015, 30, iii168-iii169.	0.7	0
80	Paradigm shift in activity assessment of IgA nephropathy – optimizing the next generation of diagnostic and therapeutic maneuvers via glycan targeting. <i>Expert Opinion on Biological Therapy</i> , 2015, 15, 583-593.	3.1	9
81	TWEAK/Fn14 system and crescent formation in IgA nephropathy. <i>BMC Nephrology</i> , 2015, 16, 27.	1.8	15
82	Altered serum glyceraldehyde-derived advanced glycation end product (AGE) and soluble AGE receptor levels indicate carbonyl stress in patients with schizophrenia. <i>Neuroscience Letters</i> , 2015, 593, 51-55.	2.1	19
83	Novel lectin-independent approach to detect galactose-deficient IgA1 in IgA nephropathy. <i>Nephrology Dialysis Transplantation</i> , 2015, 30, 1315-1321.	0.7	99
84	Phenotype of asthma related with high serum periostin levels. <i>Allergy International</i> , 2015, 64, 175-180.	3.3	102
85	Changes in Nephritogenic Serum Galactose-Deficient IgA1 in IgA Nephropathy following Tonsillectomy and Steroid Therapy. <i>PLoS ONE</i> , 2014, 9, e89707.	2.5	72
86	Dietary Zinc Is a Key Environmental Modifier in the Progression of IgA Nephropathy. <i>PLoS ONE</i> , 2014, 9, e90558.	2.5	19
87	Uncoupling of Glomerular IgA Deposition and Disease Progression in Alymphoplasia Mice with IgA Nephropathy. <i>PLoS ONE</i> , 2014, 9, e95365.	2.5	6
88	A Panel of Serum Biomarkers Differentiates IgA Nephropathy from Other Renal Diseases. <i>PLoS ONE</i> , 2014, 9, e98081.	2.5	93
89	Diagnosis and activity assessment of immunoglobulin A nephropathy: current perspectives on noninvasive testing with aberrantly glycosylated immunoglobulin A-related biomarkers. <i>International Journal of Nephrology and Renovascular Disease</i> , 2014, 7, 409.	1.8	17
90	Carbon Ion Beam Radiotherapy for Sinonasal Malignant Tumors Invading Skull Base. <i>Case Reports in Otolaryngology</i> , 2014, 2014, 1-4.	0.2	2

#	ARTICLE	IF	CITATIONS
91	Serum levels of galactose-deficient immunoglobulin (Ig) A1 and related immune complex are associated with disease activity of IgA nephropathy. <i>Clinical and Experimental Nephrology</i> , 2014, 18, 770-777.	1.6	59
92	Proposal of remission criteria for IgA nephropathy. <i>Clinical and Experimental Nephrology</i> , 2014, 18, 481-486.	1.6	38
93	Development of animal models of human IgA nephropathy. <i>Drug Discovery Today: Disease Models</i> , 2014, 11, 5-11.	1.2	24
94	Serum under-O-glycosylated IgA1 level is not correlated with glomerular IgA deposition based upon heterogeneity in the composition of immune complexes in IgA nephropathy. <i>BMC Nephrology</i> , 2014, 15, 89.	1.8	11
95	Expressions and Roles of Periostin in Otolaryngological Diseases. <i>Allergology International</i> , 2014, 63, 171-180.	3.3	49
96	The Kinetics of Glomerular Deposition of Nephritogenic IgA. <i>PLoS ONE</i> , 2014, 9, e113005.	2.5	13
97	Nationwide survey on current treatments for IgA nephropathy in Japan. <i>Clinical and Experimental Nephrology</i> , 2013, 17, 827-833.	1.6	48
98	Experimental evidence of cell dissemination playing a role in pathogenesis of IgA nephropathy in multiple lymphoid organs. <i>Nephrology Dialysis Transplantation</i> , 2013, 28, 320-326.	0.7	12
99	Amelioration of Angiotensin II-Induced Salt-Sensitive Hypertension by Liver-Type Fatty Acid-Binding Protein in Proximal Tubules. <i>Hypertension</i> , 2013, 62, 712-718.	2.7	9
100	A histologic classification of IgA nephropathy for predicting long-term prognosis: emphasis on end-stage renal disease. <i>Journal of Nephrology</i> , 2013, 26, 350-357.	2.0	88
101	Tonsillar TLR9 expression and efficacy of tonsillectomy with steroid pulse therapy in IgA nephropathy patients. <i>Nephrology Dialysis Transplantation</i> , 2012, 27, 1090-1097.	0.7	55
102	Development of a Model of Early-Onset IgA Nephropathy. <i>Journal of the American Society of Nephrology: JASN</i> , 2012, 23, 1364-1374.	6.1	51
103	Determination of Severity of Murine IgA Nephropathy by Glomerular Complement Activation by Aberrantly Glycosylated IgA and Immune Complexes. <i>American Journal of Pathology</i> , 2012, 181, 1338-1347.	3.8	42
104	Aberrant Glycosylation of IgA1 and Anti-Glycan Antibodies in IgA Nephropathy: Role of Mucosal Immune System. <i>Advances in Oto-Rhino-Laryngology</i> , 2011, 72, 60-63.	1.6	40
105	Different Pathological Roles of Toll-Like Receptor 9 on Mucosal B Cells and Dendritic Cells in Murine IgA Nephropathy. <i>Clinical and Developmental Immunology</i> , 2011, 2011, 1-10.	3.3	26
106	Reevaluation of the Mucosa-Bone Marrow Axis in IgA Nephropathy with Animal Models. <i>Advances in Oto-Rhino-Laryngology</i> , 2011, 72, 64-67.	1.6	12
107	Pathological Role of Tonsillar B Cells in IgA Nephropathy. <i>Clinical and Developmental Immunology</i> , 2011, 2011, 1-8.	3.3	34
108	A Case of Cerebellar Ataxia with EB Virus Infection. <i>Practica Otologica, Supplement</i> , 2011, 130, 124-130.	0.0	1

#	ARTICLE	IF	CITATIONS
109	A Case of Piriform Sinus Fistula Occurring Seven Times before Resolution. <i>Practica Otologica, Supplement</i> , 2011, 130, 106-111.	0.0	0
110	Prophylactic Effect and QOL in Cases of Japanese Cedar Pollinosis. <i>Practica Otologica, Supplement</i> , 2011, 130, 44-48.	0.0	0
111	Two Cases of Maxillary Sinus Hematocele Mimicking Malignant Tumors. <i>Practica Otologica, Supplement</i> , 2011, 130, 49-55.	0.0	0
112	Clinical Study of 80 Parotid Gland Tumors. <i>Practica Otologica, Supplement</i> , 2011, 130, 146-151.	0.0	0
113	A Case of Papillary Carcinoma from a Thyroglossal Duct Remnant. <i>Practica Otologica, Supplement</i> , 2011, 130, 184-189.	0.0	0
114	Aberrantly glycosylated IgA1 in IgA nephropathy patients is recognized by IgG antibodies with restricted heterogeneity. <i>Journal of Clinical Investigation</i> , 2009, 119, 1668-77.	8.2	356
115	Amelioration of Crescentic Glomerulonephritis by RhoA Kinase Inhibitor, Fasudil, through Podocyte Protection and Prevention of Leukocyte Migration. <i>American Journal of Pathology</i> , 2008, 172, 603-614.	3.8	35
116	Potential Immunopathogenic Role of the Mucosa-Bone Marrow Axis in IgA Nephropathy: Insights From Animal Models. <i>Seminars in Nephrology</i> , 2008, 28, 66-77.	1.6	43
117	Toll-Like Receptor 9 Affects Severity of IgA Nephropathy. <i>Journal of the American Society of Nephrology: JASN</i> , 2008, 19, 2384-2395.	6.1	160
118	Roles of Bone Marrow, Mucosa and Lymphoid Tissues in Pathogenesis of Murine IgA Nephropathy. , 2007, 157, 164-168.		9
119	The Mucosa-Bone-Marrow Axis in IgA Nephropathy. , 2007, 157, 70-79.		29
120	Th2 cytokine induces aberrant O-glycosylation in the hinge region of IgA1 via downregulation of core1 β , 3-galactosyltransferase and its molecular chaperone Cosmc. <i>Juntendo J, Igaku</i> , 2007, 53, 113-120.	0.1	0
121	Genome-Wide Scan in a Novel IgA Nephropathy Model Identifies a Susceptibility Locus on Murine Chromosome 10, in a Region Syntenic to Human IGAN1 on Chromosome 6q22-23. <i>Journal of the American Society of Nephrology: JASN</i> , 2005, 16, 1289-1299.	6.1	67
122	Human bronchial smooth muscle cell proliferation via thromboxane A2 receptor. <i>Prostaglandins Leukotrienes and Essential Fatty Acids</i> , 2004, 71, 375-382.	2.2	11