

# Khawla S Al-Kuraya

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4371462/publications.pdf>

Version: 2024-02-01

77  
papers

1,388  
citations

394421

19  
h-index

377865

34  
g-index

77  
all docs

77  
docs citations

77  
times ranked

2350  
citing authors

#	ARTICLE	IF	CITATIONS
1	Clinicopathological Analysis of Papillary Thyroid Cancer with <i>PIK3CA</i> Alterations in a Middle Eastern Population. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2008, 93, 611-618.	3.6	169
2	Cross-Talk between NF $\kappa$ B and the PI3-Kinase/AKT Pathway Can Be Targeted in Primary Effusion Lymphoma (PEL) Cell Lines for Efficient Apoptosis. <i>PLoS ONE</i> , 2012, 7, e39945.	2.5	138
3	Fatty Acid Synthase and AKT Pathway Signaling in a Subset of Papillary Thyroid Cancers. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2008, 93, 4088-4097.	3.6	65
4	FoxM1 and Its Association with Matrix Metalloproteinases (MMP) Signaling Pathway in Papillary Thyroid Carcinoma. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2012, 97, E1-E13.	3.6	56
5	Genetic heterogeneity and evolutionary history of high-grade ovarian carcinoma and matched distant metastases. <i>British Journal of Cancer</i> , 2020, 122, 1219-1230.	6.4	56
6	Identification of novel <i>BRCA</i> founder mutations in Middle Eastern breast cancer patients using capture and Sanger sequencing analysis. <i>International Journal of Cancer</i> , 2016, 139, 1091-1097.	5.1	52
7	Telomerase reverse transcriptase mutations are independent predictor of disease-free survival in Middle Eastern papillary thyroid cancer. <i>International Journal of Cancer</i> , 2018, 142, 2028-2039.	5.1	50
8	Loss of PTEN expression is associated with aggressive behavior and poor prognosis in Middle Eastern triple-negative breast cancer. <i>Breast Cancer Research and Treatment</i> , 2015, 151, 541-553.	2.5	43
9	Genomic Profiling of Thyroid Cancer Reveals a Role for Thyroglobulin in Metastasis. <i>American Journal of Human Genetics</i> , 2016, 98, 1170-1180.	6.2	41
10	High prevalence of mTOR complex activity can be targeted using Torin2 in papillary thyroid carcinoma. <i>Carcinogenesis</i> , 2014, 35, 1564-1572.	2.8	40
11	XIAP over-expression is an independent poor prognostic marker in Middle Eastern breast cancer and can be targeted to induce efficient apoptosis. <i>BMC Cancer</i> , 2017, 17, 640.	2.6	39
12	Prevalence of Lynch syndrome in a Middle Eastern population with colorectal cancer. <i>Cancer</i> , 2015, 121, 1762-1771.	4.1	34
13	Role of X-Linked Inhibitor of Apoptosis as a Prognostic Marker and Therapeutic Target in Papillary Thyroid Carcinoma. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2015, 100, E974-E985.	3.6	34
14	Co-targeting of Cyclooxygenase-2 and FoxM1 is a viable strategy in inducing anticancer effects in colorectal cancer cells. <i>Molecular Cancer</i> , 2015, 14, 131.	19.2	33
15	A very low incidence of BRAF mutations in Middle Eastern colorectal carcinoma. <i>Molecular Cancer</i> , 2014, 13, 168.	19.2	31
16	Whole-Exome Sequencing of Matched Primary and Metastatic Papillary Thyroid Cancer. <i>Thyroid</i> , 2020, 30, 42-56.	4.5	31
17	ALK alteration is a frequent event in aggressive breast cancers. <i>Breast Cancer Research</i> , 2015, 17, 127.	5.0	29
18	Expanding the spectrum of germline variants in cancer. <i>Human Genetics</i> , 2017, 136, 1431-1444.	3.8	23

#	ARTICLE	IF	CITATIONS
19	Evolution and Impact of Subclonal Mutations in Papillary Thyroid Cancer. American Journal of Human Genetics, 2019, 105, 959-973.	6.2	22
20	Differential expression of PD-L1 between primary and metastatic epithelial ovarian cancer and its clinico-pathological correlation. Scientific Reports, 2021, 11, 3750.	3.3	22
21	TGF $\beta$ -induced SMAD4-dependent Apoptosis Proceeded by EMT in CRC. Molecular Cancer Therapeutics, 2019, 18, 1312-1322.	4.1	20
22	Molecular markers and pathway analysis of colorectal carcinoma in the Middle East. Cancer, 2015, 121, 3799-3808.	4.1	19
23	FoxM1 and $\beta$ -catenin predicts aggressiveness in Middle Eastern ovarian cancer and their co-targeting impairs the growth of ovarian cancer cells. Oncotarget, 2018, 9, 3590-3604.	1.8	18
24	PD-L1 Is an Independent Prognostic Marker in Middle Eastern PTC and Its Expression Is Upregulated by BRAFV600E Mutation. Cancers, 2021, 13, 555.	3.7	18
25	Prognostic significance of DNMT3A alterations in Middle Eastern papillary thyroid carcinoma. European Journal of Cancer, 2019, 117, 133-144.	2.8	17
26	Overexpression of PARP is an independent prognostic marker for poor survival in Middle Eastern breast cancer and its inhibition can be enhanced with embelin co-treatment. Oncotarget, 2018, 9, 37319-37332.	1.8	17
27	APC truncating mutations in Middle Eastern Population: Tankyrase inhibitor is an effective strategy to sensitize APC mutant CRC To 5-FU chemotherapy. Biomedicine and Pharmacotherapy, 2020, 121, 109572.	5.6	15
28	FoxM1 is an independent poor prognostic marker and therapeutic target for advanced Middle Eastern breast cancer. Oncotarget, 2018, 9, 17466-17482.	1.8	15
29	Annual Hazard Rate of Recurrence in Middle Eastern Papillary Thyroid Cancer over a Long-Term Follow-Up. Cancers, 2020, 12, 3624.	3.7	13
30	<i>MED12</i> is recurrently mutated in Middle Eastern colorectal cancer. Gut, 2018, 67, gutjnl-2016-313334.	12.1	12
31	Germline POLE and POLD1 proofreading domain mutations in endometrial carcinoma from Middle Eastern region. Cancer Cell International, 2019, 19, 334.	4.1	12
32	Prevalence, spectrum, and founder effect of BRCA1 and BRCA2 mutations in epithelial ovarian cancer from the Middle East. Human Mutation, 2019, 40, 729-733.	2.5	11
33	TERT Promoter Mutations Are an Independent Predictor of Distant Metastasis in Middle Eastern Papillary Thyroid Microcarcinoma. Frontiers in Endocrinology, 2022, 13, 808298.	3.5	11
34	Dual Targeting of mTOR Activity with Torin2 Potentiates Anticancer Effects of Cisplatin in Epithelial Ovarian Cancer. Molecular Medicine, 2015, 21, 466-478.	4.4	10
35	Downregulation of SKP2 in Papillary Thyroid Cancer Acts Synergistically With TRAIL on Inducing Apoptosis via ROS. Journal of Clinical Endocrinology and Metabolism, 2018, 103, 1530-1544.	3.6	10
36	Telomerase reverse transcriptase promoter mutations in cancers derived from multiple organ sites among middle eastern population. Genomics, 2020, 112, 1746-1753.	2.9	10

#	ARTICLE	IF	CITATIONS
37	NTRK fusion analysis reveals enrichment in Middle Eastern BRAF wild-type PTC. <i>European Journal of Endocrinology</i> , 2021, 184, 503-511.	3.7	10
38	Recurrent Somatic MAP2K1 Mutations in Papillary Thyroid Cancer and Colorectal Cancer. <i>Frontiers in Oncology</i> , 2021, 11, 670423.	2.8	10
39	Colorectal carcinoma from Saudi Arabia. Analysis of MLH-1, MSH-2 and p53 genes by immunohistochemistry and tissue microarray analysis. <i>Journal of King Abdulaziz University, Islamic Economics</i> , 2006, 27, 323-8.	1.1	10
40	Clonal Evolution and Timing of Metastatic Colorectal Cancer. <i>Cancers</i> , 2020, 12, 2938.	3.7	9
41	POLE and POLD1 germline exonuclease domain pathogenic variants, a rare event in colorectal cancer from the Middle East. <i>Molecular Genetics &amp; Genomic Medicine</i> , 2020, 8, e1368.	1.2	9
42	Microscopic Extrathyroidal Extension Results in Increased Rate of Tumor Recurrence and Is an Independent Predictor of Patient's Outcome in Middle Eastern Papillary Thyroid Carcinoma. <i>Frontiers in Oncology</i> , 2021, 11, 724432.	2.8	9
43	PD-L1 Protein Expression in Middle Eastern Breast Cancer Predicts Favorable Outcome in Triple-Negative Breast Cancer. <i>Cells</i> , 2021, 10, 229.	4.1	8
44	Krupple-Like Factor 5 is a Potential Therapeutic Target and Prognostic Marker in Epithelial Ovarian Cancer. <i>Frontiers in Pharmacology</i> , 2020, 11, 598880.	3.5	7
45	PD-L1 Expression Is Associated with Deficient Mismatch Repair and Poor Prognosis in Middle Eastern Colorectal Cancers. <i>Journal of Personalized Medicine</i> , 2021, 11, 73.	2.5	7
46	High Expression of Cyclin D1 is an Independent Marker for Favorable Prognosis in Middle Eastern Breast Cancer. <i>OncoTargets and Therapy</i> , 2021, Volume 14, 3309-3318.	2.0	7
47	POLE and POLD1 pathogenic variants in the proofreading domain in papillary thyroid cancer. <i>Endocrine Connections</i> , 2020, 9, 923-932.	1.9	7
48	Lymph node ratio is superior to AJCC N stage for predicting recurrence in papillary thyroid carcinoma. <i>Endocrine Connections</i> , 2022, 11, .	1.9	7
49	Loss of ZNF677 Expression Is an Independent Predictor for Distant Metastasis in Middle Eastern Papillary Thyroid Carcinoma Patients. <i>International Journal of Molecular Sciences</i> , 2021, 22, 7833.	4.1	6
50	Risk Factors for Central Lymph Node Metastases and Benefit of Prophylactic Central Lymph Node Dissection in Middle Eastern Patients With cN0 Papillary Thyroid Carcinoma. <i>Frontiers in Oncology</i> , 2021, 11, 819824.	2.8	6
51	CHD4 Predicts Aggressiveness in PTC Patients and Promotes Cancer Stemness and EMT in PTC Cells. <i>International Journal of Molecular Sciences</i> , 2021, 22, 504.	4.1	5
52	Prognostic Value and Function of KLF5 in Papillary Thyroid Cancer. <i>Cancers</i> , 2021, 13, 185.	3.7	5
53	Cyclin-dependent kinase 9 (CDK9) predicts recurrence in Middle Eastern epithelial ovarian cancer. <i>Journal of Ovarian Research</i> , 2021, 14, 69.	3.0	5
54	Male Sex Is an Independent Predictor of Recurrence-Free Survival in Middle Eastern Papillary Thyroid Carcinoma. <i>Frontiers in Endocrinology</i> , 2022, 13, 777345.	3.5	5

#	ARTICLE	IF	CITATIONS
55	Prognostic Significance of COX-2 Overexpression in BRAF-Mutated Middle Eastern Papillary Thyroid Carcinoma. <i>International Journal of Molecular Sciences</i> , 2020, 21, 9498.	4.1	4
56	Loss of ZNF677 expression is a predictive biomarker for lymph node metastasis in Middle Eastern Colorectal Cancer. <i>Scientific Reports</i> , 2021, 11, 22346.	3.3	4
57	APOBEC SBS13 Mutational Signatureâ€™A Novel Predictor of Radioactive Iodine Refractory Papillary Thyroid Carcinoma. <i>Cancers</i> , 2022, 14, 1584.	3.7	4
58	The study of Lynch syndrome in a special population reveals a strong founder effect and an unusual mutational mechanism in familial adenomatous polyposis. <i>Gut</i> , 2020, 69, 2048-2049.	12.1	3
59	Zamzam water protects cancer cells from chemotherapy-induced apoptosis via mitogen-activated protein kinase-dependent pathway. <i>Biomedicine and Pharmacotherapy</i> , 2019, 118, 109376.	5.6	2
60	PD-L1 Expression Is an Independent Marker for Lymph Node Metastasis in Middle Eastern Endometrial Cancer. <i>Diagnostics</i> , 2021, 11, 394.	2.6	2
61	Role of Phosphatidylinositol 3- Kinase/AKT Pathway in Diffuse Large B-Cell Lymphoma Survival.. <i>Blood</i> , 2005, 106, 4808-4808.	1.4	1
62	Reply to specific gene patterns and molecular pathways related to human carcinogenesis in different populations among various geographic locations. <i>Cancer</i> , 2016, 122, 1135-1137.	4.1	0
63	Response to Yehia etÂal.. <i>American Journal of Human Genetics</i> , 2017, 100, 564-565.	6.2	0
64	Abstract 1176: CHD4 predicts aggressiveness in PTC patients and promotes cancer stemness and EMT in PTC cells. , 2021, , .		0
65	Abstract 730: Annual hazard rate of recurrence in Middle-Eastern papillary thyroid cancer over a long-term follow-up. , 2021, , .		0
66	Abstract 3190: PD-L1 protein expression in Middle Eastern breast cancer predicts favorable outcome in triple negative breast cancer. , 2021, , .		0
67	Abstract 3186: Differential expression of PD-L1 between primary and metastatic Middle-Eastern epithelial ovarian carcinoma and its clinico-pathological correlation. , 2021, , .		0
68	Abstract 1963: High expression of Cyclin D1 is an independent marker for favorable prognosis in Middle Eastern breast cancer. , 2021, , .		0
69	Abstract 2425: Prevalence of NTRK fusions and clinico-pathological characteristics of Middle Eastern papillary thyroid cancer revealed enrichment in BRAF wild-type PTC. , 2021, , .		0
70	Prognostic Molecular Features in Diffuse Large B-Cell Lymphoma from Saudi Arabia.. <i>Blood</i> , 2004, 104, 4609-4609.	1.4	0
71	Epstein-Barr Virus Infection Is Not the Sole Cause of High Prevalence for Hodgkinâ€™s Lymphoma in Saudi Arabia.. <i>Blood</i> , 2004, 104, 3120-3120.	1.4	0
72	Apigenin Induces Apoptosis in Primary Effusion Lymphoma.. <i>Blood</i> , 2005, 106, 4807-4807.	1.4	0

#	ARTICLE	IF	CITATIONS
73	Proteasome Inhibitor Treatment Down-Regulates S-Phase Kinase-Associated Protein 2 Causing Inhibition of Proliferation and Induces Apoptosis in Primary Effusion Lymphoma.. Blood, 2006, 108, 4616-4616.	1.4	0
74	Significance of Ubiquitin Ligase Subunit SKP-2 Proto-Oncogene and Proliferative Marker Ki67 Expression in Diffuse Large B-Cell Lymphoma.. Blood, 2006, 108, 4343-4343.	1.4	0
75	SUN-132 KLF5 Is a Poor Prognostic Marker and Therapeutic Target for Middle Eastern Papillary Thyroid Carcinoma. Journal of the Endocrine Society, 2020, 4, .	0.2	0
76	Genetic risk of cancer: a tale of diversity from the Middle East. Lancet Oncology, The, 2022, , .	10.7	0
77	Wingless signaling pathway family relation to colon cancer. Have we come full circle?. Journal of King Abdulaziz University, Islamic Economics, 2005, 26, 19-23.	1.1	0