

Fred Naider

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4409437/publications.pdf>

Version: 2024-02-01

96
papers

2,928
citations

172457

29
h-index

189892

50
g-index

96
all docs

96
docs citations

96
times ranked

2246
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 1 | Proline-Dependent Structural and Biological Properties of Peptides and Proteins. <i>Critical Reviews in Biochemistry and Molecular Biology</i> , 1993, 28, 31-81. | 5.2 | 511 |
| 2 | The PTR family: a new group of peptide transporters. <i>Molecular Microbiology</i> , 1995, 16, 825-834. | 2.5 | 238 |
| 3 | An oligopeptide transport gene from <i>Candida albicans</i> . <i>Microbiology (United Kingdom)</i> , 1997, 143, 387-396. | 1.8 | 96 |
| 4 | Peptides in the treatment of AIDS. <i>Current Opinion in Structural Biology</i> , 2009, 19, 473-482. | 5.7 | 95 |
| 5 | The $\hat{\pm}$ -factor mating pheromone of <i>Saccharomyces cerevisiae</i> : a model for studying the interaction of peptide hormones and G protein-coupled receptors. <i>Peptides</i> , 2004, 25, 1441-1463. | 2.4 | 89 |
| 6 | Multiplicity and regulation of genes encoding peptide transporters in <i>Saccharomyces cerevisiae</i> . <i>Molecular Membrane Biology</i> , 2001, 18, 105-112. | 2.0 | 77 |
| 7 | <i>Schizosaccharomyces pombe</i> <i>isp4</i> encodes a transporter representing a novel family of oligopeptide transporters. <i>Molecular Microbiology</i> , 2002, 28, 729-741. | 2.5 | 77 |
| 8 | A recognition component of the ubiquitin system is required for peptide transport in <i>Saccharomyces cerevisiae</i> . <i>Molecular Microbiology</i> , 1995, 15, 225-234. | 2.5 | 62 |
| 9 | A Limited Spectrum of Mutations Causes Constitutive Activation of the Yeast $\hat{\pm}$ -Factor Receptor. <i>Biochemistry</i> , 2000, 39, 6898-6909. | 2.5 | 58 |
| 10 | Oligomerization of the Yeast $\hat{\pm}$ -Factor Receptor. <i>Journal of Biological Chemistry</i> , 2006, 281, 20698-20714. | 3.4 | 54 |
| 11 | Identification of Ligand Binding Regions of the <i>Saccharomyces cerevisiae</i> $\hat{\pm}$ -Factor Pheromone Receptor by Photoaffinity Cross-Linking. <i>Biochemistry</i> , 2004, 43, 13193-13203. | 2.5 | 48 |
| 12 | Unnatural Amino Acid Replacement in a Yeast G Protein-Coupled Receptor in Its Native Environment. <i>Biochemistry</i> , 2008, 47, 5638-5648. | 2.5 | 47 |
| 13 | Structure-Activity Relationships of the Yeast $\hat{\pm}$ -Facto. <i>Critical Reviews in Biochemistry</i> , 1986, 21, 225-248. | 7.5 | 46 |
| 14 | A Fluorescent $\hat{\pm}$ -Factor Analogue Exhibits Multiple Steps on Binding to Its G Protein Coupled Receptor in Yeast. <i>Biochemistry</i> , 2004, 43, 13564-13578. | 2.5 | 45 |
| 15 | Antagonistic and synergistic peptide analogs of the tridecapeptide mating pheromone of <i>Saccharomyces cerevisiae</i> . <i>Biochemistry</i> , 1992, 31, 551-557. | 2.5 | 43 |
| 16 | PTR3, a novel gene mediating amino acid-inducible regulation of peptide transport in <i>Saccharomyces cerevisiae</i> . <i>Molecular Microbiology</i> , 1998, 29, 297-310. | 2.5 | 40 |
| 17 | The First Extracellular Loop of the <i>Saccharomyces cerevisiae</i> G Protein-coupled Receptor Ste2p Undergoes a Conformational Change upon Ligand Binding. <i>Journal of Biological Chemistry</i> , 2007, 282, 10387-10397. | 3.4 | 40 |
| 18 | Cloning of a <i>Candida albicans</i> peptide transport gene. <i>Microbiology (United Kingdom)</i> , 1995, 141, 1147-1156. | 1.8 | 39 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 19 | Identification of Residues of the <i>Saccharomyces cerevisiae</i> G Protein-coupled Receptor Contributing to $\hat{\pm}$ -Factor Pheromone Binding. <i>Journal of Biological Chemistry</i> , 2001, 276, 37950-37961. | 3.4 | 39 |
| 20 | ATR-FTIR Study of the Structure and Orientation of Transmembrane Domains of the <i>Saccharomyces cerevisiae</i> $\hat{\pm}$ -Mating Factor Receptor in Phospholipids. <i>Biochemistry</i> , 2001, 40, 8945-8954. | 2.5 | 38 |
| 21 | Identification of a Contact Region between the Tridecapeptide $\hat{\pm}$ -Factor Mating Pheromone of <i>Saccharomyces cerevisiae</i> and Its G Protein-Coupled Receptor by Photoaffinity Labeling. <i>Biochemistry</i> , 2002, 41, 6128-6139. | 2.5 | 38 |
| 22 | Biosynthesis and NMR Analysis of a 73-Residue Domain of a <i>Saccharomyces cerevisiae</i> G Protein-Coupled Receptor. <i>Biochemistry</i> , 2005, 44, 11795-11810. | 2.5 | 36 |
| 23 | Nutrient regulation of oligopeptide transport in <i>Saccharomyces cerevisiae</i> . <i>Microbiology (United Kingdom)</i> 151 1078-1084 | 1.8 | 36 |
| 24 | The solution structure of monomeric $\langle \text{CCL} \rangle_5$ in complex with a doubly sulfated N-terminal segment of $\langle \text{CCR} \rangle_5$. <i>FEBS Journal</i> , 2018, 285, 1988-2003. | 4.7 | 35 |
| 25 | Structure of a Double Transmembrane Fragment of a G-Protein-Coupled Receptor in Micelles. <i>Biophysical Journal</i> , 2009, 96, 3187-3196. | 0.5 | 32 |
| 26 | Residues in the First Extracellular Loop of a G Protein-coupled Receptor Play a Role in Signal Transduction. <i>Journal of Biological Chemistry</i> , 2002, 277, 30581-30590. | 3.4 | 31 |
| 27 | High resolution NMR analysis of the seven transmembrane domains of a heptahelical receptor in organic-aqueous medium. <i>Biopolymers</i> , 2002, 64, 161-176. | 2.4 | 30 |
| 28 | NMR Studies in Dodecylphosphocholine of a Fragment Containing the Seventh Transmembrane Helix of a G-Protein-Coupled Receptor from <i>Saccharomyces cerevisiae</i> . <i>Biophysical Journal</i> , 2007, 93, 467-482. | 0.5 | 30 |
| 29 | Solution phase synthesis of <i>Saccharomyces cerevisiae</i> $\hat{\pm}$ -mating factor and its analogs. <i>International Journal of Peptide and Protein Research</i> , 1990, 36, 362-373. | 0.1 | 29 |
| 30 | Identification of Specific Transmembrane Residues and Ligand-Induced Interface Changes Involved In Homo-Dimer Formation of a Yeast G Protein-Coupled Receptor. <i>Biochemistry</i> , 2009, 48, 10976-10987. | 2.5 | 29 |
| 31 | Direct observation of cell wall glucans in whole cells of <i>Saccharomyces cerevisiae</i> by magic-angle spinning ^{13}C -nmr. <i>Biopolymers</i> , 1994, 34, 1627-1635. | 2.4 | 28 |
| 32 | Biological activity and conformational isomerism in position 9 analogs of the des-1-tryptophan, 3-beta-cyclohexylalanine-alpha-factor from <i>Saccharomyces cerevisiae</i> . <i>Biochemistry</i> , 1985, 24, 7070-7076. | 2.5 | 27 |
| 33 | Structure of segments of a G protein-coupled receptor: CD and NMR analysis of the <i>Saccharomyces cerevisiae</i> tridecapeptide pheromone receptor. <i>Biochemistry</i> , 1998, 37, 343-357. | | 27 |
| 34 | Tyr266 in the Sixth Transmembrane Domain of the Yeast $\hat{\pm}$ -Factor Receptor Plays Key Roles in Receptor Activation and Ligand Specificity. <i>Biochemistry</i> , 2002, 41, 13681-13689. | 2.5 | 27 |
| 35 | Synthetic peptides as probes for conformational preferences of domains of membrane receptors. <i>Biopolymers</i> , 2005, 80, 199-213. | 2.4 | 27 |
| 36 | THE PREFERRED CONFORMATIONS OF PROTECTED HOMODI- AND HOMOHEPTAMETHIONINE PEPTIDES A ^1H N.M.R. Study in Deuteriochloroform Medium. <i>International Journal of Peptide and Protein Research</i> , 1979, 14, 414-436. | 0.1 | 27 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 37 | Chemical synthesis of the M-factor mating pheromone from <i>Schizosaccharomyces pombe</i> . <i>Yeast</i> , 1994, 10, 595-601. | 1.7 | 26 |
| 38 | Cross-Linking of a DOPA-Containing Peptide Ligand into Its G Protein-Coupled Receptor. <i>Biochemistry</i> , 2009, 48, 2033-2044. | 2.5 | 25 |
| 39 | Identification of Residue-to-residue Contact between a Peptide Ligand and Its G Protein-coupled Receptor Using Periodate-mediated Dihydroxyphenylalanine Cross-linking and Mass Spectrometry. <i>Journal of Biological Chemistry</i> , 2010, 285, 39425-39436. | 3.4 | 25 |
| 40 | Synthesis, Biological Activity, and Conformational Analysis of Peptidomimetic Analogues of the <i>Saccharomyces cerevisiae</i> α -Factor Tridecapeptide. <i>Biochemistry</i> , 1998, 37, 12465-12476. | 2.5 | 24 |
| 41 | Peptide fragments as models to study the structure of a G-protein coupled receptor: The α -factor receptor of <i>Saccharomyces cerevisiae</i> . <i>Biopolymers</i> , 2001, 60, 334. | 2.4 | 24 |
| 42 | Identification of residues involved in homodimer formation located within a β -strand region of the N-terminus of a Yeast G protein-coupled receptor. <i>Journal of Receptor and Signal Transduction Research</i> , 2012, 32, 65-75. | 2.5 | 24 |
| 43 | Interacting Residues in an Activated State of a G Protein-coupled Receptor. <i>Journal of Biological Chemistry</i> , 2006, 281, 2263-2272. | 3.4 | 23 |
| 44 | Differential Interactions of Fluorescent Agonists and Antagonists with the Yeast G Protein Coupled Receptor Ste2p. <i>Journal of Molecular Biology</i> , 2011, 409, 513-528. | 4.2 | 23 |
| 45 | Detection of intermolecular NOE interactions in large protein complexes. <i>Progress in Nuclear Magnetic Resonance Spectroscopy</i> , 2016, 97, 40-56. | 7.5 | 23 |
| 46 | Changes in Conformation at the Cytoplasmic Ends of the Fifth and Sixth Transmembrane Helices of a Yeast G Protein-Coupled Receptor in Response to Ligand Binding. <i>Biochemistry</i> , 2011, 50, 6841-6854. | 2.5 | 22 |
| 47 | Double-Mutant Cycle Scanning of the Interaction of a Peptide Ligand and Its G Protein-Coupled Receptor. <i>Biochemistry</i> , 2007, 46, 3476-3481. | 2.5 | 21 |
| 48 | Synthesis of N-terminal and C-terminal analogs of the <i>Saccharomyces cerevisiae</i> α -factor. <i>International Journal of Peptide and Protein Research</i> , 1991, 37, 476-486. | 0.1 | 21 |
| 49 | Sexual conjugation in yeast: A paradigm to study G-protein-coupled receptor domain structure. <i>Biopolymers</i> , 2004, 76, 119-128. | 2.4 | 19 |
| 50 | Multiple regulatory roles of the carboxy terminus of Ste2p a yeast GPCR. <i>Pharmacological Research</i> , 2012, 65, 31-40. | 7.1 | 19 |
| 51 | SYNTHESIS OF THE DODECAPEPTIDE α -MATING FACTOR OF <i>SACCHAROMYCES CEREVISIAE</i> . <i>International Journal of Peptide and Protein Research</i> , 1981, 17, 219-230. | 0.1 | 18 |
| 52 | Systematic analysis of the <i>Saccharomyces cerevisiae</i> α -factor containing lactam constraints of different ring size. <i>Biochemistry</i> , 1995, 34, 1308-1315. | 2.5 | 17 |
| 53 | Studies on the yeast β -mating factor: A model for mammalian peptide hormones. <i>Biopolymers</i> , 1992, 32, 335-339. | 2.4 | 16 |
| 54 | Synthesis of a Double Transmembrane Domain Fragment of Ste2p by Native Chemical Ligation. <i>International Journal of Peptide Research and Therapeutics</i> , 2007, 13, 251-263. | 1.9 | 16 |

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 55 | Biosynthesis and NMR-studies of a double transmembrane domain from the Y4 receptor, a human GPCR. <i>Journal of Biomolecular NMR</i> , 2008, 42, 257-269. | 2.8 | 16 |
| 56 | The N-terminus of the yeast G protein-coupled receptor Ste2p plays critical roles in surface expression, signaling, and negative regulation. <i>Biochimica Et Biophysica Acta - Biomembranes</i> , 2016, 1858, 715-724. | 2.6 | 16 |
| 57 | Biologically significant conformation of the <i>Saccharomyces cerevisiae</i> β -factor. <i>Biopolymers</i> , 1989, 28, 487-497. | 2.4 | 15 |
| 58 | ¹ H-nmr study of protected methionine homo-oligopeptides in helix-supporting environment. <i>Biopolymers</i> , 1980, 19, 1791-1799. | 2.4 | 14 |
| 59 | Synthesis and Biophysical Characterization of a Multidomain Peptide from a <i>Saccharomyces cerevisiae</i> G Protein-coupled Receptor. <i>Journal of Biological Chemistry</i> , 2003, 278, 52537-52545. | 3.4 | 14 |
| 60 | Selective labeling of a membrane peptide with ¹⁵ N-amino acids using cells grown in rich medium. <i>Biopolymers</i> , 2006, 84, 508-518. | 2.4 | 12 |
| 61 | Studies on conformational consequences of <i>i</i> to <i>i</i> +3 side-chain cyclization in model cyclic tetrapeptides. <i>International Journal of Peptide and Protein Research</i> , 1995, 45, 418-429. | 0.1 | 12 |
| 62 | Position one analogs of the <i>Saccharomyces cerevisiae</i> tridecapeptide pheromone. <i>Chemical Biology and Drug Design</i> , 1997, 50, 319-328. | 1.1 | 12 |
| 63 | Conformational analysis of cyclic analogues of the <i>Saccharomyces cerevisiae</i> β -factor pheromone. , 1998, 45, 21-34. | | 10 |
| 64 | Biosynthesis of peptide fragments of eukaryotic GPCRs in <i>Escherichia coli</i> by directing expression into inclusion bodies. <i>Journal of Peptide Science</i> , 2010, 16, 213-218. | 1.4 | 9 |
| 65 | Invited review GPCR structural characterization: Using fragments as building blocks to determine a complete structure. <i>Biopolymers</i> , 2014, 102, 223-243. | 2.4 | 8 |
| 66 | Detection of intermolecular transferred ¹ H NOEs in large protein complexes using asymmetric deuteration: ¹ H gp120 in complex with a CCR5 peptide. <i>FEBS Journal</i> , 2016, 283, 4084-4096. | 4.7 | 8 |
| 67 | Variable Dependence of Signaling Output on Agonist Occupancy of Ste2p, a G Protein-coupled Receptor in Yeast. <i>Journal of Biological Chemistry</i> , 2016, 291, 24261-24279. | 3.4 | 8 |
| 68 | Synthetic probes for the β -factor receptor. <i>Biopolymers</i> , 1990, 29, 237-245. | 2.4 | 7 |
| 69 | Probing the functional conformation of the tridecapeptide mating pheromone of <i>Saccharomyces cerevisiae</i> through study of disulfide-constrained analogs. <i>International Journal of Peptide and Protein Research</i> , 1996, 47, 131-141. | 0.1 | 7 |
| 70 | Identification of Destabilizing and Stabilizing Mutations of Ste2p, a G Protein-Coupled Receptor in <i>Saccharomyces cerevisiae</i> . <i>Biochemistry</i> , 2015, 54, 1787-1806. | 2.5 | 7 |
| 71 | Allovalency observed by transferred NOE: interactions of sulfated tyrosine residues in the N-terminal segment of CCR5 with the CCL5 chemokine. <i>FEBS Journal</i> , 2021, 288, 1648-1663. | 4.7 | 7 |
| 72 | The methyl ¹³ C-edited/ ¹³ C-filtered transferred NOE for studying protein interactions with short linear motifs. <i>Journal of Biomolecular NMR</i> , 2020, 74, 681-693. | 2.8 | 7 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 73 | Synthesis of \pm -factor analogues containing photoactivatable and labeling groups. International Journal of Peptide and Protein Research, 1995, 45, 106-115. | 0.1 | 6 |
| 74 | The C4 region as a target for HIV entry inhibitors – NMR mapping of the interacting segments of T20 and gp120. FEBS Journal, 2015, 282, 4643-4657. | 4.7 | 6 |
| 75 | NMR Investigation of Structures of G-protein Coupled Receptor Folding Intermediates. Journal of Biological Chemistry, 2016, 291, 27170-27186. | 3.4 | 6 |
| 76 | Dynamic roles for the N-terminus of the yeast G protein-coupled receptor Ste2p. Biochimica Et Biophysica Acta - Biomembranes, 2017, 1859, 2058-2067. | 2.6 | 6 |
| 77 | Identification of peptide-binding sites within BSA using rapid, laser-induced covalent cross-linking combined with high-performance mass spectrometry. Journal of Molecular Recognition, 2018, 31, e2680. | 2.1 | 6 |
| 78 | A Paradigm for Peptide Hormone-GPCR Analyses. Molecules, 2020, 25, 4272. | 3.8 | 6 |
| 79 | Synthesis of biologically active analogs of the dodecapeptide \pm -factor mating pheromone of <i>Saccharomyces cerevisiae</i> . International Journal of Peptide and Protein Research, 1990, 35, 241-248. | 0.1 | 5 |
| 80 | An extended CCR5 ECL2 peptide forms a helix that binds HIV-1 gp120 through non-specific hydrophobic interactions. FEBS Journal, 2015, 282, 1906-1921. | 4.7 | 5 |
| 81 | Synthesis and biological activity of N-acyl derivatives of a <i>Saccharomyces cerevisiae</i> mating pheromone. International Journal of Peptide and Protein Research, 1985, 25, 187-196. | 0.1 | 4 |
| 82 | Comparison of Fragments Comprising the First Two Helices of the Human Y4 and the Yeast Ste2p G-Protein-Coupled Receptors. Biophysical Journal, 2012, 103, 817-826. | 0.5 | 4 |
| 83 | Cross-linking Strategies to Study Peptide Ligand-Receptor Interactions. Methods in Enzymology, 2015, 556, 527-547. | 1.0 | 4 |
| 84 | Halo Assay for Toxic Peptides and Other Compounds in Microorganisms. Bio-protocol, 2016, 6, . | 0.4 | 4 |
| 85 | Structural characterization of triple transmembrane domain containing fragments of a yeast G protein-coupled receptor in an organic-aqueous environment by solution-state NMR spectroscopy. Journal of Peptide Science, 2015, 21, 212-222. | 1.4 | 3 |
| 86 | Novobiocin and peptide analogs of \pm -factor are positive allosteric modulators of the yeast G protein-coupled receptor Ste2p. Biochimica Et Biophysica Acta - Biomembranes, 2015, 1848, 916-924. | 2.6 | 3 |
| 87 | The Synthesis of Sulfated CCR5 Peptide Surrogates and their Use to Study Receptor-Ligand Interactions. Protein and Peptide Letters, 2019, 25, 1124-1136. | 0.9 | 3 |
| 88 | Immunofocusing using conformationally constrained V3 peptide immunogens improves HIV-1 neutralization. Vaccine, 2017, 35, 222-230. | 3.8 | 2 |
| 89 | Defining specific residue-residue interactions between the gp120 bridging sheet and the N-terminal segment of CCR5: applications of transferred NOE NMR. FEBS Journal, 2018, 285, 4296-4310. | 4.7 | 2 |
| 90 | Multiple binding modes of an N-terminal CCR5-peptide in complex with HIV-1 gp120. FEBS Journal, 2022, 289, 3132-3147. | 4.7 | 2 |

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 91 | Mobility of oligopeptides on normal-phase silica: Effect of positional isomerism. <i>Biopolymers</i> , 1983, 22, 1401-1407. | 2.4 | 1 |
| 92 | Conformational studies of nikkomycin X in aqueous solution. <i>Biopolymers</i> , 1990, 29, 1297-1306. | 2.4 | 1 |
| 93 | Oligomerization of yeast $\hat{\mu}$ -factor receptor detected by fluorescent energy transfer between ligands. <i>Biophysical Journal</i> , 2021, 120, 5090-5106. | 0.5 | 1 |
| 94 | Synthesis, Biosynthesis, and Characterization of Transmembrane Domains of a G Protein-Coupled Receptor. <i>Methods in Molecular Biology</i> , 2007, 386, 95-121. | 0.9 | 1 |
| 95 | Uptake Assay for Radiolabeled Peptides in Yeast. <i>Bio-protocol</i> , 2016, 6, . | 0.4 | 1 |
| 96 | Structural Studies on Large Fragments of G Protein Coupled Receptors. <i>Advances in Experimental Medicine and Biology</i> , 2009, 611, 309-310. | 1.6 | 0 |