
Rui M Monteiro

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/4418164/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	In the spotlight: the role of TGFβ signalling in haematopoietic stem and progenitor cell emergence. Biochemical Society Transactions, 2022, 50, 703-712.	3.4	3
2	Epigenetic Regulation of Endothelial Cell Lineages During Zebrafish Development—New Insights From Technical Advances. Frontiers in Cell and Developmental Biology, 2022, 10, .	3.7	1
3	Essential role for Gata2 in modulating lineage output from hematopoietic stem cells in zebrafish. Blood Advances, 2021, 5, 2687-2700.	5.2	21
4	<i>Hapln1b</i> , a central organizer of the ECM, modulates kit signaling to control developmental hematopoiesis in zebrafish. Blood Advances, 2021, 5, 4935-4948.	5.2	7
5	The roles and controls of GATA factors in blood and cardiac development. IUBMB Life, 2020, 72, 39-44.	3.4	21
6	Deletion of a conserved Gata2 enhancer impairs haemogenic endothelium programming and adult Zebrafish haematopoiesis. Communications Biology, 2020, 3, 71.	4.4	26
7	Genotyping and Quantification of In Situ Hybridization Staining in Zebrafish. Journal of Visualized Experiments, 2020, , .	0.3	11
8	3115 – GATA2A DETERMINES THE SURVIVAL AND HOMEOSTATIC LINEAGE OUTPUT OF HAEMATOPOIETIC STE AND PROGENITOR CELLS. Experimental Hematology, 2020, 88, S73.	M _{0.4}	0
9	Blood stem cell-forming haemogenic endothelium in zebrafish derives from arterial endothelium. Nature Communications, 2019, 10, 3577.	12.8	37
10	An optimized pipeline for parallel image-based quantification of gene expression and genotyping after <i>in situ</i> hybridization. Biology Open, 2018, 7, .	1.2	21
11	Functional analysis of a Gata2a endothelial enhancer reveals non-redundant roles for Gata2a and Gata2b in haemogenic endothelium programming and generation of haematopoietic stem cells. Experimental Hematology, 2017, 53, S77.	0.4	0
12	Transforming Growth Factor β Drives Hemogenic Endothelium Programming and the Transition to Hematopoietic Stem Cells. Developmental Cell, 2016, 38, 358-370.	7.0	75
13	BMP and Hedgehog Regulate Distinct AGM Hematopoietic Stem Cells ExÂVivo. Stem Cell Reports, 2016, 6, 383-395.	4.8	37
14	A Novel TGFβ Modulator that Uncouples R-Smad/I-Smad-Mediated Negative Feedback from R-Smad/Ligand-Driven Positive Feedback. PLoS Biology, 2015, 13, e1002051.	5.6	7
15	BMP signalling differentially regulates distinct haematopoietic stem cell types. Nature Communications, 2015, 6, 8040.	12.8	74
16	The heparan sulfate editing enzyme Sulf1 plays a novel role in zebrafish VegfA mediated arterial venous identity. Angiogenesis, 2014, 17, 77-91.	7.2	16
17	Developmental hematopoiesis: Ontogeny, genetic programming and conservation. Experimental Hematology, 2014, 42, 669-683.	0.4	110
18	Uncoupling VEGFA Functions in Arteriogenesis and Hematopoietic Stem Cell Specification. Developmental Cell, 2013, 24, 144-158.	7.0	58

RUI M MONTEIRO

#	Article	IF	CITATIONS
19	Analysis of <i>Dll4</i> regulation reveals a combinatorial role for Sox and Notch in arterial development. Proceedings of the National Academy of Sciences of the United States of America, 2013, 110, 11893-11898.	7.1	114
20	The microRNA-30 family targets DLL4 to modulate endothelial cell behavior during angiogenesis. Blood, 2012, 120, 5063-5072.	1.4	163
21	A Novel Complex, RUNX1-MYEF2, Represses Hematopoietic Genes in Erythroid Cells. Molecular and Cellular Biology, 2012, 32, 3814-3822.	2.3	32
22	Activation of the Canonical Bone Morphogenetic Protein (BMP) Pathway during Lung Morphogenesis and Adult Lung Tissue Repair. PLoS ONE, 2012, 7, e41460.	2.5	60
23	The gata1/pu.1 lineage fate paradigm varies between blood populations and is modulated by tif1γ. EMBO Journal, 2011, 30, 1093-1103.	7.8	81
24	CHAP is a newly identified Z-disc protein essential for heart and skeletal muscle function. Journal of Cell Science, 2010, 123, 1141-1150.	2.0	53
25	Real time monitoring of BMP Smads transcriptional activity during mouse development. Genesis, 2008, 46, 335-346.	1.6	70
26	Real time monitoring of BMP Smads transcriptional activity during mouse development. Genesis, 2008, 46, spcone-spcone.	1.6	3
27	Two novel type II receptors mediate BMP signalling and are required to establish left–right asymmetry in zebrafish. Developmental Biology, 2008, 315, 55-71.	2.0	54
28	Adult Neurogenesis Requires Smad4-Mediated Bone Morphogenic Protein Signaling in Stem Cells. Journal of Neuroscience, 2008, 28, 434-446.	3.6	228
29	SOST expression is restricted to the great arteries during embryonic and neonatal cardiovascular development. Developmental Dynamics, 2007, 236, 606-612.	1.8	41
30	Isl1Cre reveals a common Bmp pathway in heart and limb development. Development (Cambridge), 2006, 133, 1575-1585.	2.5	234
31	Expression of bone morphogenetic protein2 (BMP2), BMP4 and BMP receptors in the bovine ovary but absence of effects of BMP2 and BMP4 during IVM on bovine oocyte nuclear maturation and subsequent embryo development. Theriogenology, 2005, 63, 872-889.	2.1	69
32	BMP signaling mediated by ALK2 in the visceral endoderm is necessary for the generation of primordial germ cells in the mouse embryo. Genes and Development, 2004, 18, 1838-1849.	5.9	180
33	Spatio-temporal activation of Smad1 and Smad5 in vivo: monitoring transcriptional activity of Smad proteins. Journal of Cell Science, 2004, 117, 4653-4663.	2.0	81