

# Nãria Plana

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4422001/publications.pdf>

Version: 2024-02-01

56  
papers

1,451  
citations

304368

22  
h-index

360668

35  
g-index

61  
all docs

61  
docs citations

61  
times ranked

1952  
citing authors

#	ARTICLE	IF	CITATIONS
1	Effect of statin therapy on SARS-CoV-2 infection-related mortality in hospitalized patients. <i>European Heart Journal - Cardiovascular Pharmacotherapy</i> , 2022, 8, 157-164.	1.4	64
2	Lipoprotein(a) in hereditary hypercholesterolemia: Influence of the genetic cause, defective gene and type of mutation. <i>Atherosclerosis</i> , 2022, 349, 211-218.	0.4	12
3	PatrÃ³n de metilaciÃ³n en ADN de sujetos hipertrigliceridÃ©micos. <i>ClÃnica E InvestigaciÃ³n En Arteriosclerosis</i> , 2022, 34, 27-32.	0.4	2
4	El rastreo masivo de datos es una segunda oportunidad para mejorar el manejo de los pacientes fenotipo de hipercolesterolemia familiar. <i>ClÃnica E InvestigaciÃ³n En Arteriosclerosis</i> , 2021, 33, 138-147.	0.4	1
5	Maternally inherited hypercholesterolemia does not modify the cardiovascular phenotype in familial hypercholesterolemia. <i>Atherosclerosis</i> , 2021, 320, 47-52.	0.4	7
6	Low HDL and high triglycerides predict COVID-19 severity. <i>Scientific Reports</i> , 2021, 11, 7217.	1.6	122
7	Massive data screening is a second opportunity to improve the management of patients with familial hypercholesterolemia phenotype. <i>ClÃnica E InvestigaciÃ³n En Arteriosclerosis (English Edition)</i> , 2021, 33, 138-147.	0.1	2
8	Impact of statin therapy on LDL and non-HDL cholesterol levels in subjects with heterozygous familial hypercholesterolaemia. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2021, 31, 1594-1603.	1.1	9
9	Serum glycoproteins A and B assessed by 1H-NMR in familial hypercholesterolemia. <i>Atherosclerosis</i> , 2021, 330, 1-7.	0.4	9
10	Triglyceride-Rich Lipoproteins and Glycoprotein A and B Assessed by 1H-NMR in Metabolic-Associated Fatty Liver Disease. <i>Frontiers in Endocrinology</i> , 2021, 12, 775677.	1.5	4
11	Relationship Between Fatty Acid Binding Protein 4 and Liver Fat in Individuals at Increased Cardiometabolic Risk. <i>Frontiers in Physiology</i> , 2021, 12, 781789.	1.3	5
12	Derivation and validation of SIDIAP-FHP score: A new risk model predicting cardiovascular disease in familial hypercholesterolemia phenotype. <i>Atherosclerosis</i> , 2020, 292, 42-51.	0.4	9
13	Clinical and genetic differences between heterozygous familial hypercholesterolemia patients with and without type 2 diabetes. <i>Revista Espanola De Cardiologia (English Ed )</i> , 2020, 73, 718-724.	0.4	6
14	Polygenic Markers in Patients Diagnosed of Autosomal Dominant Hypercholesterolemia in Catalonia: Distribution of Weighted LDL-c-Raising SNP Scores and Refinement of Variant Selection. <i>Biomedicines</i> , 2020, 8, 353.	1.4	6
15	Reasons Why Combination Therapy Should Be the New Standard of Care to Achieve the LDL-Cholesterol Targets. <i>Current Cardiology Reports</i> , 2020, 22, 66.	1.3	26
16	Diferencias clÃnicas y genÃ©ticas de los pacientes con hipercolesterolemia familiar heterocigota con y sin diabetes mellitus tipo 2. <i>Revista Espanola De Cardiologia</i> , 2020, 73, 718-724.	0.6	11
17	Efficacy of therapeutic lifestyle changes on lipid profiles assessed by NMR in children with familial and non-familial hypercholesterolemia. <i>ClÃnica E InvestigaciÃ³n En Arteriosclerosis</i> , 2020, 32, 49-58.	0.4	3
18	Incidence of Cardiovascular Disease in Patients with Familial Hypercholesterolemia Phenotype: Analysis of 5 Years Follow-Up of Real-World Data from More than 1.5 Million Patients. <i>Journal of Clinical Medicine</i> , 2019, 8, 1080.	1.0	33

#	ARTICLE	IF	CITATIONS
19	HDL Triglycerides: A New Marker of Metabolic and Cardiovascular Risk. <i>International Journal of Molecular Sciences</i> , 2019, 20, 3151.	1.8	58
20	EstĂndares SEA 2019 para el control global del riesgo cardiovascular. <i>ClĂnica E InvestigaciĂn En Arteriosclerosis</i> , 2019, 31, 1-43.	0.4	8
21	The Circulating GRP78/BiP Is a Marker of Metabolic Diseases and Atherosclerosis: Bringing Endoplasmic Reticulum Stress into the Clinical Scenario. <i>Journal of Clinical Medicine</i> , 2019, 8, 1793.	1.0	40
22	Toward a new clinical classification of patients with familial hypercholesterolemia: One perspective from Spain. <i>Atherosclerosis</i> , 2019, 287, 89-92.	0.4	29
23	Estimated Percentage of Patients With Stable Coronary Heart Disease Candidates for PCSK9 Inhibitors. Response. <i>Revista Espanola De Cardiologia (English Ed )</i> , 2019, 72, 519-520.	0.4	1
24	Effect of lipid-lowering treatment in cardiovascular disease prevalence in familial hypercholesterolemia. <i>Atherosclerosis</i> , 2019, 284, 245-252.	0.4	55
25	Comparative efficacy between atorvastatin and rosuvastatin in the prevention of cardiovascular disease recurrence. <i>Lipids in Health and Disease</i> , 2019, 18, 216.	1.2	16
26	ActualizaciĂn de las tablas de planificaciĂn terapĂutica hipocolesterolemiantes orientadas a la obtenciĂn de los objetivos terapĂuticos. <i>ClĂnica E InvestigaciĂn En Arteriosclerosis</i> , 2019, 31, 271-277.	0.4	13
27	Valor de los parĂmetros lipĂdicos y apoproteicos para la detecciĂn de hipercolesterolemia familiar en la infancia. Proyecto DECOPIN. <i>ClĂnica E InvestigaciĂn En Arteriosclerosis</i> , 2018, 30, 170-178.	0.4	9
28	Number of Patients Eligible for PCSK9 Inhibitors Based on Real-world Data From 2.5 Million Patients. <i>Revista Espanola De Cardiologia (English Ed )</i> , 2018, 71, 1010-1017.	0.4	10
29	Clinical and pathophysiological evidence supporting the safety of extremely low LDL levelsâ€”The zero-LDL hypothesis. <i>Journal of Clinical Lipidology</i> , 2018, 12, 292-299.e3.	0.6	51
30	Lipoprotein profile assessed by 2D-1H-NMR and subclinical atherosclerosis in children with familial hypercholesterolaemia. <i>Atherosclerosis</i> , 2018, 270, 117-122.	0.4	11
31	NĂmero de pacientes candidatos a recibir inhibidores de la PCSK9 segĂn datos de 2,5 millones de participantes de la prĂctica clĂnica real. <i>Revista Espanola De Cardiologia</i> , 2018, 71, 1010-1017.	0.6	23
32	Causas de no consecuciĂn del objetivo terapĂutico del colesterol de las lipoproteĂnas de baja densidad en pacientes de alto y muy alto riesgo vascular controlados en Unidades de LĂpidos y Riesgo Vascular. Estudio EROMOT. <i>ClĂnica E InvestigaciĂn En Arteriosclerosis</i> , 2018, 30, 1-9.	0.4	8
33	Plasma inducible degrader of the LDLR, soluble low-density lipoprotein receptor, and proprotein convertase subtilisin/kexin type 9 levels as potential biomarkers of familial hypercholesterolemia in children. <i>Journal of Clinical Lipidology</i> , 2018, 12, 211-218.	0.6	14
34	Autosomal dominant hypercholesterolemia in Catalonia: Correspondence between clinical-biochemical and genetic diagnostics in 967 patients studied in a multicenter clinical setting. <i>Journal of Clinical Lipidology</i> , 2018, 12, 1452-1462.	0.6	14
35	Detecting familial hypercholesterolemia earlier in life by actively searching for affected children: The DECOPIN project. <i>Atherosclerosis</i> , 2018, 278, 210-216.	0.4	18
36	Value of the Definition of Severe Familial Hypercholesterolemia for Stratification of Heterozygous Patients. <i>American Journal of Cardiology</i> , 2017, 119, 742-748.	0.7	17

#	ARTICLE	IF	CITATIONS
37	How many familial hypercholesterolemia patients are eligible for PCSK9 inhibition?. <i>Atherosclerosis</i> , 2017, 262, 107-112.	0.4	22
38	Familial hypercholesterolemia in a European Mediterranean populationâ€™Prevalence and clinical data from 2.5 million primary care patients. <i>Journal of Clinical Lipidology</i> , 2017, 11, 1013-1022.	0.6	61
39	Effect of LDL cholesterol, statins and presence of mutations on the prevalence of type 2 diabetes in heterozygous familial hypercholesterolemia. <i>Scientific Reports</i> , 2017, 7, 5596.	1.6	41
40	Hipercolesterolemia familiar en la infancia y la adolescencia: una realidad oculta. <i>ClĂnica E InvestigaciĂn En Arteriosclerosis</i> , 2017, 29, 129-140.	0.4	8
41	Registro Nacional de Dislipemias de la Sociedad EspaĂola de Arteriosclerosis: situaciĂn actual. <i>ClĂnica E InvestigaciĂn En Arteriosclerosis</i> , 2017, 29, 248-253.	0.4	20
42	Lipoprotein hydrophobic core lipids are partially extruded to surface in smaller HDL: â€™Herniatedâ€™HDL, a common feature in diabetes. <i>Scientific Reports</i> , 2016, 6, 19249.	1.6	25
43	MĂxima reducciĂn de colesterol unido a lipoproteĂnas de baja densidad alcanzable con combinaciones farmacolĂgicas. Cuando 50 mĂs 20 suma 60. <i>Revista Espanola De Cardiologia</i> , 2016, 69, 342-343.	0.6	24
44	Circulating PCSK9 levels and CETP plasma activity are independently associated in patients with metabolic diseases. <i>Cardiovascular Diabetology</i> , 2016, 15, 107.	2.7	28
45	Impact of epidermal fatty acid binding protein on 2D-NMRâ€™assessed atherogenic dyslipidemia and related disorders. <i>Journal of Clinical Lipidology</i> , 2016, 10, 330-338.e2.	0.6	9
46	Circulating PCSK9 in patients with type 2 diabetes and related metabolic disorders. <i>ClĂnica E InvestigaciĂn En Arteriosclerosis</i> , 2016, 28, 71-78.	0.4	35
47	Novel mutations in the GPIHBP1 gene identified in 2 patients with recurrent acute pancreatitis. <i>Journal of Clinical Lipidology</i> , 2016, 10, 92-100.e1.	0.6	27
48	Letter to Editor: Increased Presence of Remnant Lipoprotein Cholesterol in The Hdl of Diabetic Subjects. <i>Annals of Clinical and Laboratory Science</i> , 2016, 46, 229-32.	0.2	0
49	Liposcale: a novel advanced lipoprotein test based on 2D diffusion-ordered 1H NMR spectroscopy. <i>Journal of Lipid Research</i> , 2015, 56, 737-746.	2.0	133
50	APOA5 variants predispose hyperlipidemic patients to atherogenic dyslipidemia and subclinical atherosclerosis. <i>Atherosclerosis</i> , 2015, 240, 98-104.	0.4	28
51	Remarkable quantitative and qualitative differences in HDL after niacin or fenofibrate therapy in type 2 diabetic patients. <i>Atherosclerosis</i> , 2015, 238, 213-219.	0.4	23
52	Low-carbohydrate, high-protein, high-fat diet alters small peripheral artery reactivity in metabolic syndrome patients. <i>ClĂnica E InvestigaciĂn En Arteriosclerosis</i> , 2014, 26, 58-65.	0.4	5
53	Increasing long-chain n-3PUFA consumption improves small peripheral artery function in patients at intermediateâ€™high cardiovascular risk. <i>Journal of Nutritional Biochemistry</i> , 2014, 25, 642-646.	1.9	19
54	Heterozygous Familial Hypercholesterolaemic Patients have Increased Arterial Stiffness, as Determined using the Augmentation Index. <i>Journal of Atherosclerosis and Thrombosis</i> , 2011, 18, 1110-1116.	0.9	18

#	ARTICLE	IF	CITATIONS
55	Comparison of Genetic Versus Clinical Diagnosis in Familial Hypercholesterolemia. American Journal of Cardiology, 2008, 102, 1187-1193.e1.	0.7	153
56	Physicochemical changes in HDL3 after bezafibrate treatment: influence on free cholesterol efflux from human fibroblasts. Cardiovascular Drugs and Therapy, 1997, 11, 653-658.	1.3	7