

Gerard C Blobe

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4442487/publications.pdf>

Version: 2024-02-01

122
papers

11,288
citations

44042

48
h-index

29127

104
g-index

123
all docs

123
docs citations

123
times ranked

15407
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 1 | <sc>TGF</sc> β^2 superfamily co-receptors in cancer. <i>Developmental Dynamics</i> , 2022, 251, 117-143. | 0.8 | 33 |
| 2 | Cabozantinib and Panitumumab for RAS Wild-Type Metastatic Colorectal Cancer. <i>Oncologist</i> , 2021, 26, 465-e917. | 1.9 | 13 |
| 3 | ALK1 regulates the internalization of endoglin and the type III TGF- β^2 receptor. <i>Molecular Biology of the Cell</i> , 2021, 32, 605-621. | 0.9 | 8 |
| 4 | The role of the extracellular matrix protein TGFBI in cancer. <i>Cellular Signalling</i> , 2021, 84, 110028. | 1.7 | 38 |
| 5 | KEYlargo: A phase II study of first-line pembrolizumab (P), capecitabine (C), and oxaliplatin (O) in HER2-negative gastroesophageal (GE) adenocarcinoma. <i>Journal of Clinical Oncology</i> , 2021, 39, 228-228. | 0.8 | 2 |
| 6 | A phase Ib study of the combination regorafenib with PF-03446962 in patients with refractory metastatic colorectal cancer (REGAL-1 trial). <i>Cancer Chemotherapy and Pharmacology</i> , 2019, 84, 909-917. | 1.1 | 13 |
| 7 | Imaging & Biomarker Correlates on Outcomes in a Phase II Trial of Neoadjuvant Gemcitabine/Nab-Paclitaxel and Hypofractionated Image-Guided Radiotherapy (HIGRT) in Potentially Resectable Pancreas Cancer. <i>International Journal of Radiation Oncology Biology Physics</i> , 2019, 105, E238-E239. | 0.4 | 0 |
| 8 | Increased type III TGF- β^2 receptor shedding decreases tumorigenesis through induction of epithelial-to-mesenchymal transition. <i>Oncogene</i> , 2019, 38, 3402-3414. | 2.6 | 15 |
| 9 | Surgical and Pathologic Outcomes in Patients on a Phase II Trial of Neoadjuvant Chemotherapy and Hypofractionated Image-Guided Intensity Modulated Radiation Therapy (HIGRT) in Resectable and Borderline Resectable Pancreatic Cancer. <i>International Journal of Radiation Oncology Biology Physics</i> , 2018, 102, S180. | 0.4 | 0 |
| 10 | Stromal Fibroblasts Mediate Anti-PD-1 Resistance via MMP-9 and Dictate TGF- β^2 Inhibitor Sequencing in Melanoma. <i>Cancer Immunology Research</i> , 2018, 6, 1459-1471. | 1.6 | 81 |
| 11 | Endoglin interacts with VEGFR2 to promote angiogenesis. <i>FASEB Journal</i> , 2018, 32, 2934-2949. | 0.2 | 56 |
| 12 | A phase II trial of neoadjuvant gemcitabine/nab-paclitaxel and SBRT for potentially resectable pancreas cancer: An evaluation of acute toxicity. <i>Journal of Clinical Oncology</i> , 2018, 36, 4121-4121. | 0.8 | 3 |
| 13 | Heparin-binding epidermal growth factor-like growth factor promotes neuroblastoma differentiation. <i>FASEB Journal</i> , 2017, 31, 1903-1915. | 0.2 | 14 |
| 14 | Endoglin Mediates Vascular Maturation by Promoting Vascular Smooth Muscle Cell Migration and Spreading. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2017, 37, 1115-1126. | 1.1 | 27 |
| 15 | Phase I study of pazopanib plus TH-302 in advanced solid tumors. <i>Cancer Chemotherapy and Pharmacology</i> , 2017, 79, 611-619. | 1.1 | 8 |
| 16 | TGF- β^2 -induced stromal CYR61 promotes resistance to gemcitabine in pancreatic ductal adenocarcinoma through downregulation of the nucleoside transporters hENT1 and hCNT3. <i>Carcinogenesis</i> , 2016, 37, 1041-1051. | 1.3 | 67 |
| 17 | Dichotomous roles of TGF- β^2 in human cancer. <i>Biochemical Society Transactions</i> , 2016, 44, 1441-1454. | 1.6 | 91 |
| 18 | An Automated High-throughput Array Microscope for Cancer Cell Mechanics. <i>Scientific Reports</i> , 2016, 6, 27371. | 1.6 | 5 |

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 19 | Abstract 1182: Heparin-binding epidermal growth factor-like growth factor is a pro-differentiating factor in neuroblastoma. , 2016, , . | | 3 |
| 20 | Phase Ib study of cabozantinib plus panitumumab in KRAS wild-type (WT) metastatic colorectal cancer (mCRC).. Journal of Clinical Oncology, 2016, 34, 3548-3548. | 0.8 | 4 |
| 21 | Phase Ib study of regorafenib (rego) and PF-03446962 (PF) in patients with refractory metastatic colorectal cancer (mCRC) (REGAL).. Journal of Clinical Oncology, 2016, 34, e15013-e15013. | 0.8 | 2 |
| 22 | TÎ²RIII independently binds type I and type II TGF-Î² receptors to inhibit TGF-Î² signaling. Molecular Biology of the Cell, 2015, 26, 3535-3545. | 0.9 | 28 |
| 23 | Fibulin-3 is a novel TGF-Î² pathway inhibitor in the breast cancer microenvironment. Oncogene, 2015, 34, 5635-5647. | 2.6 | 41 |
| 24 | Regulation of TGF-Î² receptor hetero-oligomerization and signaling by endoglin. Molecular Biology of the Cell, 2015, 26, 3117-3127. | 0.9 | 35 |
| 25 | Angiotensin II stimulates canonical TGF-Î² signaling pathway through angiotensin type 1 receptor to induce granulation tissue contraction. Journal of Molecular Medicine, 2015, 93, 289-302. | 1.7 | 42 |
| 26 | Dalantercept. ALK-1 ligand trap, Angiogenesis inhibitor, treatment of solid tumors. Drugs of the Future, 2015, 40, 0633. | 0.0 | 0 |
| 27 | Safety, Pharmacokinetics, Pharmacodynamics, and Antitumor Activity of Dalantercept, an Activin Receptorâ€™like Kinase-1 Ligand Trap, in Patients with Advanced Cancer. Clinical Cancer Research, 2014, 20, 480-489. | 3.2 | 52 |
| 28 | Role of TGF-Î² receptor III localization in polarity and breast cancer progression. Molecular Biology of the Cell, 2014, 25, 2291-2304. | 0.9 | 17 |
| 29 | TGF-Î² regulates LARG and GEF-H1 during EMT to affect stiffening response to force and cell invasion. Molecular Biology of the Cell, 2014, 25, 3528-3540. | 0.9 | 53 |
| 30 | Ectodomain shedding of TÎ²RIII is required for TÎ²RIII-mediated suppression of TGF-Î² signaling and breast cancer migration and invasion. Molecular Biology of the Cell, 2014, 25, 2320-2332. | 0.9 | 39 |
| 31 | Phase I study of dasatinib in combination with capecitabine, oxaliplatin and bevacizumab followed by an expanded cohort in previously untreated metastatic colorectal cancer. Investigational New Drugs, 2014, 32, 330-339. | 1.2 | 18 |
| 32 | Heparan sulfate signaling in cancer. Trends in Biochemical Sciences, 2014, 39, 277-288. | 3.7 | 154 |
| 33 | Novel bone morphogenetic protein signaling through Smad2 and Smad3 to regulate cancer progression and development. FASEB Journal, 2014, 28, 1248-1267. | 0.2 | 80 |
| 34 | Effects of the combination of TRC105 and bevacizumab on endothelial cell biology. Investigational New Drugs, 2014, 32, 851-859. | 1.2 | 40 |
| 35 | The Balance of Cell Surface and Soluble Type III TGF-Î² Receptor Regulates BMP Signaling in Normal and Cancerous Mammary Epithelial Cells. Neoplasia, 2014, 16, 489-500. | 2.3 | 22 |
| 36 | Angiotensin II Stimulates Canonical TGFÎ² Signaling Pathway through Angiotensin Receptor 1 to Induce Granulation Tissue Contraction. Journal of the American College of Surgeons, 2014, 219, S84. | 0.2 | 0 |

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 37 | Stromal heparan sulfate differentiates neuroblasts to suppress neuroblastoma growth. Journal of Clinical Investigation, 2014, 124, 3016-3031. | 3.9 | 28 |
| 38 | Combinatorial TGF- β 2 signaling blockade and anti-CTLA-4 antibody immunotherapy in a murine BRAF ^{V600E} -PTEN ^{-/-} transgenic model of melanoma.. Journal of Clinical Oncology, 2014, 32, 3011-3011. | 0.8 | 25 |
| 39 | Abstract 2674: Stroma biology identifies heparins as differentiating agents in neuroblastoma. , 2014, , . | | 0 |
| 40 | T β RIII/ β 2-arrestin2 regulates integrin α 5 β 1 trafficking, function, and localization in epithelial cells. Oncogene, 2013, 32, 1416-1427. | 2.6 | 26 |
| 41 | A phase I study of ABT \times 510 plus bevacizumab in advanced solid tumors. Cancer Medicine, 2013, 2, 316-324. | 1.3 | 31 |
| 42 | The type β III TGF β 2 receptor regulates filopodia formation via a Cdc42-mediated IRSp53 \times N-WASP interaction in epithelial cells. Biochemical Journal, 2013, 454, 79-89. | 1.7 | 16 |
| 43 | A Phase II Study of Capecitabine, Oxaliplatin, and Bevacizumab in the Treatment of Metastatic Esophagogastric Adenocarcinomas. Oncologist, 2013, 18, 271-272. | 1.9 | 38 |
| 44 | Type III TGF- β 2 receptor promotes FGF2-mediated neuronal differentiation in neuroblastoma. Journal of Clinical Investigation, 2013, 123, 4786-4798. | 3.9 | 36 |
| 45 | Emerging Roles of TGF- β 2 Co-receptors in Human Disease. , 2013, , 59-89. | | 1 |
| 46 | Type III TGF- β 2 receptor downregulation generates an immunotolerant tumor microenvironment. Journal of Clinical Investigation, 2013, 123, 3925-3940. | 3.9 | 94 |
| 47 | Abstract 5041: The type III TGF-beta receptor promotes FGF2-mediated neuronal differentiation in neuroblastoma.. , 2013, , . | | 0 |
| 48 | Abstract C61: Phase I Study of pazopanib in combination with the investigational hypoxia-targeted drug TH-302.. , 2013, , . | | 0 |
| 49 | Endoglin regulates PI3-kinase/Akt trafficking and signaling to alter endothelial capillary stability during angiogenesis. Molecular Biology of the Cell, 2012, 23, 2412-2423. | 0.9 | 41 |
| 50 | Endoglin mediates fibronectin/ α 5 β 1 integrin and TGF- β 2 pathway crosstalk in endothelial cells. EMBO Journal, 2012, 31, 3885-3900. | 3.5 | 73 |
| 51 | Endocardial cell epithelial-mesenchymal transformation requires Type III TGF β 2 receptor interaction with GIPC. Cellular Signalling, 2012, 24, 247-256. | 1.7 | 34 |
| 52 | A phase I study of bevacizumab, everolimus and panitumumab in advanced solid tumors. Cancer Chemotherapy and Pharmacology, 2012, 70, 95-102. | 1.1 | 40 |
| 53 | Effect of the loss of the type III TGF β 2 receptor during tumor progression on tumor microenvironment: Preclinical development of TGF β 2 inhibition and TGF β 2-related biomarkers to enhance immunotherapy efficacy.. Journal of Clinical Oncology, 2012, 30, 10563-10563. | 0.8 | 3 |
| 54 | Effect of the vaccine Ad5 [E1-, E2b-]-CEA(6D) on CEA-directed CMI responses in patients with advanced CEA-expressing malignancies in a phase I/II clinical trial.. Journal of Clinical Oncology, 2012, 30, 2585-2585. | 0.8 | 38 |

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 55 | Abstract 3548: Loss of the type III TGF- β 2 receptor during cancer progression generates an immunotolerant tumor microenvironment: Translational implications for TGF- β 2 inhibition and immunotherapy biomarker development. , 2012, , . | | 0 |
| 56 | Abstract 3035: Bone morphogenetic proteins signal through Smad2 and Smad3 to regulate cell migration and proliferation. , 2012, , . | | 0 |
| 57 | Type III TGF- β 2 Receptor Enhances Colon Cancer Cell Migration and Anchorage-Independent Growth. Neoplasia, 2011, 13, 758-IN28. | 2.3 | 56 |
| 58 | A Phase II Study of Oxaliplatin, Dose-intense Capecitabine, and High-dose Bevacizumab in the Treatment of Metastatic Colorectal Cancer. Clinical Colorectal Cancer, 2011, 10, 210-216. | 1.0 | 10 |
| 59 | A phase I study of bevacizumab (B) in combination with everolimus (E) and erlotinib (E) in advanced cancer (BEE). Cancer Chemotherapy and Pharmacology, 2011, 67, 465-474. | 1.1 | 30 |
| 60 | A Phase II Trial of Bevacizumab plus Everolimus for Patients with Refractory Metastatic Colorectal Cancer. Oncologist, 2011, 16, 1131-1137. | 1.9 | 58 |
| 61 | BMP-2 and TGF β 2 Shared Pathways Regulate Endocardial Cell Transformation. Cells Tissues Organs, 2011, 194, 1-12. | 1.3 | 30 |
| 62 | The type III transforming growth factor- β 2 receptor inhibits proliferation, migration, and adhesion in human myeloma cells. Molecular Biology of the Cell, 2011, 22, 1463-1472. | 0.9 | 48 |
| 63 | Mechanical Stiffness Grades Metastatic Potential in Patient Tumor Cells and in Cancer Cell Lines. Cancer Research, 2011, 71, 5075-5080. | 0.4 | 597 |
| 64 | Molecular Characterization of the Tumor-Suppressive Function of Nischarin in Breast Cancer. Journal of the National Cancer Institute, 2011, 103, 1513-1528. | 3.0 | 54 |
| 65 | Phase I study of dasatinib in combination with capecitabine, oxaliplatin, and bevacizumab followed by an expanded cohort in previously untreated metastatic colorectal cancer.. Journal of Clinical Oncology, 2011, 29, 3586-3586. | 0.8 | 3 |
| 66 | A phase II study of capecitabine, oxaliplatin, bevacizumab and cetuximab in the treatment of metastatic colorectal cancer. Anticancer Research, 2011, 31, 255-61. | 0.5 | 22 |
| 67 | Deep sequencing of the small RNA transcriptome of normal and malignant human B cells identifies hundreds of novel microRNAs. Blood, 2010, 116, e118-e127. | 0.6 | 188 |
| 68 | Gfi-1B controls human erythroid and megakaryocytic differentiation by regulating TGF- β 2 signaling at the bipotent erythro-megakaryocytic progenitor stage. Blood, 2010, 115, 2784-2795. | 0.6 | 73 |
| 69 | Roles for the type III TGF- β 2 receptor in human cancer. Cellular Signalling, 2010, 22, 1163-1174. | 1.7 | 154 |
| 70 | Loss of type III transforming growth factor- β 2 receptor expression is due to methylation silencing of the transcription factor GATA3 in renal cell carcinoma. Oncogene, 2010, 29, 2905-2915. | 2.6 | 41 |
| 71 | The type III TGF- β 2 receptor suppresses breast cancer progression through GIPC-mediated inhibition of TGF- β 2 signaling. Carcinogenesis, 2010, 31, 175-183. | 1.3 | 57 |
| 72 | ALK5 phosphorylation of the endoglin cytoplasmic domain regulates Smad1/5/8 signaling and endothelial cell migration. Carcinogenesis, 2010, 31, 435-441. | 1.3 | 63 |

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 73 | Abstract 3971: The type III $\text{tgf-}\hat{\text{T}}^2$ receptor mediates bmp signaling in normal and cancerous mammary epithelial cells. , 2010, , . | | 0 |
| 74 | A phase I dose-escalation study of imatinib mesylate (Gleevec/STI571) plus capecitabine (Xeloda) in advanced solid tumors. <i>Anticancer Research</i> , 2010, 30, 1251-6. | 0.5 | 3 |
| 75 | Casein kinase $2\hat{\text{T}}^2$ as a novel enhancer of activin $\hat{\text{A}}$ -like receptor $\hat{\text{A}}^1$ signaling. <i>FASEB Journal</i> , 2009, 23, 3712-3721. | 0.2 | 26 |
| 76 | Bone morphogenetic proteins induce pancreatic cancer cell invasiveness through a Smad1-dependent mechanism that involves matrix metalloproteinase-2. <i>Carcinogenesis</i> , 2009, 30, 238-248. | 1.3 | 108 |
| 77 | The type III TGF- $\hat{\text{T}}^2$ receptor regulates directional migration: New tricks for an old dog. <i>Cell Cycle</i> , 2009, 8, 3069-3070. | 1.3 | 18 |
| 78 | The type III TGF- $\hat{\text{T}}^2$ receptor regulates epithelial and cancer cell migration through $\hat{\text{T}}^2$ -arrestin2-mediated activation of Cdc42. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2009, 106, 8221-8226. | 3.3 | 129 |
| 79 | The type III transforming growth factor- $\hat{\text{A}}$ receptor negatively regulates nuclear factor kappa B signaling through its interaction with $\hat{\text{A}}$ -arrestin2. <i>Carcinogenesis</i> , 2009, 30, 1281-1287. | 1.3 | 52 |
| 80 | The Transforming Growth Factor- $\hat{\text{T}}^2$ Type III Receptor Mediates Distinct Subcellular Trafficking and Downstream Signaling of Activin-like Kinase (ALK)3 and ALK6 Receptors. <i>Molecular Biology of the Cell</i> , 2009, 20, 4362-4370. | 0.9 | 37 |
| 81 | Proteoglycan signaling co-receptors: Roles in cell adhesion, migration and invasion. <i>Cellular Signalling</i> , 2009, 21, 1548-1558. | 1.7 | 123 |
| 82 | TGF- $\hat{\text{T}}^2$ Signaling. , 2009, , 137-149. | | 0 |
| 83 | Role of transforming growth factor- $\hat{\text{T}}^2$ superfamily signaling pathways in human disease. <i>Biochimica Et Biophysica Acta - Molecular Basis of Disease</i> , 2008, 1782, 197-228. | 1.8 | 544 |
| 84 | Endocytosis of the Type III Transforming Growth Factor- $\hat{\text{T}}^2$ (TGF- $\hat{\text{T}}^2$) Receptor through the Clathrin-independent/Lipid Raft Pathway Regulates TGF- $\hat{\text{T}}^2$ Signaling and Receptor Down-regulation. <i>Journal of Biological Chemistry</i> , 2008, 283, 34808-34818. | 1.6 | 57 |
| 85 | Expression of the type III TGF- $\hat{\text{T}}^2$ receptor is negatively regulated by TGF- $\hat{\text{T}}^2$. <i>Carcinogenesis</i> , 2008, 29, 905-912. | 1.3 | 47 |
| 86 | Bone Morphogenetic Proteins Signal through the Transforming Growth Factor- $\hat{\text{T}}^2$ Type III Receptor. <i>Journal of Biological Chemistry</i> , 2008, 283, 7628-7637. | 1.6 | 161 |
| 87 | $\hat{\text{T}}^2$ RIII suppresses non-small cell lung cancer invasiveness and tumorigenicity. <i>Carcinogenesis</i> , 2008, 29, 528-535. | 1.3 | 110 |
| 88 | Endoglin Promotes Transforming Growth Factor $\hat{\text{T}}^2$ -mediated Smad 1/5/8 Signaling and Inhibits Endothelial Cell Migration through Its Association with GIPC. <i>Journal of Biological Chemistry</i> , 2008, 283, 32527-32533. | 1.6 | 104 |
| 89 | Loss of type III transforming growth factor $\hat{\text{T}}^2$ receptor expression increases motility and invasiveness associated with epithelial to mesenchymal transition during pancreatic cancer progression. <i>Carcinogenesis</i> , 2008, 29, 252-262. | 1.3 | 119 |
| 90 | Initial results of a phase II study of oxaliplatin (OX), capecitabine (CAP), bevacizumab (BV), and cetuximab (CET) in the treatment of metastatic colorectal cancer (mCRC). <i>Journal of Clinical Oncology</i> , 2008, 26, 4063-4063. | 0.8 | 15 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|-----|-----------|
| 91 | The Type III Transforming Growth Factor (TGF)- β 2 Receptor Regulates Growth, Proliferation and Adhesion in Multiple Myeloma Cells. <i>Blood</i> , 2008, 112, 5158-5158. | 0.6 | 0 |
| 92 | The type III TGF- receptor signals through both Smad3 and the p38 MAP kinase pathways to contribute to inhibition of cell proliferation. <i>Carcinogenesis</i> , 2007, 28, 2491-2500. | 1.3 | 55 |
| 93 | Loss of Betaglycan Expression in Ovarian Cancer: Role in Motility and Invasion. <i>Cancer Research</i> , 2007, 67, 5231-5238. | 0.4 | 108 |
| 94 | The Interaction of Endoglin with β 2-Arrestin2 Regulates Transforming Growth Factor- β 2-mediated ERK Activation and Migration in Endothelial Cells. <i>Journal of Biological Chemistry</i> , 2007, 282, 21507-21517. | 1.6 | 83 |
| 95 | The Type III Transforming Growth Factor- β 2 Receptor as a Novel Tumor Suppressor Gene in Prostate Cancer. <i>Cancer Research</i> , 2007, 67, 1090-1098. | 0.4 | 167 |
| 96 | A Phase I study of capecitabine, carboplatin, and paclitaxel with external beam radiation therapy for esophageal carcinoma. <i>International Journal of Radiation Oncology Biology Physics</i> , 2007, 67, 1002-1007. | 0.4 | 11 |
| 97 | Bevacizumab, Oxaliplatin, and Capecitabine With Radiation Therapy in Rectal Cancer: Phase I Trial Results. <i>International Journal of Radiation Oncology Biology Physics</i> , 2007, 68, 472-478. | 0.4 | 135 |
| 98 | The type III TGF- β 2 receptor suppresses breast cancer progression. <i>Journal of Clinical Investigation</i> , 2007, 117, 206-217. | 3.9 | 212 |
| 99 | 281: The Role of the Type III TGF-BETA Receptor in Prostate Cancer. <i>Journal of Urology</i> , 2007, 177, 94-94. | 0.2 | 0 |
| 100 | Role of transforming growth factor- β 2 in hematologic malignancies. <i>Blood</i> , 2006, 107, 4589-4596. | 0.6 | 228 |
| 101 | Increased Toxicity With Gefitinib, Capecitabine, and Radiation Therapy in Pancreatic and Rectal Cancer: Phase I Trial Results. <i>Journal of Clinical Oncology</i> , 2006, 24, 656-662. | 0.8 | 134 |
| 102 | Cell-surface co-receptors: emerging roles in signaling and human disease. <i>Trends in Biochemical Sciences</i> , 2005, 30, 611-621. | 3.7 | 68 |
| 103 | Role of Transforming Growth Factor Beta in Human Cancer. <i>Journal of Clinical Oncology</i> , 2005, 23, 2078-2093. | 0.8 | 614 |
| 104 | Escape of Tumors From the Immune System. , 2004, , 85-95. | | 1 |
| 105 | Inhibiting the TGF- β 2 signalling pathway as a means of cancer immunotherapy. <i>Expert Opinion on Biological Therapy</i> , 2003, 3, 251-261. | 1.4 | 19 |
| 106 | Development of Human Protein Reference Database as an Initial Platform for Approaching Systems Biology in Humans. <i>Genome Research</i> , 2003, 13, 2363-2371. | 2.4 | 954 |
| 107 | β 2-Arrestin 2 Mediates Endocytosis of Type III TGF- β Receptor and Down-Regulation of Its Signaling. <i>Science</i> , 2003, 301, 1394-1397. | 6.0 | 246 |
| 108 | Inhibiting the TGF-beta signalling pathway as a means of cancer immunotherapy. <i>Expert Opinion on Biological Therapy</i> , 2003, 3, 251-261. | 1.4 | 25 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|------|-----------|
| 109 | Regulation of ALK-1 Signaling by the Nuclear Receptor LXR \hat{I}^2 . Journal of Biological Chemistry, 2002, 277, 50788-50794. | 1.6 | 25 |
| 110 | Context-specific Effects of Fibulin-5 (DANCE/EVEC) on Cell Proliferation, Motility, and Invasion. Journal of Biological Chemistry, 2002, 277, 27367-27377. | 1.6 | 141 |
| 111 | Functional Roles for the Cytoplasmic Domain of the Type III Transforming Growth Factor \hat{I}^2 Receptor in Regulating Transforming Growth Factor \hat{I}^2 Signaling. Journal of Biological Chemistry, 2001, 276, 24627-24637. | 1.6 | 123 |
| 112 | A Novel Mechanism for Regulating Transforming Growth Factor \hat{I}^2 (TGF- \hat{I}^2) Signaling. Journal of Biological Chemistry, 2001, 276, 39608-39617. | 1.6 | 169 |
| 113 | Two Patients With Sarcoma. Journal of Clinical Oncology, 2000, 18, 2343-2344. | 0.8 | 1 |
| 114 | Role of Transforming Growth Factor \hat{I}^2 in Human Disease. New England Journal of Medicine, 2000, 342, 1350-1358. | 13.9 | 2,264 |
| 115 | The p53 tumor suppressor gene in head and neck cancer. Current Opinion in Otolaryngology and Head and Neck Surgery, 1998, 6, 129-133. | 0.8 | 1 |
| 116 | Phosphorylation Specificities of Protein Kinase C Isozymes for Bovine Cardiac Troponin I and Troponin T and Sites within These Proteins and Regulation of Myofilament Properties. Journal of Biological Chemistry, 1996, 271, 23277-23283. | 1.6 | 163 |
| 117 | Protein Kinase C \hat{I}^{2II} Specifically Binds to and Is Activated by F-actin. Journal of Biological Chemistry, 1996, 271, 15823-15830. | 1.6 | 116 |
| 118 | Arachidonic acid and free fatty acids as second messengers and the role of protein kinase C. Cellular Signalling, 1995, 7, 171-184. | 1.7 | 221 |
| 119 | Regulation of protein kinase C and role in cancer biology. Cancer and Metastasis Reviews, 1994, 13, 411-431. | 2.7 | 234 |
| 120 | Alk1. The AFCS-nature Molecule Pages, 0, , . | 0.2 | 1 |
| 121 | TGF-beta type II receptor. The AFCS-nature Molecule Pages, 0, , . | 0.2 | 1 |
| 122 | TGF-beta type I receptor. The AFCS-nature Molecule Pages, 0, , . | 0.2 | 0 |