

Brooke Levis

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4459519/publications.pdf>

Version: 2024-02-01

68
papers

3,489
citations

257450

24
h-index

161849

54
g-index

70
all docs

70
docs citations

70
times ranked

4932
citing authors

#	ARTICLE	IF	CITATIONS
1	Accuracy of Patient Health Questionnaire-9 (PHQ-9) for screening to detect major depression: individual participant data meta-analysis. <i>BMJ: British Medical Journal</i> , 2019, 365, l1476.	2.3	822
2	Accuracy of the Edinburgh Postnatal Depression Scale (EPDS) for screening to detect major depression among pregnant and postpartum women: systematic review and meta-analysis of individual participant data. <i>BMJ, The</i> , 2020, 371, m4022.	6.0	298
3	Preferred reporting items for systematic review and meta-analysis of diagnostic test accuracy studies (PRISMA-DTA): explanation, elaboration, and checklist. <i>BMJ, The</i> , 2020, 370, m2632.	6.0	262
4	Accuracy of the PHQ-2 Alone and in Combination With the PHQ-9 for Screening to Detect Major Depression. <i>JAMA - Journal of the American Medical Association</i> , 2020, 323, 2290.	7.4	242
5	Equivalency of the diagnostic accuracy of the PHQ-8 and PHQ-9: a systematic review and individual participant data meta-analysis. <i>Psychological Medicine</i> , 2020, 50, 1368-1380.	4.5	175
6	Patient Health Questionnaire-9 scores do not accurately estimate depression prevalence: individual participant data meta-analysis. <i>Journal of Clinical Epidemiology</i> , 2020, 122, 115-128.e1.	5.0	113
7	Effects of screening for psychological distress on patient outcomes in cancer: A systematic review. <i>Journal of Psychosomatic Research</i> , 2013, 75, 1-17.	2.6	111
8	Does Evidence Support the American Heart Association's Recommendation to Screen Patients for Depression in Cardiovascular Care? An Updated Systematic Review. <i>PLoS ONE</i> , 2013, 8, e52654.	2.5	109
9	Depression screening and patient outcomes in pregnancy or postpartum: A systematic review. <i>Journal of Psychosomatic Research</i> , 2014, 76, 433-446.	2.6	88
10	Accuracy of the Hospital Anxiety and Depression Scale Depression subscale (HADS-D) to screen for major depression: systematic review and individual participant data meta-analysis. <i>BMJ, The</i> , 2021, 373, n972.	6.0	77
11	A systematic review of validated screening tools for anxiety disorders and PTSD in low to middle income countries. <i>BMC Psychiatry</i> , 2020, 20, 338.	2.6	73
12	The diagnostic accuracy of the Patient Health Questionnaire-2 (PHQ-2), Patient Health Questionnaire-8 (PHQ-8), and Patient Health Questionnaire-9 (PHQ-9) for detecting major depression: protocol for a systematic review and individual patient data meta-analyses. <i>Systematic Reviews</i> , 2014, 3, 124.	5.3	71
13	The Accuracy of the Patient Health Questionnaire-9 Algorithm for Screening to Detect Major Depression: An Individual Participant Data Meta-Analysis. <i>Psychotherapy and Psychosomatics</i> , 2020, 89, 25-37.	8.8	67
14	Comparison of depression prevalence estimates in meta-analyses based on screening tools and rating scales versus diagnostic interviews: a meta-research review. <i>BMC Medicine</i> , 2019, 17, 65.	5.5	64
15	Accuracy of the Patient Health Questionnaire-9 for screening to detect major depression: updated systematic review and individual participant data meta-analysis. <i>BMJ, The</i> , 2021, 375, n2183.	6.0	64
16	Psychosocial Aspects of Scleroderma. <i>Rheumatic Disease Clinics of North America</i> , 2015, 41, 519-528.	1.9	58
17	Association between antihypertensive treatment and adverse events: systematic review and meta-analysis. <i>BMJ, The</i> , 2021, 372, n189.	6.0	58
18	Probability of major depression diagnostic classification using semi-structured versus fully structured diagnostic interviews. <i>British Journal of Psychiatry</i> , 2018, 212, 377-385.	2.8	53

#	ARTICLE	IF	CITATIONS
19	Diagnostic accuracy of the Edinburgh Postnatal Depression Scale (EPDS) for detecting major depression in pregnant and postnatal women: protocol for a systematic review and individual patient data meta-analyses. <i>BMJ Open</i> , 2015, 5, e009742.	1.9	46
20	Selective Cutoff Reporting in Studies of Diagnostic Test Accuracy: A Comparison of Conventional and Individual-Patient-Data Meta-Analyses of the Patient Health Questionnaire-9 Depression Screening Tool. <i>American Journal of Epidemiology</i> , 2017, 185, 954-964.	3.4	45
21	Comparison of major depression diagnostic classification probability using the SCID, CIDI, and MINI diagnostic interviews among women in pregnancy or postpartum: An individual participant data meta-analysis. <i>International Journal of Methods in Psychiatric Research</i> , 2019, 28, e1803.	2.1	34
22	Completeness of Reporting of Systematic Reviews of Diagnostic Test Accuracy Based on the PRISMA-DTA Reporting Guideline. <i>Clinical Chemistry</i> , 2019, 65, 291-301.	3.2	33
23	Probability of major depression diagnostic classification based on the SCID, CIDI and MINI diagnostic interviews controlling for Hospital Anxiety and Depression Scale "Depression subscale scores: An individual participant data meta-analysis of 73 primary studies. <i>Journal of Psychosomatic Research</i> , 2020, 129, 109892.	2.6	33
24	CT/MRI and CEUS LI-RADS Major Features Association with Hepatocellular Carcinoma: Individual Patient Data Meta-Analysis. <i>Radiology</i> , 2022, 302, 326-335.	7.3	32
25	Depression prevalence based on the Edinburgh Postnatal Depression Scale compared to Structured Clinical Interview for DSM Disorders classification: Systematic review and individual participant data meta-analysis. <i>International Journal of Methods in Psychiatric Research</i> , 2021, 30, e1860.	2.1	30
26	Are MEDLINE searches sufficient for systematic reviews and meta-analyses of the diagnostic accuracy of depression screening tools? A review of meta-analyses. <i>Journal of Psychosomatic Research</i> , 2016, 87, 7-13.	2.6	29
27	Evaluation of Journal Registration Policies and Prospective Registration of Randomized Clinical Trials of Nonregulated Health Care Interventions. <i>JAMA Internal Medicine</i> , 2019, 179, 624.	5.1	25
28	Diagnostic accuracy of the Geriatric Depression Scale-30, Geriatric Depression Scale-15, Geriatric Depression Scale-5 and Geriatric Depression Scale-4 for detecting major depression: protocol for a systematic review and individual participant data meta-analysis. <i>BMJ Open</i> , 2018, 8, e026598.	1.9	24
29	Effects of a multi-faceted education and support programme on anxiety symptoms among people with systemic sclerosis and anxiety during COVID-19 (SPIN-CHAT): a two-arm parallel, partially nested, randomised, controlled trial. <i>Lancet Rheumatology</i> , The, 2021, 3, e427-e437.	3.9	24
30	Rates and correlates of sexual activity and impairment among women with systemic sclerosis. <i>Arthritis Care and Research</i> , 2012, 64, 340-350.	3.4	23
31	Diagnostic accuracy of the Depression subscale of the Hospital Anxiety and Depression Scale (HADS-D) for detecting major depression: protocol for a systematic review and individual patient data meta-analyses. <i>BMJ Open</i> , 2016, 6, e011913.	1.9	22
32	Protocol for a partially nested randomised controlled trial to evaluate the effectiveness of the scleroderma patient-centered intervention network COVID-19 home-isolation activities together (SPIN-CHAT) program to reduce anxiety among at-risk scleroderma patients. <i>Journal of Psychosomatic Research</i> , 2020, 135, 110132.	2.6	21
33	Validation of the COVID-19 Fears Questionnaires for Chronic Medical Conditions: A Scleroderma Patient-centered Intervention Network COVID-19 Cohort study. <i>Journal of Psychosomatic Research</i> , 2020, 139, 110271.	2.6	20
34	Probability of Major Depression Classification Based on the SCID, CIDI, and MINI Diagnostic Interviews: A Synthesis of Three Individual Participant Data Meta-Analyses. <i>Psychotherapy and Psychosomatics</i> , 2021, 90, 28-40.	8.8	20
35	Depression prevalence using the HADS-D compared to SCID major depression classification: An individual participant data meta-analysis. <i>Journal of Psychosomatic Research</i> , 2020, 139, 110256.	2.6	19
36	Sexual Activity and Impairment in Women with Systemic Sclerosis Compared to Women from a General Population Sample. <i>PLoS ONE</i> , 2012, 7, e52129.	2.5	17

#	ARTICLE	IF	CITATIONS
37	Shortening self-report mental health symptom measures through optimal test assembly methods: Development and validation of the Patient Health Questionnaire-Depression-4. <i>Depression and Anxiety</i> , 2019, 36, 82-92.	4.1	16
38	Impact of Reference Standard on CT, MRI, and Contrast-enhanced US LI-RADS Diagnosis of Hepatocellular Carcinoma: A Meta-Analysis. <i>Radiology</i> , 2022, 303, 544-545.	7.3	15
39	Selective cutoff reporting in studies of the accuracy of the Patient Health Questionnaire-9 and Edinburgh Postnatal Depression Scale: Comparison of results based on published cutoffs versus all cutoffs using individual participant data meta-analysis. <i>International Journal of Methods in Psychiatric Research</i> , 2021, 30, e1873.	2.1	12
40	Using Marital Status and Continuous Marital Satisfaction Ratings to Predict Depressive Symptoms in Married and Unmarried Women With Systemic Sclerosis: A Canadian Scleroderma Research Group Study. <i>Arthritis Care and Research</i> , 2016, 68, 1143-1149.	3.4	10
41	Factors associated with symptoms of depression among informal caregivers of people with systemic sclerosis: a cross-sectional study. <i>Disability and Rehabilitation</i> , 2020, 42, 394-399.	1.8	10
42	The association of sociodemographic and objectively-assessed disease variables with fatigue in systemic sclerosis: an analysis of 785 Canadian Scleroderma Research Group Registry patients. <i>Clinical Rheumatology</i> , 2017, 36, 373-379.	2.2	9
43	An empirical comparison of three methods for multiple cutoff diagnostic test meta-analysis of the Patient Health Questionnaire-9 (PHQ-9) depression screening tool using published data vs individual level data. <i>Research Synthesis Methods</i> , 2020, 11, 833-848.	8.7	9
44	Overestimation of Postpartum Depression Prevalence Based on a 5-item Version of the EPDS: Systematic Review and Individual Participant Data Meta-analysis. <i>Canadian Journal of Psychiatry</i> , 2020, 65, 835-844.	1.9	9
45	Comparison of different scoring methods based on latent variable models of the PHQ-9: an individual participant data meta-analysis. <i>Psychological Medicine</i> , 2022, 52, 3472-3483.	4.5	9
46	Data-driven methods distort optimal cutoffs and accuracy estimates of depression screening tools: a simulation study using individual participant data. <i>Journal of Clinical Epidemiology</i> , 2021, 137, 137-147.	5.0	9
47	Pain levels and associated factors in the Scleroderma Patient-centered Intervention Network (SPIN) cohort: a multicentre cross-sectional study. <i>Lancet Rheumatology</i> , The, 2021, 3, e844-e854.	3.9	9
48	Reliability and Validity of Three Versions of the Brief Fear of Negative Evaluation Scale in Patients With Systemic Sclerosis: A Scleroderma Patient-centered Intervention Network Cohort Study. <i>Arthritis Care and Research</i> , 2018, 70, 1646-1652.	3.4	8
49	Reporting of drug trial funding sources and author financial conflicts of interest in Cochrane and non-Cochrane meta-analyses: a cross-sectional study. <i>BMJ Open</i> , 2020, 10, e035633.	1.9	8
50	PRIME-IPD SERIES Part 2. Retrieving, checking, and harmonizing data are underappreciated challenges in individual participant data meta-analyses. <i>Journal of Clinical Epidemiology</i> , 2021, 136, 221-223.	5.0	5
51	Shortening the Edinburgh postnatal depression scale using optimal test assembly methods: Development of the EPDS-Dep-5. <i>Acta Psychiatrica Scandinavica</i> , 2021, 143, 348-362.	4.5	5
52	A comparison of bivariate, multivariate random-effects, and Poisson correlated gamma frailty models to meta-analyze individual patient data of ordinal scale diagnostic tests. <i>Biometrical Journal</i> , 2017, 59, 1317-1338.	1.0	4
53	The Scleroderma Patient-centered Intervention Network Self-Management (SPIN-SELF) Program: protocol for a two-arm parallel partially nested randomized controlled feasibility trial with progression to full-scale trial. <i>Trials</i> , 2021, 22, 856.	1.6	4
54	Validation of the Body Concealment Scale for Scleroderma (BCSS): Replication in the Scleroderma Patient-centered Intervention Network (SPIN) Cohort. <i>Body Image</i> , 2017, 20, 99-106.	4.3	3

#	ARTICLE	IF	CITATIONS
55	External validation of a shortened screening tool using individual participant data meta-analysis: A case study of the Patient Health Questionnaire-Dep-4. <i>Methods</i> , 2022, 204, 300-311.	3.8	3
56	Randomized feasibility trial of the Scleroderma Patient-centered Intervention Network Self-Management (SPIN-SELF) Program. <i>Pilot and Feasibility Studies</i> , 2022, 8, 45.	1.2	3
57	Sample size and precision of estimates in studies of depression screening tool accuracy: A meta-analysis research review of studies published in 2018-2021. <i>International Journal of Methods in Psychiatric Research</i> , 2022, 31, e1910.	2.1	3
58	Inclusion of currently diagnosed or treated individuals in studies of depression screening tool accuracy: a meta-research review of studies published in 2018-2021. <i>General Hospital Psychiatry</i> , 2022, 76, 25-30.	2.4	3
59	The Comparability of Functional Assessment of Chronic Illness Therapy - Fatigue Scores between Cancer and Systemic Sclerosis. <i>Journal of Scleroderma and Related Disorders</i> , 2017, 2, 57-63.	1.7	2
60	Reducing Waste and Increasing the Usability of Psychiatry Research: The Family of EQUATOR Reporting Guidelines and One of Its Newest Members: The PRISMA-DTA Statement. <i>Canadian Journal of Psychiatry</i> , 2018, 63, 509-512.	1.9	2
61	Intensity of care and perceived burden among informal caregivers to persons with chronic medical conditions: a systematic review and meta-analysis. <i>Disability and Rehabilitation</i> , 2022, 44, 6230-6246.	1.8	2
62	PHQ-8 scores and estimation of depression prevalence. <i>Lancet Public Health</i> , The, 2021, 6, e793.	10.0	2
63	A protocol for the VISION study: An individual patient data meta-analysis of randomised trials comparing MRI-targeted biopsy to standard transrectal ultrasound guided biopsy in the detection of prostate cancer. <i>PLoS ONE</i> , 2022, 17, e0263345.	2.5	2
64	Are Couple-Oriented Interventions Effective Across Chronic Illnesses? A Commentary on Martire et al.. <i>Annals of Behavioral Medicine</i> , 2011, 42, 134-135.	2.9	1
65	Reassessing the clinical utility of the Patient Health Questionnaire (PHQ)-9 for depression screening in prenatal women: a commentary on Sidebottom et al.. <i>Archives of Women's Mental Health</i> , 2013, 16, 253-254.	2.6	1
66	Resident Physicians With Depression or Depressive Symptoms. <i>JAMA - Journal of the American Medical Association</i> , 2016, 315, 2347.	7.4	1
67	THREE AUTHORS REPLY. <i>American Journal of Epidemiology</i> , 2017, 186, 895-895.	3.4	1
68	Group sample sizes in nonregulated health care intervention trials described as randomized controlled trials were overly similar. <i>Journal of Clinical Epidemiology</i> , 2020, 120, 8-16.	5.0	1