Michael Maeng

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/4459634/publications.pdf

Version: 2024-02-01

245 papers 13,120 citations

47409 49 h-index 28425 109 g-index

285 all docs

285 docs citations

times ranked

285

11576 citing authors

| # | Article | IF | Citations |
|----|--|------|-----------|
| 1 | Dual Antithrombotic Therapy with Dabigatran after PCI in Atrial Fibrillation. New England Journal of Medicine, 2017, 377, 1513-1524. | 13.9 | 1,099 |
| 2 | Thrombus Aspiration during ST-Segment Elevation Myocardial Infarction. New England Journal of Medicine, 2013, 369, 1587-1597. | 13.9 | 943 |
| 3 | Instantaneous Wave-free Ratio versus Fractional Flow Reserve to Guide PCI. New England Journal of Medicine, 2017, 376, 1813-1823. | 13.9 | 740 |
| 4 | Randomized Study on Simple Versus Complex Stenting of Coronary Artery Bifurcation Lesions. Circulation, 2006, 114, 1955-1961. | 1.6 | 666 |
| 5 | Percutaneous coronary angioplasty versus coronary artery bypass grafting in treatment of unprotected left main stenosis (NOBLE): a prospective, randomised, open-label, non-inferiority trial. Lancet, The, 2016, 388, 2743-2752. | 6.3 | 620 |
| 6 | System Delay and Mortality Among Patients With STEMI Treated With Primary Percutaneous Coronary Intervention. JAMA - Journal of the American Medical Association, 2010, 304, 763. | 3.8 | 519 |
| 7 | Duration of Triple Therapy in Patients Requiring Oral Anticoagulation After Drug-Eluting Stent Implantation. Journal of the American College of Cardiology, 2015, 65, 1619-1629. | 1.2 | 401 |
| 8 | ISAR-SAFE: a randomized, double-blind, placebo-controlled trial of 6 vs. 12 months of clopidogrel therapy after drug-eluting stenting. European Heart Journal, 2015, 36, 1252-1263. | 1.0 | 366 |
| 9 | Outcomes 1 Year after Thrombus Aspiration for Myocardial Infarction. New England Journal of Medicine, 2014, 371, 1111-1120. | 13.9 | 337 |
| 10 | 2-Year Clinical Outcomes After Implantation of Sirolimus-Eluting, Paclitaxel-Eluting, and Bare-Metal Coronary Stents. Journal of the American College of Cardiology, 2009, 53, 658-664. | 1.2 | 316 |
| 11 | Randomized Comparison of Final Kissing Balloon Dilatation Versus No Final Kissing Balloon Dilatation in Patients With Coronary Bifurcation Lesions Treated With Main Vessel Stenting. Circulation, 2011, 123, 79-86. | 1.6 | 269 |
| 12 | Routine Thrombectomy in Percutaneous Coronary Intervention for Acute ST-Segment–Elevation Myocardial Infarction. Circulation, 2006, 114, 40-47. | 1.6 | 242 |
| 13 | Stent Thrombosis, Myocardial Infarction, and Death After Drug-Eluting and Bare-Metal Stent Coronary Interventions. Journal of the American College of Cardiology, 2007, 50, 463-470. | 1.2 | 229 |
| 14 | Effect of remote ischaemic conditioning on clinical outcomes in patients with acute myocardial infarction (CONDI-2/ERIC-PPCI): a single-blind randomised controlled trial. Lancet, The, 2019, 394, 1415-1424. | 6.3 | 223 |
| 15 | Identification of vulnerable plaques and patients by intracoronary near-infrared spectroscopy and ultrasound (PROSPECT II): a prospective natural history study. Lancet, The, 2021, 397, 985-995. | 6.3 | 208 |
| 16 | Efficacy and safety of zotarolimus-eluting and sirolimus-eluting coronary stents in routine clinical care (SORT OUT III): a randomised controlled superiority trial. Lancet, The, 2010, 375, 1090-1099. | 6.3 | 198 |
| 17 | Biolimus-eluting biodegradable polymer-coated stent versus durable polymer-coated sirolimus-eluting stent in unselected patients receiving percutaneous coronary intervention (SORT OUT V): a randomised non-inferiority trial. Lancet, The, 2013, 381, 661-669. | 6.3 | 173 |
| 18 | Long-Term Results After Simple Versus Complex Stenting of Coronary Artery Bifurcation Lesions. Journal of the American College of Cardiology, 2013, 62, 30-34. | 1.2 | 168 |

| # | Article | IF | CITATIONS |
|----|--|-------------|-----------|
| 19 | Evaluation of Coronary Artery Stenosis by Quantitative Flow Ratio During Invasive Coronary Angiography. Circulation: Cardiovascular Imaging, 2018, 11, e007107. | 1.3 | 157 |
| 20 | Randomized Comparison of Coronary Bifurcation Stenting With the Crush Versus the Culotte Technique Using Sirolimus Eluting Stents. Circulation: Cardiovascular Interventions, 2009, 2, 27-34. | 1.4 | 156 |
| 21 | Randomized Comparison of Everolimus-Eluting and Sirolimus-Eluting Stents in Patients Treated With Percutaneous Coronary Intervention. Circulation, 2012, 125, 1246-1255. | 1.6 | 149 |
| 22 | Existing data sources for clinical epidemiology: The Western Denmark Heart Registry. Clinical Epidemiology, 2010, 2, 137. | 1.5 | 147 |
| 23 | Percutaneous Coronary Intervention for Vulnerable Coronary Atherosclerotic Plaque. Journal of the American College of Cardiology, 2020, 76, 2289-2301. | 1.2 | 123 |
| 24 | Safety and Efficacy of Everolimus-VersusÂSirolimus-Eluting Stents. Journal of the American College of Cardiology, 2016, 67, 751-762. | 1.2 | 116 |
| 25 | Safety of the Deferral of Coronary Revascularization on the Basis of Instantaneous Wave-Free Ratio and Fractional Flow Reserve Measurements in Stable Coronary Artery Disease and Acute Coronary Syndromes. JACC: Cardiovascular Interventions, 2018, 11, 1437-1449. | 1.1 | 111 |
| 26 | Zotarolimus-eluting durable-polymer-coated stent versus a biolimus-eluting biodegradable-polymer-coated stent in unselected patients undergoing percutaneous coronary intervention (SORT OUT VI): a randomised non-inferiority trial. Lancet, The, 2015, 385, 1527-1535. | 6. 3 | 107 |
| 27 | Effect of Ischemic Postconditioning During Primary Percutaneous Coronary Intervention for Patients With ST-Segment Elevation Myocardial Infarction. JAMA Cardiology, 2017, 2, 490. | 3.0 | 105 |
| 28 | Randomized Comparison of a Biodegradable Polymer Ultrathin Strut Sirolimus-Eluting Stent With a Biodegradable Polymer Biolimus-Eluting Stent in Patients Treated With Percutaneous Coronary Intervention. Circulation: Cardiovascular Interventions, 2016, 9, . | 1.4 | 104 |
| 29 | Coronary bifurcation lesions treated with simple or complex stenting: 5-year survival from patient-level pooled analysis of the Nordic Bifurcation Study and the British Bifurcation Coronary Study. European Heart Journal, 2016, 37, 1923-1928. | 1.0 | 103 |
| 30 | The EBC TWO Study (European Bifurcation Coronary TWO). Circulation: Cardiovascular Interventions, 2016, 9, . | 1.4 | 102 |
| 31 | Differential clinical outcomes after 1 year versus 5 years in a randomised comparison of zotarolimus-eluting and sirolimus-eluting coronary stents (the SORT OUT III study): a multicentre, open-label, randomised superiority trial. Lancet, The, 2014, 383, 2047-2056. | 6.3 | 96 |
| 32 | Serial Multimodality Imaging and 2-Year Clinical Outcomes of the NovelÂDESolve Novolimus-Eluting Bioresorbable Coronary Scaffold SystemÂfor the Treatment of Single DeÂNovo CoronaryÂLesions. JACC: Cardiovascular Interventions, 2016, 9, 565-574. | 1.1 | 91 |
| 33 | The Western Denmark Heart Registry. Journal of the American College of Cardiology, 2018, 71, 1259-1272. | 1.2 | 90 |
| 34 | Risk Associated With Surgery WithinÂ12ÂMonths After Coronary Drug-Eluting StentÂlmplantation. Journal of the American College of Cardiology, 2016, 68, 2622-2632. | 1.2 | 89 |
| 35 | Primary Angioplasty Versus Fibrinolysis in Acute Myocardial Infarction. Circulation, 2010, 121, 1484-1491. | 1.6 | 83 |
| 36 | Impact of Side Branch Modeling on Computation of Endothelial Shear Stress in Coronary Artery Disease. Journal of the American College of Cardiology, 2015, 66, 125-135. | 1.2 | 75 |

| # | Article | IF | CITATIONS |
|----|---|------------|----------------|
| 37 | Hypothermia during reperfusion does not reduce myocardial infarct size in pigs. Basic Research in Cardiology, 2006, 101, 61-68. | 2.5 | 72 |
| 38 | The Danish multicentre randomized study of fibrinolytic therapy vs. primary angioplasty in acute myocardial infarction (the DANAMI-2 trial): outcome after 3 years follow-up. European Heart Journal, 2007, 29, 1259-1266. | 1.0 | 71 |
| 39 | Outcomes after primary percutaneous coronary intervention in octogenarians and nonagenarians with STâ€segment elevation myocardial infarction: From the Western Denmark heart registry. Catheterization and Cardiovascular Interventions, 2013, 81, 912-919. | 0.7 | 68 |
| 40 | Lack of cardioprotection from subcutaneously and preischemic administered Liraglutide in a closed chest porcine ischemia reperfusion model. BMC Cardiovascular Disorders, 2009, 9, 31. | 0.7 | 65 |
| 41 | Side branch fractional flow reserve measurements after main vessel stenting: a Nordic-Baltic Bifurcation Study III substudy. EuroIntervention, 2012, 7, 1155-1161. | 1.4 | 59 |
| 42 | Negative vascular remodelling after implantation of bioabsorbable magnesium alloy stents in porcine coronary arteries: a randomised comparison with bare-metal and sirolimus-eluting stents. Heart, 2008, 95, 241-246. | 1,2 | 57 |
| 43 | Comparison of the Sirolimus-Eluting Versus Paclitaxel-Eluting Coronary Stent in Patients With Diabetes Mellitus: The Diabetes and Drug-Eluting Stent (DiabeDES) Randomized Angiography Trialâ€â€A list of participating centers and investigators appears in the Appendix American Journal of Cardiology, 2009, 103, 345-349. | 0.7 | 55 |
| 44 | Layered Fibrotic Plaques Are the Predominant Component in CardiacÂAllograft Vasculopathy. JACC: Cardiovascular Imaging, 2017, 10, 773-784. | 2.3 | 55 |
| 45 | Nonculprit Stenosis Evaluation Using Instantaneous Wave-Free Ratio in PatientsÂWith ST-Segment Elevation Myocardial Infarction. JACC: Cardiovascular Interventions, 2017, 10, 2528-2535. | 1.1 | 55 |
| 46 | Influence of Diabetes Mellitus on Clinical Outcomes Following Primary Percutaneous Coronary Intervention in Patients With ST-Segment Elevation Myocardial Infarction. American Journal of Cardiology, 2012, 109, 629-635. | 0.7 | 54 |
| 47 | Randomised comparison of manual compression and FemoSealª vascular closure device for closure after femoral artery access coronary angiography: the CLOSure dEvices Used in everyday Practice (CLOSE-UP) study. EuroIntervention, 2014, 10, 183-190. | 1.4 | 54 |
| 48 | System Delay and Timing of Intervention in Acute Myocardial Infarction (from the Danish Acute) Tj ETQq0 0 0 rg | BT /Overlo | ock 10 Tf 50 3 |
| 49 | Clinical Outcome After Crush Versus Culotte Stenting of Coronary Artery Bifurcation Lesions. JACC: Cardiovascular Interventions, 2013, 6, 1160-1165. | 1.1 | 51 |
| 50 | Comparison of Outcomes in Patients With Versus Without Diabetes Mellitus After Revascularization With Everolimus- and Sirolimus-Eluting Stents (from the SORT OUT IV Trial). American Journal of Cardiology, 2012, 110, 1585-1591. | 0.7 | 48 |
| 51 | Randomized Comparison of the Polymer-Free Biolimus-Coated BioFreedom Stent With the Ultrathin Strut Biodegradable Polymer Sirolimus-Eluting Orsiro Stent in an All-Comers Population Treated With Percutaneous Coronary Intervention. Circulation, 2020, 141, 2052-2063. | 1.6 | 48 |
| 52 | Time to Treatment and Three-Year Mortality After Primary Percutaneous Coronary Intervention for ST-Segment Elevation Myocardial Infarctionâ€"a DANish Trial in Acute Myocardial Infarction-2 (DANAMI-2) Substudy. American Journal of Cardiology, 2010, 105, 1528-1534. | 0.7 | 45 |
| 53 | Computed tomography derived fractional flow reserve testing in stable patients with typical angina pectoris: influence on downstream rate of invasive coronary angiography. European Heart Journal Cardiovascular Imaging, 2018, 19, 405-414. | 0.5 | 45 |
| 54 | Quantitative flow ratio for immediate assessment of nonculprit lesions in patients with STâ€segment elevation myocardial infarctionâ€"An iSTEMI substudy. Catheterization and Cardiovascular Interventions, 2019, 94, 686-692. | 0.7 | 45 |

| # | Article | IF | CITATIONS |
|----|--|-----|-----------|
| 55 | 3-Year Clinical Outcomes in the Randomized SORT OUT III Superiority Trial Comparing Zotarolimus- and Sirolimus-Eluting Coronary Stents. JACC: Cardiovascular Interventions, 2012, 5, 812-818. | 1.1 | 43 |
| 56 | Danish study of Non-Invasive testing in Coronary Artery Disease (Dan-NICAD): study protocol for a randomised controlled trial. Trials, 2016, 17, 262. | 0.7 | 43 |
| 57 | The SABRE Trial (Sirolimus Angioplasty Balloon forÂCoronary In-Stent Restenosis). JACC: Cardiovascular Interventions, 2017, 10, 2029-2037. | 1.1 | 43 |
| 58 | Intravascular ultrasound assessment of remodelling and reference segment plaque burden in type-2 diabetic patients. European Heart Journal, 2007, 28, 1759-1764. | 1.0 | 42 |
| 59 | 2-Year Patient-Related Versus Stent-Related Outcomes. Journal of the American College of Cardiology, 2012, 60, 1140-1147. | 1.2 | 42 |
| 60 | Paclitaxel and sirolimus eluting stents versus bare metal stents: long-term risk of stent thrombosis and other outcomes. From the Western Denmark Heart Registry. EuroIntervention, 2010, 5, 898-905. | 1.4 | 42 |
| 61 | Long-Term Outcomes After Percutaneous Coronary Intervention in Patients With and Without Diabetes Mellitus in Western Denmark. American Journal of Cardiology, 2010, 105, 1513-1519. | 0.7 | 41 |
| 62 | Co-registration of optical coherence tomography and X-ray angiography in percutaneous coronary intervention. The Does Optical Coherence Tomography Optimize Revascularization (DOCTOR) fusion study. International Journal of Cardiology, 2015, 182, 272-278. | 0.8 | 41 |
| 63 | Prospective, randomized trial of bioresorbable scaffolds vs. everolimus-eluting stents in patients undergoing coronary stenting for myocardial infarction: the Intracoronary Scaffold Assessment a Randomized evaluation of Absorb in Myocardial Infarction (ISAR-Absorb MI) trial. European Heart Journal, 2019, 40, 167-176. | 1.0 | 40 |
| 64 | Neointimal hyperplasia after sirolimus-eluting and paclitaxel-eluting stent implantation in diabetic patients: The Randomized Diabetes and Drug-Eluting Stent (DiabeDES) Intravascular Ultrasound Trial. European Heart Journal, 2008, 29, 2733-2741. | 1.0 | 39 |
| 65 | Outcome of Sirolimus-Eluting Versus Zotarolimus-Eluting Coronary Stent Implantation in Patients With and Without Diabetes Mellitus (a SORT OUT III Substudy). American Journal of Cardiology, 2011, 108, 1232-1237. | 0.7 | 39 |
| 66 | 16-year follow-up of the Danish Acute Myocardial Infarction 2 (DANAMI-2) trial: primary percutaneous coronary intervention vs. fibrinolysis in ST-segment elevation myocardial infarction. European Heart Journal, 2020, 41, 847-854. | 1.0 | 39 |
| 67 | Comparison of Durable-Polymer Zotarolimus-Eluting and Biodegradable-Polymer Biolimus-Eluting Coronary Stents in Patients With Coronary Artery Disease. JACC: Cardiovascular Interventions, 2017, 10, 255-264. | 1.1 | 38 |
| 68 | Patients With Diabetes Without Significant Angiographic Coronary Artery Disease Have the Same Risk of Myocardial Infarction as Patients Without Diabetes in a Real-World Population Receiving Appropriate Prophylactic Treatment. Diabetes Care, 2017, 40, 1103-1110. | 4.3 | 37 |
| 69 | Culprit only or multivessel percutaneous coronary interventions in patients with ST-segment elevation myocardial infarction and multivessel disease. EuroIntervention, 2012, 8, 456-464. | 1.4 | 37 |
| 70 | Clinical Validation of a Virtual Planner for Coronary Interventions Based on Coronary CT Angiography. JACC: Cardiovascular Imaging, 2022, 15, 1242-1255. | 2.3 | 36 |
| 71 | Randomised comparison of provisional side branch stenting versus a two-stent strategy for treatment of true coronary bifurcation lesions involving a large side branch: the Nordic-Baltic Bifurcation Study IV. Open Heart, 2020, 7, e000947. | 0.9 | 34 |
| 72 | Evaluation and Management of Nonculprit Lesions in STEMI. JACC: Cardiovascular Interventions, 2020, 13, 1145-1154. | 1.1 | 33 |

| # | Article | IF | CITATIONS |
|----|--|-----|-----------|
| 73 | Should the Presence or Extent of Coronary Artery Disease be Quantified in the CHA2DS2-VASc Score in Atrial Fibrillation? A Report from the Western Denmark Heart Registry. Thrombosis and Haemostasis, 2018, 118, 2162-2170. | 1.8 | 32 |
| 74 | Diabetes Mellitus Is Associated With Increased Risk of Ischemic Stroke in Patients With and Without Coronary Artery Disease. Stroke, 2019, 50, 3347-3354. | 1.0 | 32 |
| 75 | Cardiovascular risk and mortality in rheumatoid arthritis compared with diabetes mellitus and the general population. Rheumatology, 2021, 60, 1400-1409. | 0.9 | 32 |
| 76 | Comparison of Stent Thrombosis, Myocardial Infarction, and Mortality Following Drug-Eluting Versus Bare-Metal Stent Coronary Intervention in Patients With Diabetes Mellitus. American Journal of Cardiology, 2008, 102, 165-172. | 0.7 | 31 |
| 77 | Clinical Outcome After Primary Percutaneous Coronary Intervention With Drug-Eluting and Bare Metal Stents in Patients With ST-Segment Elevation Myocardial Infarction. Circulation: Cardiovascular Interventions, 2008, 1, 176-184. | 1.4 | 30 |
| 78 | Evaluation of algorithms for registry-based detection of acute myocardial infarction following percutaneous coronary intervention. Clinical Epidemiology, 2016, Volume 8, 415-423. | 1.5 | 30 |
| 79 | Development of heart failure in patients with rheumatoid arthritis: A Danish populationâ€based study. European Journal of Clinical Investigation, 2018, 48, e12915. | 1.7 | 30 |
| 80 | 5-Year Outcomes of PCI Guided by Measurement of Instantaneous Wave-Free Ratio Versus Fractional FlowÂReserve. Journal of the American College of Cardiology, 2022, 79, 965-974. | 1,2 | 30 |
| 81 | Lack of acute cardioprotective effect from preischaemic erythropoietin administration in a porcine coronary occlusion model. Clinical Physiology and Functional Imaging, 2005, 25, 305-310. | 0.5 | 28 |
| 82 | Three-Year Outcomes After Revascularization With Everolimus- andÂSirolimus-Eluting Stents From theÂSORT OUT IV Trial. JACC: Cardiovascular Interventions, 2014, 7, 840-848. | 1.1 | 28 |
| 83 | Comparison of Intravascular Ultrasound and Angiographic Assessment of Coronary Reference Segment Size in Patients With Type 2 Diabetes Mellitus. American Journal of Cardiology, 2008, 101, 590-595. | 0.7 | 27 |
| 84 | Clopidogrel discontinuation within the first year after coronary drug-eluting stent implantation: an observational study. BMC Cardiovascular Disorders, 2014, 14, 100. | 0.7 | 27 |
| 85 | Everolimus-Eluting Versus Biolimus-Eluting Stents With Biodegradable Polymers in UnselectedÂPatients Undergoing Percutaneous Coronary Intervention. JACC: Cardiovascular Interventions, 2019, 12, 624-633. | 1.1 | 27 |
| 86 | Validation of the DAPT score in patients randomized to 6 or 12 months clopidogrel after predominantly second-generation drug-eluting stents. Thrombosis and Haemostasis, 2017, 117, 1989-1999. | 1.8 | 26 |
| 87 | Randomized Clinical Comparison of the Dual-Therapy CD34 Antibody-Covered Sirolimus-Eluting Combo Stent With the Sirolimus-Eluting Orsiro Stent in Patients Treated With Percutaneous Coronary Intervention: The SORT OUT X Trial. Circulation, 2021, 143, 2155-2165. | 1.6 | 25 |
| 88 | A meta-analysis of specifically designed randomized trials of sirolimus-eluting versus paclitaxel-eluting stents in diabetic patients with coronary artery disease. American Heart Journal, 2011, 162, 740-747. | 1.2 | 24 |
| 89 | Outcome in high risk patients with unprotected left main coronary artery stenosis treated with percutaneous coronary intervention. Catheterization and Cardiovascular Interventions, 2010, 75, 101-108. | 0.7 | 23 |
| 90 | Coronary artery disease and risk of adverse cardiac events and stroke. European Journal of Clinical Investigation, 2017, 47, 819-828. | 1.7 | 23 |

| # | Article | IF | CITATIONS |
|-----|--|-----------------|--------------|
| 91 | Detection of early changes in the coronary artery microstructure after heart transplantation: A prospective optical coherence tomography study. Journal of Heart and Lung Transplantation, 2018, 37, 486-495. | 0.3 | 23 |
| 92 | Late lumen loss and intima hyperplasia after sirolimus-eluting and zotarolimus-eluting stent implantation in diabetic patients: the diabetes and drug-eluting stent (DiabeDES III) angiography and intravascular ultrasound trial. EuroIntervention, 2011, 7, 323-331. | 1.4 | 23 |
| 93 | Influence of multivessel disease with or without additional revascularization on mortality in patients with ST-segment elevation myocardial infarction. American Heart Journal, 2015, 170, 70-78. | 1.2 | 21 |
| 94 | Severe Mental Illness and Clinical Outcome After Primary Percutaneous Coronary Intervention. American Journal of Cardiology, 2017, 120, 550-555. | 0.7 | 21 |
| 95 | Association of Coronary Plaque With Low-Density Lipoprotein Cholesterol Levels and Rates of Cardiovascular Disease Events Among Symptomatic Adults. JAMA Network Open, 2022, 5, e2148139. | 2.8 | 21 |
| 96 | The impact of acquisition angle differences on threeâ€dimensional quantitative coronary angiography. Catheterization and Cardiovascular Interventions, 2011, 78, 214-222. | 0.7 | 20 |
| 97 | The natural history of collagen and α-actin expression after coronary angioplasty. Cardiovascular Pathology, 2004, 13, 260-267. | 0.7 | 19 |
| 98 | Rationale and design of The Intracoronary Stenting and Antithrombotic Regimen—Testing of a six-week versus a six-month clopidogrel treatment Regimen In Patients with concomitant aspirin and oraL anticoagulant therapy following drug-Eluting stenting (ISAR-TRIPLE) study. American Heart Journal, 2014, 167, 459-465.e1. | 1.2 | 19 |
| 99 | Timing, Causes, and Predictors of Death After Three Years' Follow-Up in the Danish Multicenter Randomized Study of Fibrinolysis Versus Primary Angioplasty in Acute Myocardial Infarction (DANAMI-2) Trial. American Journal of Cardiology, 2009, 104, 210-215. | 0.7 | 18 |
| 100 | CAD Is an Independent Risk Factor for Stroke Among Patients With AtrialÂFibrillation. Journal of the American College of Cardiology, 2018, 72, 2540-2542. | 1.2 | 18 |
| 101 | Comparison of Outcomes of Patients ≥80 Years of Age Having Percutaneous Coronary Intervention According to Presentation (Stable vs Unstable Angina Pectoris/Non–ST-Segment Elevation Myocardial) Tj ETQq. 1395-1400. | 1 1.9.7843 | 314 rgBT /0 |
| 102 | Randomized comparison of a sirolimus-eluting Orsiro stent with a biolimus-eluting Nobori stent in patients treated with percutaneous coronary intervention: Rationale and study design of the Scandinavian Organization for Randomized Trials with Clinical Outcome VII trial. American Heart Journal, 2015, 170, 210-215. | 1.2 | 17 |
| 103 | Two-year outcome after biodegradable polymer sirolimus- and biolimus-eluting coronary stents (from) Tj ETQq $1\ 1$ | 0,784314 1.4 | l rgBT /Over |
| 104 | The risk and prognostic impact of definite stent thrombosis or in-stent restenosis after coronary stent implantation. EuroIntervention, 2012, 8, 591-598. | 1.4 | 17 |
| 105 | Target lesion revascularisation in patients treated with a sirolimus-eluting or paclitaxel-eluting stent. Heart, 2007, 93, 694-697. | 1.2 | 16 |
| 106 | Serial Intravascular Ultrasound Analysis of Peri-Stent Remodeling and Proximal and Distal Edge Effects After Sirolimus-Eluting or Paclitaxel-Eluting Stent Implantation in Patients With Diabetes Mellitus. American Journal of Cardiology, 2009, 103, 1083-1088. | 0.7 | 16 |
| 107 | Assessing the Nationwide Impact of a Registry-Based Randomized Clinical Trial on Cardiovascular Practice. Circulation: Cardiovascular Interventions, 2019, 12, e007381. | 1.4 | 16 |
| 108 | Impact of rheumatoid arthritis on major cardiovascular events in patients with and without coronary artery disease. Annals of the Rheumatic Diseases, 2020, 79, 1182-1188. | 0.5 | 16 |

| # | Article | IF | CITATIONS |
|-----|---|---------------------|-------------|
| 109 | Peripheral artery disease, lower limb revascularization, and amputation in diabetes patients with and without coronary artery disease: a cohort study from the Western Denmark Heart Registry. BMJ Open Diabetes Research and Care, 2021, 9, e001803. | 1.2 | 16 |
| 110 | Interplay of Risk Factors and CoronaryÂArtery Calcium for CHD Risk inÂYoung Patients. JACC: Cardiovascular Imaging, 2021, 14, 2387-2396. | 2.3 | 16 |
| 111 | Development and validation of an artificial neural network algorithm to predict mortality and admission to hospital for heart failure after myocardial infarction: a nationwide population-based study. The Lancet Digital Health, 2022, 4, e37-e45. | 5.9 | 16 |
| 112 | Quantitative coronary analysis in the Nordic Bifurcation studies. International Journal of Cardiovascular Imaging, 2011, 27, 175-180. | 0.7 | 15 |
| 113 | Long-Term Outcome of Sirolimus-Eluting and Zotarolimus-Eluting Coronary Stent Implantation in Patients With and Without Diabetes Mellitus (A Danish Organization for Randomized Trials on) Tj ETQq1 1 0.7843 | 3 b4 7rgBT / | Oværlock 10 |
| 114 | Dual anti-platelet therapy after coronary drug-eluting stent implantation and surgery-associated major adverse events. Thrombosis and Haemostasis, 2016, 116, 172-180. | 1.8 | 15 |
| 115 | Risk stratification by assessment of coronary artery disease using coronary computed tomography angiography in diabetes and non-diabetes patients: a study from the Western Denmark Cardiac Computed Tomography Registry. European Heart Journal Cardiovascular Imaging, 2019, 20, 1271-1278. | 0.5 | 15 |
| 116 | SARS-CoV-2 infection and adverse outcomes in users of ACE inhibitors and angiotensin-receptor blockers: a nationwide case-control and cohort analysis. Thorax, 2021, 76, 370-379. | 2.7 | 15 |
| 117 | Intravascular Ultrasound Assessment of Expansion of the Sirolimus-Eluting (Cypher Select) and Paclitaxel-Eluting (Taxus Express-2) Stent in Patients With Diabetes Mellitus. American Journal of Cardiology, 2008, 102, 19-26. | 0.7 | 14 |
| 118 | Concomitant use of clopidogrel and statins and risk of major adverse cardiovascular events following coronary stent implantation. British Journal of Clinical Pharmacology, 2012, 74, 161-170. | 1.1 | 14 |
| 119 | Use of clopidogrel and calcium channel blockers and risk of major adverse cardiovascular events. European Journal of Clinical Investigation, 2012, 42, 266-274. | 1.7 | 14 |
| 120 | Six Versus Twelve Months Clopidogrel Therapy After Drug-Eluting Stenting in Patients With Acute Coronary Syndrome: An ISAR-SAFE Study Subgroup Analysis. Scientific Reports, 2016, 6, 33054. | 1.6 | 14 |
| 121 | Editor's Choice-Acute versus subacute angiography in patients with non-ST-elevation myocardial infarction – the NONSTEMI trial phase I. European Heart Journal: Acute Cardiovascular Care, 2017, 6, 490-499. | 0.4 | 14 |
| 122 | Rationale and design of the precise percutaneous coronary intervention plan (<scp>P3</scp>) study: Prospective evaluation of a virtual computed tomographyâ€based percutaneous intervention planner. Clinical Cardiology, 2021, 44, 446-454. | 0.7 | 14 |
| 123 | Nationwide Trends in Cardiac Risk and Mortality in Patients With Incident Type 2 Diabetes: A Danish Cohort Study. Diabetes Care, 2021, 44, 2353-2360. | 4.3 | 14 |
| 124 | Angiographic and clinical outcomes of STEMI patients treated with bioresorbable or metallic everolimus-eluting stents: a pooled analysis of individual patient data. EuroIntervention, 2020, 15, 1451-1457. | 1.4 | 14 |
| 125 | Adventitial Myofibroblasts Play no Major Role in Neointima Formation After Angioplasty. Scandinavian Cardiovascular Journal, 2003, 37, 34-42. | 0.4 | 13 |
| 126 | Incidence of definite stent thrombosis or inâ€stent restenosis after drugâ€eluting stent implantation for treatment of coronary inâ€stent restenosis: From Western Denmark heart registry. Catheterization and Cardiovascular Interventions, 2013, 81, 260-265. | 0.7 | 13 |

| # | Article | IF | CITATIONS |
|-----|---|-----|-----------|
| 127 | A 10â€month angiographic and 4â€year clinical outcome of everolimusâ€eluting versus sirolimusâ€eluting coronary stents in patients with diabetes mellitus (the diabedES IV randomized angiography trial). Catheterization and Cardiovascular Interventions, 2015, 86, 1161-1167. | 0.7 | 13 |
| 128 | <p>Extent of coronary artery disease is associated with myocardial infarction and mortality in patients with diabetes mellitus</p> . Clinical Epidemiology, 2019, Volume 11, 419-428. | 1.5 | 13 |
| 129 | Dabigatran Dual Therapy Versus Warfarin Triple Therapy Post–PCI in Patients WithÂAtrial Fibrillation and Diabetes. JACC: Cardiovascular Interventions, 2019, 12, 2346-2355. | 1.1 | 13 |
| 130 | Nurse-led Motivational Telephone Follow-up After Same-day Percutaneous Coronary Intervention Reduces Readmission and Contacts to General Practice. Journal of Cardiovascular Nursing, 2019, 34, 222-230. | 0.6 | 13 |
| 131 | Myocardial perfusion imaging with 99mTc sestamibi early after reperfusion reliably reflects infarct size reduction by ischaemic preconditioning in an experimental porcine model. Nuclear Medicine Communications, 2004, 25, 495-500. | 0.5 | 12 |
| 132 | Pain and discomfort in closure of femoral access coronary angiography. The CLOSuredEvices Used in everyday Practice (CLOSE-UP) pain sub study. European Journal of Cardiovascular Nursing, 2014, 13, 221-226. | 0.4 | 12 |
| 133 | Randomized clinical comparison of the dual-therapy CD34 antibody-covered sirolimus-eluting Combo stent with the sirolimus-eluting Orsiro stent in patients treated with percutaneous coronary intervention: Rationale and study design of the Scandinavian Organization for Randomized Trials with Clinical Outcome (SORT OUT) X trial. American Heart Journal, 2018, 202, 49-53. | 1.2 | 12 |
| 134 | Validation of the European Society of Cardiology and European Society of Anaesthesiology non-cardiac surgery risk score in patients treated with coronary drug-eluting stent implantation. European Heart Journal Quality of Care & Dinical Outcomes, 2019, 5, 22-27. | 1.8 | 12 |
| 135 | External applicability of the COMPASS trial: the Western Denmark Heart Registry. European Heart Journal - Cardiovascular Pharmacotherapy, 2019, 5, 192-199. | 1.4 | 12 |
| 136 | Estimated Pulse Wave Velocity Is Associated With Allâ€Cause Mortality During 8.5 Years Followâ€up in Patients Undergoing Elective Coronary Angiography. Journal of the American Heart Association, 2022, 11, e025173. | 1.6 | 12 |
| 137 | Longâ€ŧerm outcome following percutaneous coronary intervention with drugâ€eluting stents compared with bareâ€metal stents in saphenous vein graft lesions: From Western Denmark heart registry. Catheterization and Cardiovascular Interventions, 2014, 83, 1035-1042. | 0.7 | 11 |
| 138 | Regional systems-of-care for primary percutaneous coronary intervention in ST-elevation myocardial infarction. Coronary Artery Disease, 2015, 26, 713-722. | 0.3 | 11 |
| 139 | Impact of diabetes on clinical outcomes after revascularization with sirolimusâ€eluting and biolimusâ€eluting stents with biodegradable polymer from the SORT OUT VII trial. Catheterization and Cardiovascular Interventions, 2019, 93, 567-573. | 0.7 | 11 |
| 140 | Interaction of ischaemic postconditioning and thrombectomy in patients with ST-elevation myocardial infarction. Heart, 2020, 106, 24-32. | 1.2 | 11 |
| 141 | Final five-year outcomes after implantation of biodegradable polymer-coated biolimus-eluting stents versus durable polymer-coated sirolimus-eluting stents. EuroIntervention, 2017, 13, 1336-1344. | 1.4 | 11 |
| 142 | Effects of Levosimendan on Myocardial Infarct Size and Hemodynamics in a Closed-Chest Porcine Ischemia–Reperfusion Model. Cardiovascular Drugs and Therapy, 2006, 20, 335-342. | 1.3 | 10 |
| 143 | Effects of pentoxifylline on the vascular response to injury after angioplasty in rabbit iliac arteries. Basic Research in Cardiology, 2008, 103, 257-264. | 2.5 | 10 |
| 144 | Zotarolimusâ€eluting vs. sirolimusâ€eluting coronary stents in patients with and without acute coronary syndromes: a SORT OUT III substudy. European Journal of Clinical Investigation, 2012, 42, 1047-1054. | 1.7 | 10 |

| # | Article | IF | Citations |
|-----|---|-----|-----------|
| 145 | Safety of therapeutic hypothermia combined with primary percutaneous coronary intervention after out-of-hospital cardiac arrest. European Heart Journal: Acute Cardiovascular Care, 2015, 4, 60-63. | 0.4 | 10 |
| 146 | Impact of thrombus aspiration during ST-Elevation Myocardial Infarction: a six month composite endpoint and risk of stroke analyses of the TASTE trial. BMC Cardiovascular Disorders, 2016, 16, 62. | 0.7 | 10 |
| 147 | Comparison of Acute Versus Subacute Coronary Angiography in Patients With NON-ST-Elevation Myocardial Infarction (from the NONSTEMI Trial). American Journal of Cardiology, 2019, 124, 825-832. | 0.7 | 10 |
| 148 | Comparison of the polymer-free biolimus-coated BioFreedom stent with the thin-strut biodegradable polymer sirolimus-eluting Orsiro stent in an all-comers population treated with percutaneous coronary intervention: Rationale and design of the randomized SORT OUT IX trial. American Heart Journal, 2019, 213, 1-7. | 1,2 | 10 |
| 149 | Comparison of Frequency of Ischemic Stroke in Patients With Versus Without Coronary Heart Disease and Without Atrial Fibrillation. American Journal of Cardiology, 2019, 123, 153-158. | 0.7 | 10 |
| 150 | Smoking is the dominating modifiable risk factor in younger patients with STEMI. European Heart Journal: Acute Cardiovascular Care, 2020, 9, 70-75. | 0.4 | 9 |
| 151 | Clinical outcomes three-year after revascularization with biodegradable polymer stents: ultrathin-strut sirolimus-eluting stent versus biolimus-eluting stent: from the Scandinavian organization for randomized trials with clinical outcome VII trial. Coronary Artery Disease, 2020, 31, 485-492. | 0.3 | 9 |
| 152 | Design and Rationale of a Randomized Trial of COBRA PzF Stenting to REDUCE Duration of Triple Therapy (COBRA-REDUCE). Cardiovascular Revascularization Medicine, 2022, 34, 17-24. | 0.3 | 9 |
| 153 | Use of three-dimensional optical coherence tomography to verify correct wire position in a jailed side branch after main vessel stent implantation. EuroIntervention, 2011, 7, 528-529. | 1.4 | 9 |
| 154 | Comparison of MynxGrip vascular closure device and manual compression for closure after femoral access angiography: a randomized controlled trial: the closure devices used in every day practice study, CLOSE-UP III trial. BMC Cardiovascular Disorders, 2022, 22, 68. | 0.7 | 9 |
| 155 | Nonsteroidal Antiinflammatory Drug Use and Cardiovascular Risks After Coronary Stent Implantation. Pharmacotherapy, 2011, 31, 458-468. | 1.2 | 8 |
| 156 | Intimal hyperplasia and vascular remodeling after everolimusâ€eluting and sirolimusâ€eluting stent implantation in diabetic patients: The randomized diabetes and drugâ€eluting stent (DiabeDES) IV intravascular ultrasound trial. Catheterization and Cardiovascular Interventions, 2014, 83, 864-872. | 0.7 | 8 |
| 157 | Heart Failure and Ischemic Heart Disease in Patients With Rheumatoid Arthritis. Journal of the American College of Cardiology, 2017, 70, 3069-3071. | 1.2 | 8 |
| 158 | Association between anti-diabetes treatments and cardiovascular risk in diabetes patients with and without coronary artery disease. Diabetes and Vascular Disease Research, 2019, 16, 351-359. | 0.9 | 8 |
| 159 | Outcomes after revascularisation with everolimus- and sirolimus-eluting stents in patients with acute coronary syndromes and stable angina pectoris: a substudy of the SORT OUT IV trial. EuroIntervention, 2014, 10, 212-223. | 1.4 | 8 |
| 160 | Dual antithrombotic treatment in chronic coronary syndrome: European Society of Cardiology criteria vs. CHADS-P2A2RC score. European Heart Journal, 2022, 43, 996-1004. | 1.0 | 8 |
| 161 | Myocardial cooling for reperfusion injury protection achieved by organ specific hypothermic autologous perfusion. Scandinavian Cardiovascular Journal, 2003, 37, 297-303. | 0.4 | 7 |
| 162 | Positive Predictive Value of Clinically Suspected ST-Segment Elevation Myocardial Infarction Using Angiographic Verification. American Journal of Cardiology, 2013, 112, 923-927. | 0.7 | 7 |

| # | Article | IF | CITATIONS |
|-----|--|-----|-----------|
| 163 | A Novel Model for Prediction of Thromboembolic and Cardiovascular Events in Patients Without Atrial Fibrillation. American Journal of Cardiology, 2020, 131, 40-48. | 0.7 | 7 |
| 164 | Renin–Angiotensin System Blockers and Adverse Outcomes of Influenza and Pneumonia: A Danish Cohort Study. Journal of the American Heart Association, 2020, 9, e017297. | 1.6 | 7 |
| 165 | Polymer-free biolimus-coated stents versus ultrathin-strut biodegradable polymer sirolimus-eluting stents: two-year outcomes of the randomised SORT OUT IX trial. EuroIntervention, 2022, 18, e124-e131. | 1.4 | 7 |
| 166 | Lack of cardioprotection from metabolic support with glutamine or glutamate in a porcine coronary occlusion model. Scandinavian Cardiovascular Journal, 2005, 39, 115-120. | 0.4 | 6 |
| 167 | Staged re-evaluation of non-culprit lesions in ST segment elevation myocardial infarction: a retrospective study. Open Heart, 2016, 3, e000427. | 0.9 | 6 |
| 168 | Everolimus-Eluting Versus Biolimus-Eluting Coronary Stent Implantation in Patients With and Without Diabetes Mellitus. American Journal of Cardiology, 2019, 124, 671-677. | 0.7 | 6 |
| 169 | Ten-year cardiovascular risk in diabetes patients without obstructive coronary artery disease: a retrospective Western Denmark cohort study. Cardiovascular Diabetology, 2021, 20, 23. | 2.7 | 6 |
| 170 | Ten-year outcomes from a randomised comparison of zotarolimus-eluting and sirolimus-eluting stents: the SORT OUT III study. EuroIntervention, 2019, 15, e1022-e1024. | 1.4 | 6 |
| 171 | Effectiveness and Safety of Ticagrelor Implementation in Patients with Acute Coronary Syndrome undergoing Percutaneous Coronary Intervention: A Cohort Study in Western Denmark. Lancet Regional Health - Europe, The, 2022, 14, 100301. | 3.0 | 6 |
| 172 | Coronary Edema Demonstrated by Cardiovascular Magnetic Resonance in Patients With Peri-Stent Inflammation and Aneurysm Formation After Treatment by Drug-Eluting Stents. Circulation: Cardiovascular Imaging, 2013, 6, 352-354. | 1.3 | 5 |
| 173 | Gastroscopy-related adverse cardiac events and bleeding complications among patients treated with coronary stents and dual antiplatelet therapy. Endoscopy International Open, 2016, 04, E527-E533. | 0.9 | 5 |
| 174 | Myocardial infarction with non-obstructive thrombus validated by optical coherence tomography. Scandinavian Cardiovascular Journal, 2017, 51, 61-68. | 0.4 | 5 |
| 175 | Randomized comparison of sirolimus eluting, and biolimus eluting bioresorbable polymer stents: the SORT-OUT VII optical coherence tomography study. European Heart Journal Cardiovascular Imaging, 2018, 19, 329-338. | 0.5 | 5 |
| 176 | Procedural findings and early healing response after implantation of a self-apposing bioresorbable scaffold in coronary bifurcation lesions. International Journal of Cardiovascular Imaging, 2019, 35, 1199-1210. | 0.7 | 5 |
| 177 | Predicting stroke in patients without atrial fibrillation. European Journal of Clinical Investigation, 2019, 49, e13103. | 1.7 | 5 |
| 178 | Ten-Year Outcomes of Sirolimus-Eluting Versus Zotarolimus-Eluting Coronary Stents in Patients With Versus Without Diabetes Mellitus (SORT OUT III). American Journal of Cardiology, 2020, 125, 349-353. | 0.7 | 5 |
| 179 | Diabetes is not a risk factor for myocardial infarction in patients without coronary artery disease: A study from the Western Denmark Heart Registry. Diabetes and Vascular Disease Research, 2020, 17, 147916412094180. | 0.9 | 5 |
| 180 | Insulin-treated versus noninsulin-treated diabetes and risk of ischemic stroke in patients with atrial fibrillation. Vascular Pharmacology, 2021, 136, 106809. | 1.0 | 5 |

| # | Article | IF | Citations |
|-----|--|-------------|-----------|
| 181 | Evaluation of Dual Versus Triple Therapy by Landmark Analysis in the RE-DUAL PCIÂTrial. JACC: Cardiovascular Interventions, 2021, 14, 768-780. | 1.1 | 5 |
| 182 | Enhanced platelet inhibition by clopidogrel and risk of bleeding in patients requiring oral anticoagulation after drug-eluting stent implantation. EuroIntervention, 2019, 15, 700-706. | 1.4 | 5 |
| 183 | Ten-year patterns of stent thrombosis after percutaneous coronary intervention with new- versus early-generation drug-eluting stents: insights from the DECADE cooperation. Revista Espanola De Cardiologia (English Ed), 2022, , . | 0.4 | 5 |
| 184 | A phase of increased ST elevation during coronary occlusion following ischemic preconditioning. Basic Research in Cardiology, 2006, 101, 140-148. | 2. 5 | 4 |
| 185 | Deliver the drug and be resorbed: evidence from ABSORB Japan. European Heart Journal, 2015, 36, 3343-3345. | 1.0 | 4 |
| 186 | ST Elevation Infarction after Heart Transplantation Induced by Coronary Spasms and Mural Thrombus Detected by Optical Coherence Tomography. Case Reports in Transplantation, 2016, 2016, 1-4. | 0.1 | 4 |
| 187 | Agreement between nonculprit stenosis follow-up iFR and FFR after STEMI (iSTEMI substudy). BMC Research Notes, 2020, 13, 410. | 0.6 | 4 |
| 188 | Instantaneous wave-free ratio guided multivessel revascularisation during percutaneous coronary intervention for acute myocardial infarction: study protocol of the randomised controlled iMODERN trial. BMJ Open, 2021, 11, e044035. | 0.8 | 4 |
| 189 | Microvascular disease increases the risk of lower limb amputation – A Western Danish cohort study. European Journal of Clinical Investigation, 2022, 52, e13812. | 1.7 | 4 |
| 190 | Association between REDUCE-IT criteria, coronary artery disease severity, and cardiovascular events: the Western Denmark Heart Registry. European Journal of Preventive Cardiology, 2022, 29, 1802-1810. | 0.8 | 4 |
| 191 | Clinical use of optical coherence tomography to identify angiographic silent stent thrombosis. Scandinavian Cardiovascular Journal, 2014, 48, 156-160. | 0.4 | 3 |
| 192 | Reply. Journal of the American College of Cardiology, 2015, 66, 1089-1090. | 1.2 | 3 |
| 193 | Coronary stent implantation and adverse cardiac events after surgery. European Journal of Clinical Investigation, 2018, 48, e13030. | 1.7 | 3 |
| 194 | Risk of Myocardial Infarction in Patients Without Angiographic Coronary Artery Disease Compared With the General Population. American Journal of Cardiology, 2020, 132, 8-14. | 0.7 | 3 |
| 195 | Thirteen-year trends in cardiovascular risk in men and women with chronic coronary syndrome. European Heart Journal Quality of Care & Clinical Outcomes, 2022, 8, 437-446. | 1.8 | 3 |
| 196 | Five-year safety and performance data of a novel third-generation novolimus-eluting bioresorbable scaffold in single de novo lesions. EuroIntervention, 2019, 15, 685-687. | 1.4 | 3 |
| 197 | Cardiovascular risk in patients with and without diabetes presenting with chronic coronary syndrome in 2004–2016. BMC Cardiovascular Disorders, 2021, 21, 579. | 0.7 | 3 |
| 198 | Clinical outcomes of everolimus-eluting bioresorbable scaffolds or everolimus-eluting stents in patients with acute myocardial infarction: two-year results of the randomised ISAR-Absorb MI trial. EuroIntervention, 2022, 17, 1348-1351. | 1.4 | 3 |

| # | Article | IF | CITATIONS |
|-----|---|-----|-----------|
| 199 | Acute haemodynamic effects of erythropoietin alone and in combination with dopamine in a porcine model. Clinical Physiology and Functional Imaging, 2006, 26, 283-287. | 0.5 | 2 |
| 200 | Zotarolimus-eluting versus sirolimus-eluting coronary stent implantation. Interventional Cardiology, 2010, 2, 807-812. | 0.0 | 2 |
| 201 | Comparison of zotarolimus-eluting and sirolimus-eluting coronary stents: a study from the Western Denmark Heart Registry. BMC Cardiovascular Disorders, 2012, 12, 84. | 0.7 | 2 |
| 202 | TCT-433 Feasibility, self-correcting properties and one-month results after implantation of a novolimus eluting bioresorbable stent in coronary bifurcations. The BIFSORB pilot study. Journal of the American College of Cardiology, 2016, 68, B174-B175. | 1.2 | 2 |
| 203 | Perioperative myocardial infarction: in the twilight zone between surgery and cardiology. European Heart Journal, 2017, 38, 2418-2420. | 1.0 | 2 |
| 204 | Risk of Myocardial Infarction and Death After Noncardiac Surgery Performed Within the First Year After Coronary Drug-Eluting Stent Implantation for Acute Coronary Syndrome or Stable Angina Pectoris. American Journal of Cardiology, 2021, 160, 14-20. | 0.7 | 2 |
| 205 | EXCEL and NOBLE: stents or surgery for left main stem stenosis?. EuroIntervention, 2017, 13, e604-e608. | 1.4 | 2 |
| 206 | Impact of diabetes on clinical outcomes after revascularization with the dual therapy CD34 antibodyâ€covered sirolimusâ€eluting Combo stentÂand the sirolimusâ€eluting Orsiro stent. Catheterization and Cardiovascular Interventions, 2022, , . | 0.7 | 2 |
| 207 | OUP accepted manuscript. European Heart Journal, 2022, , . | 1.0 | 2 |
| 208 | TCT-297 Stent edge dissections detected by Optical Coherence Tomography, incidence, predictors and 12-month outcome. Journal of the American College of Cardiology, 2012, 60, B84. | 1.2 | 1 |
| 209 | TCT-641 Improved Safety and Reduction in Stent Thrombosis After Everolimus-Eluting Stents Versus Sirolimus-Eluting Stents in Patients With Coronary Artery Disease: SORT OUT IV 5-year. Journal of the American College of Cardiology, 2014, 64, B186-B187. | 1.2 | 1 |
| 210 | TCT-318 Ten-year All-cause Mortality after Simple versus Complex Stenting of Coronary Artery Bifurcation Lesions in the Randomized Nordic BifurcationÂStudy. Journal of the American College of Cardiology, 2016, 68, B131-B132. | 1.2 | 1 |
| 211 | The year in cardiology 2017: coronary interventions. European Heart Journal, 2018, 39, 914-924. | 1.0 | 1 |
| 212 | <p>Extent of coronary artery disease is associated with myocardial infarction and mortality in patients with diabetes mellitus [Response to Letter]</p> . Clinical Epidemiology, 2019, Volume 11, 721-722. | 1.5 | 1 |
| 213 | Instantaneous wave-free ratio cutoff values for nonculprit stenosis classification in patients with ST-segment elevation myocardial infarction (an iSTEMI substudy). Coronary Artery Disease, 2020, 31, 411-416. | 0.3 | 1 |
| 214 | Statin but not aspirin treatment is associated with reduced cardiovascular risk in patients with diabetes without obstructive coronary artery disease: a cohort study from the Western Denmark Heart Registry. European Heart Journal - Cardiovascular Pharmacotherapy, 2022, 8, 434-441. | 1.4 | 1 |
| 215 | Optical coherence tomography tissue coverage and characterization at six months after implantation of bioresorbable scaffolds versus conventional everolimus eluting stents in the ISAR-Absorb MI trial. International Journal of Cardiovascular Imaging, 2021, 37, 2815-2826. | 0.7 | 1 |
| 216 | Impact of drug eluting stents on target vessel revascularization. A report from the Western Denmark heart registry from 2000 to 2004. EuroIntervention, 2006, 1, 391-5. | 1.4 | 1 |

| # | Article | IF | CITATIONS |
|-----|--|-----|-----------|
| 217 | Long-Term Outcomes of Perioperative Versus Nonoperative Myocardial Infarction: A Danish Population-Based Cohort Study (2000–2016). Circulation: Cardiovascular Quality and Outcomes, 2022, 15, . | 0.9 | 1 |
| 218 | Intravascular ultrasound and angiographic assessment of reference segment size in type-2 diabetic patients. International Journal of Cardiology, 2007, 119, S27-S28. | 0.8 | 0 |
| 219 | Response to Letter Regarding Article, "Primary Angioplasty Versus Fibrinolysis in Acute Myocardial Infarction: Long-Term Follow-Up in the Danish Acute Myocardial Infarction 2 Trial― Circulation, 2010, 122, . | 1.6 | 0 |
| 220 | COMMENTARY: Deliver the Drug and Disappear: Is the Bioabsorbable Magnesium Stent Growing Up or Still Shrinking?. Journal of Endovascular Therapy, 2011, 18, 416-417. | 0.8 | 0 |
| 221 | TCT-301 Feasibility and Results of Novel Side Branch Evaluation by Reconstructed 3-Dimensional Optical Coherence Tomography. Matched Analysis of Baseline and 12-month follow-up in Native, Jailed and Opened Side Branches. Journal of the American College of Cardiology, 2012, 60, B85-B86. | 1.2 | 0 |
| 222 | TCT-347 Clinical Outcome Of Biolimus-Eluting Versus Sirolimus-Eluting Coronary Stent Implantation In Patients With And Without Diabetes Mellitus: A SORT OUT V Substudy. Journal of the American College of Cardiology, 2012, 60, B98-B99. | 1.2 | 0 |
| 223 | TCT-531 Positive Predictive Value Of Clinically Suspected ST-Segment Elevation Myocardial Infarction Using Angiographic Verification. Journal of the American College of Cardiology, 2012, 60, B154. | 1.2 | O |
| 224 | Clinical outcomes after treatment of multiple lesions with zotarolimus-eluting versus sirolimus-eluting coronary stents (a SORT OUT III substudy). BMC Cardiovascular Disorders, 2012, 12, 18. | 0.7 | 0 |
| 225 | TCT-598 Mechanisms of Incomplete Stent Apposition After Implantation of Drug-eluting Stents in patients with ST-segment Elevation Myocardial Infarction. Journal of the American College of Cardiology, 2013, 62, B181. | 1.2 | 0 |
| 226 | TCT-595 Early Healing After Treatment Of Coronary Lesions By Everolimus, Or Biolimus Eluting Bioresorbable Polymer Stents. One-month Results In The SORT-OUT VIII Optical Coherence Tomography Study. Journal of the American College of Cardiology, 2014, 64, B174. | 1.2 | 0 |
| 227 | Coronary stents and non-cardiac surgery: to bridge or not to bridge?. Thrombosis and Haemostasis, 2015, 114, 211-213. | 1.8 | 0 |
| 228 | Reply. Journal of the American College of Cardiology, 2015, 66, 2266-2267. | 1.2 | 0 |
| 229 | TCT-261 Impact of Diabetes on Clinical Outcomes after Revascularization with Sirolimus-eluting and Birolimus-Eluting Stents with biodegradable polymer. From the SORT OUT VII Trial. Journal of the American College of Cardiology, 2016, 68, B106. | 1.2 | 0 |
| 230 | TCT-321 Definite and probable stent thrombosis after revascularization with drug-eluting stents with a biodegradable polymer. From the randomized SORT OUT VII Trial. Journal of the American College of Cardiology, 2016, 68, B133. | 1.2 | 0 |
| 231 | TCT-347 Culprit Lesion Types and Vascular Healing in Patients with ST-segment Elevation Myocardial Infarction after Primary Percutaneous Intervention. Journal of the American College of Cardiology, 2018, 72, B141. | 1.2 | 0 |
| 232 | TCT-792 Everolimus-eluting versus biolimus-eluting stent implantation in unselected patients with and without diabetes: a SORT OUT VIII substudy. Journal of the American College of Cardiology, 2018, 72, B315. | 1.2 | 0 |
| 233 | TCT-290 Outcomes After Revascularization With a Polymer-Free Biolimus-Eluting BioFreedom Stent or a Biodegradable Polymer Ultrathin Strut Sirolimus-Eluting Orsiro Stent in Patients With and Without Acute Coronary Syndromes: From the SORT OUT IX Trial. Journal of the American College of Cardiology, 2019, 74, B289. | 1.2 | 0 |
| 234 | TCT-11 Everolimus-Eluting Bioresorbable Scaffolds Versus Drug-Eluting Stents in Patients With Acute Myocardial Infarction: 2-Year Results of the Randomized ISAR-Absorb MI Trial. Journal of the American College of Cardiology, 2019, 74, B11. | 1.2 | 0 |

| # | Article | IF | CITATIONS |
|-----|---|-----|-----------|
| 235 | Research update for articles published in EJCI in 2017. European Journal of Clinical Investigation, 2019, 49, e13163. | 1.7 | 0 |
| 236 | Very late Absorb scaffold thrombosis. Coronary Artery Disease, 2019, 30, 232-233. | 0.3 | 0 |
| 237 | Response to †Correspondence on †Impact of rheumatoid arthritis on major cardiovascular events in patients with and without coronary artery disease†by Jong et al. Annals of the Rheumatic Diseases, 2020, , annrheumdis-2020-219231. | 0.5 | 0 |
| 238 | Reply. JACC: Cardiovascular Interventions, 2020, 13, 658-659. | 1.1 | 0 |
| 239 | Prognostic Value of Coronary Artery Calcium in Symptomatic Young Individuals Age 18 to 45 Years. Journal of the American College of Cardiology, 2021, 77, 2980-2982. | 1.2 | 0 |
| 240 | Comment on: Cardiovascular risk and mortality in rheumatoid arthritis compared with diabetes mellitus and the general population: reply. Rheumatology, 2021, 60, e419-e420. | 0.9 | 0 |
| 241 | Culprit lesion morphology in patients with ST-segment elevation myocardial infarction assessed by optical coherence tomography. Coronary Artery Disease, 2020, 31, 671-677. | 0.3 | 0 |
| 242 | Cardiovascular risks associated with smoking in patients without obstructive coronary artery disease. European Journal of Preventive Cardiology, 2022, 29, e14-e17. | 0.8 | 0 |
| 243 | CHA 2 DS 2 â€VASc impact on risk following percutaneous coronary intervention in atrial fibrillation. European Journal of Clinical Investigation, 2021, , e13717. | 1.7 | 0 |
| 244 | Early vascular healing after implantation of the polymer-free biolimus-eluting stent or the ultrathin strut biodegradable polymer sirolimus-eluting stent in patients with ST-segment elevation myocardial infarction. Coronary Artery Disease, 2022, Publish Ahead of Print, . | 0.3 | 0 |
| 245 | Impact of diabetes on 1â€year clinical outcome in patients undergoing revascularization with the BioFreedom stents or the Orsiro stents from the SORT OUT IX trial. Catheterization and Cardiovascular Interventions, 2022, , . | 0.7 | 0 |