Eva TesaÅð₩Ã;

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/4489341/publications.pdf

Version: 2024-02-01

		159585	2	33421
120	2,875	30		45
papers	citations	h-index		g-index
			_	
100	100	100		0016
120	120	120		2316
all docs	docs citations	times ranked		citing authors

#	Article	IF	CITATIONS
1	Phosphorothioate oligonucleotides separation in ion-pairing reversed-phase liquid chromatography: Effect of ion-pairing system. Journal of Chromatography A, 2022, 1676, 463201.	3.7	11
2	Enantioseparation performance of superficially porous particle vancomycin-based chiral stationary phases in supercritical fluid chromatography and high performance liquid chromatography; applicability for psychoactive substances. Journal of Chromatography A, 2021, 1637, 461846.	3.7	20
3	Characterization and comparison of mixed-mode and reversed-phase columns; interaction abilities and applicability for peptide separation. Journal of Chromatography A, 2021, 1648, 462182.	3.7	13
4	The effect of tandem coupling of NicoShell and TeicoShell columns in sub/supercritical fluid chromatography on enantioresolution. Journal of Separation Science, 2021, 44, 4048-4057.	2.5	4
5	Enantioselective potential of teicoplanin- and vancomycin-based superficially porous particles-packed columns for supercritical fluid chromatography. Journal of Chromatography A, 2020, 1612, 460687.	3.7	18
6	The effect of particle and ligand types on retention and peak shape in liquid chromatography. Microchemical Journal, 2020, 159, 105466.	4. 5	6
7	Chiral separation of beta-blockers by high-performance liquid chromatography and determination of bisoprolol enantiomers in surface waters. Arhiv Za Higijenu Rada I Toksikologiju, 2020, 71, 56-62.	0.7	5
8	Method for evaluation of ionic interactions in liquid chromatography. Journal of Chromatography A, 2020, 1625, 461301.	3.7	18
9	Enantiorecognition ability of different chiral selectors for separation of liquid crystals in supercritical fluid chromatography; critical evaluation. Journal of Chromatography A, 2020, 1622, 461138.	3.7	11
10	Evaluation of separation properties of stationary phases in supercritical fluid chromatography; deazapurine nucleosides case study. Microchemical Journal, 2019, 150, 104137.	4.5	7
11	The degree of substitution affects the enantioselectivity of sulfobutyletherâ€Î²â€cyclodextrin chiral stationary phases. Electrophoresis, 2019, 40, 1972-1977.	2.4	14
12	Fast enantioseparation of indole phytoalexins in additive free supercritical fluid chromatography. Journal of Chromatography A, 2019, 1596, 209-216.	3.7	11
13	Systematic evaluation of selected supercritical fluid chromatography diol―and diethylamineâ€based columns for application in hydrophilic interaction liquid chromatography. Separation Science Plus, 2019, 2, 81-88.	0.6	9
14	Cellulose trisâ€(3,5â€dimethylphenylcarbamate)â€based chiral stationary phase for the enantioseparation of drugs in supercritical fluid chromatography: comparison with HPLC. Journal of Separation Science, 2018, 41, 1471-1478.	2.5	37
15	Chromatographic behavior of new deazapurine ribonucleosides in hydrophilic interaction liquid chromatography. Electrophoresis, 2018, 39, 2144-2151.	2.4	5
16	Sulfated Metabolites of Flavonolignans and 2,3-Dehydroflavonolignans: Preparation and Properties. International Journal of Molecular Sciences, 2018, 19, 2349.	4.1	23
17	Performance comparison of three trypsin columns used in liquid chromatography. Journal of Chromatography A, 2017, 1490, 126-132.	3.7	20
18	Enantioselective potential of polysaccharideâ€based chiral stationary phases in supercritical fluid chromatography. Chirality, 2017, 29, 239-246.	2.6	22

#	Article	IF	CITATIONS
19	Immobilized strychnine as a new chiral stationary phase for HPLC. Electrophoresis, 2017, 38, 1956-1963.	2.4	6
20	Cyclic Oligosaccharide-Based Chiral Stationary Phases Applicable to Drug Purity Control; A Review. Current Medicinal Chemistry, 2017, 24, 829-848.	2.4	7
21	Enantiomeric Ratio of Amino Acids as a Tool for Determination of Aging and Disease Diagnostics by Chromatographic Measurement. Separations, 2016, 3, 30.	2.4	18
22	The empirical comparison of cyclofructans and cyclodextrins as chiral selectors in capillary electrophoretic separation of atropisomers of <i>R</i> , <i>S</i> â€1,1'â€binaphthaleneâ€2,2'â€diyl hydrog phosphate. Journal of Separation Science, 2016, 39, 973-979.	g e n5	10
23	Enantioselective separation of biologically active basic compounds in ultra-performance supercritical fluid chromatography. Analytica Chimica Acta, 2016, 932, 98-105.	5.4	29
24	Chromatographic methods enabling the characterization of stationary phases and retention prediction in highâ€performance liquid chromatography and supercritical fluid chromatography. Journal of Separation Science, 2016, 39, 115-131.	2.5	21
25	Chromatographic Characterization of a New Cationic Î ² -CD Based Stationary Phase Prepared by Dynamic Coating. Chromatographia, 2016, 79, 529-536.	1.3	8
26	Sulfobutylether- \hat{l}^2 -cyclodextrin as a chiral selector for separation of amino acids and dipeptides in chromatography. Journal of Chromatography A, 2016, 1467, 356-362.	3.7	23
27	Development of separation methods for the chiral resolution of hexahelicenes. Journal of Chromatography A, 2016, 1476, 130-134.	3.7	17
28	Direct CE and HPLC methods for enantioseparation of tryptophan and its unnatural derivatives. Separation and Purification Technology, 2016, 158, 24-30.	7.9	12
29	Evaluation of differences between Chiralpak IA and Chiralpak ADâ€RH amyloseâ€based chiral stationary phases in reversedâ€phase highâ€performance liquid chromatography. Journal of Separation Science, 2015, 38, 711-719.	2.5	15
30	Pharmacokinetics of pure silybin diastereoisomers and identification of their metabolites in rat plasma. Journal of Functional Foods, 2015, 14, 570-580.	3.4	25
31	Immobilized Polysaccharide-Based Stationary Phases for Enantioseparation in Normal Versus Reversed Phase HPLC. Chromatographia, 2015, 78, 909-915.	1.3	13
32	A supercritical fluid chromatography method for the systematic toxicology analysis of cannabinoids and their metabolites. Analytical Methods, 2015, 7, 6056-6059.	2.7	16
33	Insight into Trypsin Miscleavage: Comparison of Kinetic Constants of Problematic Peptide Sequences. Analytical Chemistry, 2015, 87, 7636-7643.	6.5	77
34	Effect of Buffer Constituents on Retention and Separation in Achiral and Chiral HPLC Systems with \hat{l}^2 -Cyclodextrin-Based Stationary Phase. Chromatographia, 2015, 78, 917-921.	1.3	4
35	Supercritical fluid chromatography as a tool for enantioselective separation; A review. Analytica Chimica Acta, 2014, 821, 1-33.	5.4	144
36	Isopropyl derivative of cyclofructan 6 as chiral selector in liquid chromatography and capillary electrophoresis. Journal of Chromatography A, 2014, 1338, 197-200.	3.7	23

#	Article	IF	CITATIONS
37	Phototransformation of benzimidazole and thiabendazole inside cucurbit[8]uril. Photochemical and Photobiological Sciences, 2014, 13, 310-315.	2.9	17
38	Characterization of novel metallacarborane-based sorbents by linear solvation energy relationships. Journal of Chromatography A, 2014, 1371, 220-226.	3.7	6
39	Enantioselective potential of chiral stationary phases based on immobilized polysaccharides in reversed phase mode. Journal of Chromatography A, 2014, 1363, 155-161.	3.7	55
40	HPLC Method for Chiral Separation and Quantification of Antidepressant Citalopram and Its Precursor Citadiol. Chromatographia, 2013, 76, 483-489.	1.3	23
41	Complexation of Buffer Constituents with Neutral Complexation Agents: Part I. Impact on Common Buffer Properties. Analytical Chemistry, 2013, 85, 8518-8525.	6.5	31
42	Complexation of Buffer Constituents with Neutral Complexation Agents: Part II. Practical Impact in Capillary Zone Electrophoresis. Analytical Chemistry, 2013, 85, 8526-8534.	6.5	30
43	Properties of two amideâ€based hydrophilic interaction liquid chromatography columns. Journal of Separation Science, 2013, 36, 2421-2429.	2.5	27
44	Rapid Supercritical Fluid Chromatography Method for Separation of Chlorthalidone Enantiomers. Analytical Letters, 2013, 46, 2860-2869.	1.8	12
45	Hydrophilic interaction liquid chromatography with tandem mass spectrometric detection applied for analysis of pteridines in two Graphosoma species (Insecta: Heteroptera). Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences, 2013, 930, 82-89.	2.3	16
46	Enantioselective separation of unusual amino acids by high performance liquid chromatography. Separation and Purification Technology, 2013, 119, 123-128.	7.9	27
47	An insight into the use of dimethylphenyl carbamate cyclofructan 7 chiral stationary phase in supercritical fluid chromatography: The basic comparison with HPLC. Journal of Separation Science, 2013, 36, 1711-1719.	2.5	30
48	Determination of nitrite and nitrate in cerebrospinal fluid by microchip electrophoresis with microsolid phase extraction pre-treatment. Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences, 2013, 930, 41-47.	2.3	28
49	Effect of silica gel modification with cyclofructans on properties of hydrophilic interaction liquid chromatography stationary phases. Journal of Chromatography A, 2012, 1257, 58-65.	3.7	35
50	Chiral HPLC Separation on Derivatized Cyclofructan Versus Cyclodextrin Stationary Phases. Analytical Letters, 2012, 45, 2344-2358.	1.8	18
51	Study on the use of boromycin as a chiral selector in capillary electrophoresis. Journal of Chromatography A, 2012, 1237, 128-132.	3.7	30
52	Recentchiral selectors for separation in HPLC and CE. Open Chemistry, 2012, 10, 450-471.	1.9	36
53	Use of Capillary Zone Electrophoresis and Micellar Electrokinetic Chromatography for Separations of Anthraquinone Derivatives. Analytical Letters, 2011, 44, 1783-1795.	1.8	1
54	Methods for determination of all binding parameters in systems with simultaneous borate and cyclodextrin complexation. Journal of Chromatography A, 2011, 1218, 7211-7218.	3.7	7

#	Article	IF	Citations
55	Cyclofructan 6 based stationary phases for hydrophilic interaction liquid chromatography. Journal of Chromatography A, 2011, 1218, 270-279.	3.7	73
56	Characterization of cyclofructanâ€based chiral stationary phases by linear free energy relationship. Journal of Separation Science, 2011, 34, 2639-2644.	2.5	30
57	Accuracy and sensitivity of the determination of rate constants of interconversion in achiral and chiral environments by dynamic enantioselective electrophoresis. Electrophoresis, 2011, 32, 595-603.	2.4	6
58	Development of a solid-phase extraction with capillary liquid chromatography tandem mass spectrometry for analysis of estrogens in environmental water samples. Journal of Chromatography A, 2011, 1218, 2127-2132.	3.7	24
59	Characterization of new R-naphthylethyl cyclofructan 6 chiral stationary phase and its comparison with R-naphthylethyl \hat{I}^2 -cyclodextrin-based column. Journal of Chromatography A, 2011, 1218, 1393-1398.	3.7	37
60	Monolithic columns based on a poly(styrene-divinylbenzene-methacrylic acid) copolymer for capillary liquid chromatography of small organic molecules. Journal of Chromatography A, 2011, 1218, 1544-1547.	3.7	37
61	Separation and Quantification of 1,4-benzodiazepines: HPLC versus CZE. Croatica Chemica Acta, 2011, 84, 367-373.	0.4	5
62	Enhanced selectivity in CZE multiâ€chiral selector enantioseparation systems: Proposed separation mechanism. Electrophoresis, 2010, 31, 1435-1441.	2.4	54
63	Comparison of HPLC enantioseparation of substituted binaphthyls on CDâ€, polysaccharideâ€, and synthetic polymerâ€based chiral stationary phases. Journal of Separation Science, 2010, 33, 1244-1254.	2.5	14
64	Pluronic Fâ€127 as the buffer additive in capillary entangled polymer electrophoresis: Some fundamental aspects. Journal of Separation Science, 2010, 33, 2458-2464.	2.5	10
65	Study of interaction mechanisms on zirconiaâ€based polystyrene HPLC column. Journal of Separation Science, 2010, 33, 3043-3051.	2.5	15
66	Sterility testing by CE: A comparison of online preconcentration approaches in capillaries with greater internal diameters. Electrophoresis, 2009, 30, 3870-3876.	2.4	20
67	Separation of inorganic and small organic anions by CE using phosphoniumâ€based mono†and dicationic reagents. Electrophoresis, 2009, 30, 3955-3963.	2.4	27
68	Comparison of enantioselective HPLC separation of structurally diverse compounds on chiral stationary phases with different teicoplanin coverage and distinct linkage chemistry. Journal of Separation Science, 2009, 32, 1704-1711.	2.5	22
69	Occurrence and behavior of system peaks in RP HPLC with solely aqueous mobile phases. Journal of Separation Science, 2009, 32, 2864-2870.	2.5	2
70	A spectroelectrochemical approach to the electrodeposition of bismuth film electrodes and their use in stripping analysis. Analytica Chimica Acta, 2008, 608, 140-146.	5.4	16
71	Cellulose tris(3,5â€dimethylphenylcarbamate)â€based chiral stationary phases as effective tools for enantioselective HPLC separation of structurally different disubstituted binaphthyls. Chirality, 2008, 20, 900-909.	2.6	9
72	System peaks in micellar electrophoresis: I. Utilization of system peaks for determination of critical micelle concentration. Electrophoresis, 2008, 29, 1189-1195.	2.4	13

#	Article	IF	CITATIONS
73	HPLC method for enantioselective analysis of cloprostenol. Journal of Pharmaceutical and Biomedical Analysis, 2008, 46, 892-897.	2.8	7
74	LC with a Teicoplanin Aglycone Chiral Sorbent for the Separation of the Enantiomers of Non-Steroidal Anti-Inflammatory Drugs: An Evaluation of Chiral Capillary Columns. Chromatographia, 2008, 67, 33-40.	1.3	6
75	Model of CE enantioseparation systems with a mixture of chiral selectors. Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences, 2008, 875, 35-41.	2.3	20
76	On-line preconcentration of weak electrolytes by electrokinetic accumulation in CE: Experiment and simulation. Electrophoresis, 2007, 28, 1540-1547.	2.4	34
77	Separation and quantification of 9â€(alkylthio)acridines by capillary micellar electrokinetic chromatography and capillary liquid chromatography. Journal of Separation Science, 2007, 30, 2123-2129.	2.5	2
78	Effects of capillary coating and \hat{l}^2 -cyclodextrin additive to the background electrolyte on separation of sulphonated azodyes by capillary zone electrophoresis. Journal of Chromatography A, 2007, 1149, 358-367.	3.7	16
79	Vancomycin as chiral selector for enantioselective separation of selected profen nonsteroidal anti-inflammatory drugs incapillary liquid chromatography. Chirality, 2006, 18, 531-538.	2.6	24
80	Eigenmobilities in background electrolytes for CZE. V. Intensity (amplitudes) of system peaks. Electrophoresis, 2006, 27, 4610-4617.	2.4	24
81	Evaluation and comparison of a methylated teicoplanin aglycone to teicoplanin aglycone and natural teicoplanin chiral stationary phases. Journal of Separation Science, 2006, 29, 429-445.	2.5	43
82	Linear free energy relationship as a tool for characterization of three teicoplanin-based chiral stationary phases under various mobile phase compositions. Journal of Separation Science, 2006, 29, 1476-1485.	2.5	17
83	Chiral separation of tamsulosin by capillary electrophoresis. Journal of Pharmaceutical and Biomedical Analysis, 2005, 39, 691-696.	2.8	30
84	Comparison of vancomycin-based stationary phases with different chiral selector coverage for enantioselective separation of selected drugs in high-performance liquid chromatography. Journal of Chromatography A, 2005, 1088, 94-103.	3.7	55
85	Comparison of zirconia- and silica-based reversed stationary phases for separation of enkephalins. Journal of Chromatography A, 2005, 1087, 104-111.	3.7	10
86	Chiral separation of beta-adrenergic antagonists, profen non-steroidal anti-inflammatory drugs and chlorophenoxypropionic acid herbicides using teicoplanin as the chiral selector in capillary liquid chromatography. Journal of Chromatography A, 2005, 1088, 82-93.	3.7	37
87	Comparative study of three teicoplanin-based chiral stationary phases using the linear free energy relationship model. Journal of Chromatography A, 2005, 1088, 57-66.	3.7	21
88	System peaks and their positive and negative aspects in chromatographic techniques. Journal of Separation Science, 2005, 28, 1263-1270.	2.5	14
89	Dynamics of interconversion of enantiomers in chiral separation systems: A novel approach for determination of all rate constants involved in the interconversion. Electrophoresis, 2004, 25, 733-742.	2.4	32
90	Effects of partial/asymmetrical filling of micelles and chiral selectors on capillary electrophoresis enantiomeric separation: Generation of a gradient. Electrophoresis, 2004, 25, 2693-2700.	2.4	17

#	Article	IF	CITATIONS
91	Optimization of binary porogen solvent composition for preparation of butyl methacrylate monoliths in capillary liquid chromatography. Journal of Chromatography A, 2004, 1049, 43-49.	3.7	41
92	Optimization of binary porogen solvent composition for preparation of butyl methacrylate monoliths in capillary liquid chromatography. Journal of Chromatography A, 2004, 1049, 43-49.	3.7	41
93	Enantioseparation of dihydrofurocoumarin derivatives by various separation modes of capillary electrophoresis. Electrophoresis, 2003, 24, 2650-2656.	2.4	15
94	Comparison of enantioseparation of selected benzodiazepine and phenothiazine derivatives on chiral stationary phases based on \hat{l}^2 -cyclodextrin and macrocyclic antibiotics. Journal of Separation Science, 2003, 26, 661-668.	2.5	9
95	Capillary liquid chromatography as a tool for separation of hydrophobic basic drugs. Relation between tests for column characterization and real analysis. Journal of Separation Science, 2003, 26, 686-692.	2.5	5
96	Determination of new pyridoquinoline derivatives and their quantification in urine by capillary liquid chromatography. Journal of Separation Science, 2003, 26, 1582-1588.	2.5	4
97	Methacrylate monolithic columns for capillary liquid chromatography polymerized using ammonium peroxodisulfate as initiator. Journal of Separation Science, 2003, 26, 1623-1628.	2.5	45
98	Quantification and purity determination of newly synthesized thioacridines by capillary liquid chromatography. Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences, 2002, 770, 183-189.	2.3	12
99	Study of the stability of promethazine enantiomers by liquid chromatography using a vancomycin-bonded chiral stationary phase. Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences, 2002, 770, 63-69.	2.3	23
100	Methacrylate monolithic columns of 320 $\hat{l}\frac{1}{4}$ m I.D. for capillary liquid chromatography. Journal of Chromatography A, 2002, 946, 99-106.	3.7	98
101	Study on the aggregation of teicoplanin. Talanta, 2001, 54, 643-653.	5.5	18
102	Hybrid Polymeric Micelles with Hydrophobic Cores and Mixed Polyelectrolyte/Nonelectrolyte Shells in Aqueous Media. 1. Preparation and Basic Characterization. Langmuir, 2001, 17, 4240-4244.	3.5	88
103	Enantioseparation of selected N-tertbutyloxycarbonyl amino acids in high-performance liquid chromatography and capillary electrophoresis with a teicoplanin chiral selector. Journal of Chromatography A, 2000, 879, 147-156.	3.7	30
104	Capillary Electrokinetic Chromatography with Charged Linear Polymers as a Nonmicellar PseudoStationary Phase:Â Determination of Capacity Factors and Characterization by Solvation Parameters. Analytical Chemistry, 2000, 72, 74-80.	6.5	31
105	Comparison of enantioselective separation of N-tertbutyloxycarbonyl amino acids and their non-blocked analogues on teicoplanin-based chiral stationary phase. Journal of Chromatography A, 1999, 838, 121-129.	3.7	46
106	Electromigration behavior of metal ions in the presence of complexing polymer. Journal of Chromatography A, 1999, 838, 101-109.	3.7	4
107	Enantioseparation of semisynthetic ergot alkaloids on vancomycin and teicoplanin stationary phases. Journal of Chromatography A, 1999, 844, 137-147.	3.7	36
108	Enantioselective Separations. Journal of Chromatography Library, 1998, 60, 197-256.	0.1	3

#	Article	IF	CITATIONS
109	Selected Derivatization Reactions. Journal of Chromatography Library, 1998, 60, 141-196.	0.1	2
110	Structural study of flobufen II. An unexpected role of packing effects on the dihedral angle of phenyl rings in crystal structures of 2,4-difluorobiphenyls. Journal of Fluorine Chemistry, 1997, 83, 111-116.	1.7	5
111	Interactions of basic compounds in reversed-phase high-performance liquid chromatography influence of sorbent character, mobile phase composition, and pH on retention of basic compounds. Journal of Chromatography A, 1997, 758, 37-51.	3.7	92
112	Physical factors negatively affecting evaluation of long-term biodegradation experiments of polychlorinated biphenyls. Chemosphere, 1996, 33, 2411-2421.	8.2	10
113	Enantiomer separation of dihydropyridine calcium antagonists with cyclodextrins as chiral selectors: structural correlation. Biomedical Applications, 1996, 681, 133-141.	1.7	35
114	New Organic Monosized Microspheres for Use in Enantiomer Separations by High-Performance Liquid Chromatography. Journal of Liquid Chromatography and Related Technologies, 1995, 18, 3187-3203.	1.0	10
115	Chiral separation by analytical electromigration methods. Journal of Chromatography A, 1992, 609, 1-17.	3.7	102
116	Separation of type IX collagen from other cartilage collagens by hydrophobic interaction chromatography. Biomedical Applications, 1988, 434, 423-427.	1.7	0
117	Reversed-phase thin-layer chromatography of phenolic compounds. Journal of High Resolution Chromatography, 1987, 10, 404-408.	1.4	2
118	High-performance liquid chromatography of biphenols and bis (hydroxyphenyl) propanes (dianes) with voltammetric and UV photometric detection. Chromatographia, 1987, 23, 102-108.	1.3	4
119	Gas and high-performance liquid chromatography of phenols. Chromatographia, 1983, 17, 269-284.	1.3	68
120	Separation and behaviour of s-triazine derivatives on a NH2-chemically bonded stationary phase by high-performance liquid chromatography. Journal of Chromatography A, 1980, 191, 115-120.	3.7	22