

Akira Watanabe

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4493210/publications.pdf>

Version: 2024-02-01

60
papers

677
citations

567281

15
h-index

677142

22
g-index

60
all docs

60
docs citations

60
times ranked

1153
citing authors

#	ARTICLE	IF	CITATIONS
1	Systematic review of resectable intraductal tubulopapillary neoplasm with special reference to recurrence patterns. <i>Surgery Today</i> , 2022, 52, 532-541.	1.5	2
2	Direct Bilirubin in Drainage Fluid 3 days after Hepatectomy Is Useful for Detecting Severe Bile Leakage. <i>European Surgical Research</i> , 2022, 63, 33-39.	1.3	1
3	Significance of zinc deficiency in patients with hepatocellular carcinoma undergoing hepatic resection. <i>Hepatology Research</i> , 2022, 52, 210-220.	3.4	6
4	Study on the portal ramification pattern of the right anterior sector of the liver and a unique medial branch (PV8c) of the right anterior portal vein. <i>Annals of Gastroenterological Surgery</i> , 2022, 6, 679-687.	2.4	2
5	Mac-2 Binding Protein Glycosylation Isomer as a Prognostic Marker for Hepatocellular Carcinoma With Sustained Virological Response. <i>Anticancer Research</i> , 2022, 42, 245-251.	1.1	1
6	The ratio of creatinine and cystatin C estimated glomerular filtration rates as a surrogate marker in patients with hepatocellular carcinoma undergoing hepatic resection. <i>Journal of Hepato-Biliary-Pancreatic Sciences</i> , 2022, 29, 964-973.	2.6	4
7	Unclassified hepatocellular adenoma with beta-catenin mutation: a case report. <i>Surgical Case Reports</i> , 2021, 7, 46.	0.6	2
8	Conophylline Inhibits Hepatocellular Carcinoma by Inhibiting Activated Cancer-associated Fibroblasts Through Suppression of G Protein-coupled Receptor 68. <i>Molecular Cancer Therapeutics</i> , 2021, 20, 1019-1028.	4.1	13
9	Hepatic stellate cell as a Mac-2-binding protein-producing cell in patients with liver fibrosis. <i>Hepatology Research</i> , 2021, 51, 1058-1063.	3.4	7
10	FDG-PET for preoperative evaluation of tumor invasion in ampullary cancer: A retrospective analysis. <i>Journal of Surgical Oncology</i> , 2021, 124, 317-323.	1.7	3
11	Impact of preoperative nutritional support and rehabilitation therapy in patients undergoing pancreaticoduodenectomy. <i>International Journal of Clinical Oncology</i> , 2021, 26, 1698-1706.	2.2	21
12	Gastrectomy history as a predictor of post-hepatectomy complications. <i>Surgery Today</i> , 2021, , 1.	1.5	0
13	Skeletal Muscle Loss and Octogenarian Status Are Associated with S-1 Adjuvant Therapy Discontinuation and Poor Prognosis after Pancreatectomy. <i>Cancers</i> , 2021, 13, 4105.	3.7	3
14	Pancreatic Xanthogranuloma with High Accumulation in Fluorodeoxyglucose-Positron Emission Tomography. <i>Nihon Gekakei Rengo Gakkaishi (Journal of Japanese College of Surgeons)</i> , 2021, 46, 90-96.	0.0	0
15	FDG-PET reflects tumor viability on SUV in colorectal cancer liver metastasis. <i>International Journal of Clinical Oncology</i> , 2020, 25, 322-329.	2.2	8
16	High Co-expression of Large Tenascin C Splice Variants in Stromal Tissue and Annexin A2 in Cancer Cell Membranes is Associated with Poor Prognosis in Pancreatic Cancer. <i>Annals of Surgical Oncology</i> , 2020, 27, 924-930.	1.5	12
17	Predictors of Postoperative Ascites After Hepatic Resection in Patients With Hepatocellular Carcinoma. <i>Anticancer Research</i> , 2020, 40, 4343-4349.	1.1	4
18	Diffusion-Weighted MRI Predicts Lymph Node Metastasis and Tumor Aggressiveness in Resectable Pancreatic Neuroendocrine Tumors. <i>World Journal of Surgery</i> , 2020, 44, 4136-4141.	1.6	8

#	ARTICLE	IF	CITATIONS
19	Mac-2-binding protein glycan isomer enhances the aggressiveness of hepatocellular carcinoma by activating mTOR signaling. <i>British Journal of Cancer</i> , 2020, 123, 1145-1153.	6.4	20
20	Nintedanib inhibits intrahepatic cholangiocarcinoma aggressiveness via suppression of cytokines extracted from activated cancer-associated fibroblasts. <i>British Journal of Cancer</i> , 2020, 122, 986-994.	6.4	44
21	Albumin-Indocyanine Green Evaluation (ALICE) grade predicts bile leakage after hepatic resection. <i>Surgery Today</i> , 2020, 50, 849-854.	1.5	9
22	Right hepatic venous system variation in living donors: a three-dimensional CT analysis. <i>British Journal of Surgery</i> , 2020, 107, 1192-1198.	0.3	7
23	Unclassified hepatocellular adenoma with histological brown pigment deposition and serum PIVKA-II level elevation: a case report. <i>Surgical Case Reports</i> , 2020, 6, 94.	0.6	5
24	Rare metastatic pancreatic tumors from lung cancer with cystic changes resembling intraductal papillary mucinous neoplasm: a case report. <i>Surgical Case Reports</i> , 2020, 6, 117.	0.6	0
25	Liver metastasis from papillary thyroid carcinoma treated by laparoscopic hepatectomy 43 years after resection of the primary tumor: a case report. <i>Surgical Case Reports</i> , 2020, 6, 142.	0.6	4
26	Multimodal approach to portal hypertension and gastric varices before hepatic resection for hepatocellular carcinoma: a case report. <i>Surgical Case Reports</i> , 2020, 6, 190.	0.6	1
27	Author response to: Comment on: Right hepatic venous system variation in living donors: a three-dimensional CT analysis. <i>British Journal of Surgery</i> , 2020, 107, e608.	0.3	0
28	Author response to: Comment on: Right hepatic venous system variation in living donors: a three-dimensional CT analysis. <i>British Journal of Surgery</i> , 2020, 107, e653.	0.3	0
29	Prognostic significance of neutrophil-lymphocyte ratio in resectable pancreatic neuroendocrine tumors with special reference to tumor-associated macrophages. <i>Pancreatology</i> , 2019, 19, 897-902.	1.1	26
30	Successful resection of intrahepatic cholangiocarcinoma with idiopathic thrombocytopenic purpura using thrombopoietin receptor agonist: a case report. <i>Surgical Case Reports</i> , 2019, 5, 56.	0.6	2
31	Hepatic pleomorphic leiomyosarcoma after surgery for gastric gastrointestinal stromal tumor: a case report. <i>Surgical Case Reports</i> , 2019, 5, 62.	0.6	2
32	Preoperative Mac-2 binding protein glycosylation isomer level predicts postoperative ascites in patients with hepatic resection for hepatocellular carcinoma. <i>Hepatology Research</i> , 2019, 49, 1398-1405.	3.4	12
33	Conophylline suppresses pancreatic cancer desmoplasia and cancer-promoting cytokines produced by cancer-associated fibroblasts. <i>Cancer Science</i> , 2019, 110, 334-344.	3.9	28
34	Impact of Metabolic Parameters of ¹⁸ F-Fluorodeoxyglucose Positron-emission Tomography After Hepatic Resection in Patients With Intrahepatic Cholangiocarcinoma. <i>Anticancer Research</i> , 2019, 39, 971-977.	1.1	5
35	A Surgically Resected Pancreatic Schwannoma with Obstructive Jaundice with Special Reference to Differential Diagnosis from Other Cystic Lesions in the Pancreas. <i>Case Reports in Gastroenterology</i> , 2018, 12, 85-91.	0.6	7
36	D-dimer predicts postoperative recurrence and prognosis in patients with liver metastasis of colorectal cancer. <i>International Journal of Clinical Oncology</i> , 2018, 23, 689-697.	2.2	26

#	ARTICLE	IF	CITATIONS
37	Evaluation of the International Consensus Guidelines for the Surgical Resection of Intraductal Papillary Mucinous Neoplasms. <i>Digestive Diseases and Sciences</i> , 2018, 63, 860-867.	2.3	10
38	Fairly rare clear cell adenocarcinoma mimicking liver cancer: a case report. <i>Surgical Case Reports</i> , 2018, 4, 97.	0.6	5
39	Successful treatment of isolated bile leakage after hepatectomy combination therapy with percutaneous transhepatic portal embolization and bile duct ablation with ethanol: a case report. <i>Surgical Case Reports</i> , 2018, 4, 61.	0.6	4
40	A new strategy based on fluorodeoxyglucose-positron emission tomography for managing liver metastasis from colorectal cancer. <i>Journal of Surgical Oncology</i> , 2018, 118, 1088-1095.	1.7	8
41	Skeletal Muscle Volume and Intramuscular Adipose Tissue Are Prognostic Predictors of Postoperative Complications After Hepatic Resection. <i>Anticancer Research</i> , 2018, 38, 4933-4939.	1.1	14
42	High expression of forkhead box protein C2 is associated with aggressive phenotypes and poor prognosis in clinical hepatocellular carcinoma. <i>BMC Cancer</i> , 2018, 18, 597.	2.6	21
43	High STMN1 Expression Is Associated with Tumor Differentiation and Metastasis in Clinical Patients with Pancreatic Cancer. <i>Anticancer Research</i> , 2018, 38, 939-944.	1.1	15
44	Perioperative management of hepatectomy in patients with interstitial pneumonia: a report of three cases and a literature review. <i>Surgery Today</i> , 2017, 47, 1173-1179.	1.5	3
45	Stathmin 1 promotes the proliferation and malignant transformation of pancreatic intraductal papillary mucinous neoplasms. <i>Oncology Letters</i> , 2017, 13, 1783-1788.	1.8	5
46	Enhanced karyopherin- β 2 expression is associated with carcinogenesis in patients with intraductal papillary mucinous neoplasms. <i>Pancreatology</i> , 2017, 17, 611-616.	1.1	7
47	Poor prognosis in cholangiocarcinoma patients with low FBXW7 expression is improved by chemotherapy. <i>Oncology Letters</i> , 2017, 13, 3653-3661.	1.8	9
48	High STMN1 Expression is Associated with Cancer Progression and Chemo-Resistance in Lung Squamous Cell Carcinoma. <i>Annals of Surgical Oncology</i> , 2017, 24, 4017-4024.	1.5	31
49	Complete resection of a huge pancreatic undifferentiated carcinoma with osteoclast-like giant cells. <i>International Cancer Conference Journal</i> , 2017, 6, 193-196.	0.5	5
50	Small cholangiolocellular carcinoma that was difficult to distinguish from cholangiocellular carcinoma: a case report. <i>Surgical Case Reports</i> , 2017, 3, 103.	0.6	8
51	Reduced FBXW7 expression in pancreatic cancer correlates with poor prognosis and chemotherapeutic resistance via accumulation of MCL1. <i>Oncotarget</i> , 2017, 8, 112636-112646.	1.8	20
52	Overexpression of natural killer group 2 member D ligands predicts favorable prognosis in cholangiocarcinoma. <i>Cancer Science</i> , 2016, 107, 116-122.	3.9	32
53	High expression of karyopherin- β 2 and stathmin 1 is associated with proliferation potency and transformation in the bile duct and gall bladder epithelia in the cases of pancreaticobiliary maljunction. <i>Journal of Surgical Oncology</i> , 2016, 114, 462-468.	1.7	6
54	Presence of Cytokeratin 19-Expressing Cholangiocarcinoma-Like Tumour in a Liver Metastatic Lesion of Rectal Neuroendocrine Tumour. <i>Case Reports in Gastroenterology</i> , 2016, 10, 431-439.	0.6	3

#	ARTICLE	IF	CITATIONS
55	Xanthogranulomatous cholecystitis: Difficulty in differentiating from gallbladder cancer. <i>World Journal of Gastroenterology</i> , 2015, 21, 10166-10173.	3.3	43
56	Nesfatin-1 Suppresses Gastric Contractions and Inhibits Interdigestive Migrating Contractions in Conscious Dogs. <i>Digestive Diseases and Sciences</i> , 2015, 60, 1595-1602.	2.3	22
57	Relationship between 18-F-fluoro-deoxy-d-glucose uptake and expression of glucose transporter 1 and pyruvate kinase M2 in intrahepatic cholangiocarcinoma. <i>Digestive and Liver Disease</i> , 2015, 47, 590-596.	0.9	18
58	Stathmin1 regulates p27 expression, proliferation and drug resistance, resulting in poor clinical prognosis in cholangiocarcinoma. <i>Cancer Science</i> , 2014, 105, 690-696.	3.9	46
59	A case of mesenteric cystic lymphangioma in an adult which caused duodenal stenosis after resection. <i>International Journal of Surgery Case Reports</i> , 2013, 4, 212-215.	0.6	17
60	Forkhead box protein C2 contributes to invasion and metastasis of extrahepatic cholangiocarcinoma, resulting in a poor prognosis. <i>Cancer Science</i> , 2013, 104, 1427-1432.	3.9	30