

Jing-Ping Zhao

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/450132/publications.pdf>

Version: 2024-02-01

203
papers

5,623
citations

94433

37
h-index

144013

57
g-index

207
all docs

207
docs citations

207
times ranked

7548
citing authors

#	ARTICLE	IF	CITATIONS
1	Reduced default mode network functional connectivity in patients with recurrent major depressive disorder. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2019, 116, 9078-9083.	7.1	441
2	De novo genic mutations among a Chinese autism spectrum disorder cohort. <i>Nature Communications</i> , 2016, 7, 13316.	12.8	293
3	Emotional Roles of Mono-Aminergic Neurotransmitters in Major Depressive Disorder and Anxiety Disorders. <i>Frontiers in Psychology</i> , 2018, 9, 2201.	2.1	126
4	Minocycline supplementation for treatment of negative symptoms in early-phase schizophrenia: A double blind, randomized, controlled trial. <i>Schizophrenia Research</i> , 2014, 153, 169-176.	2.0	119
5	Inherited and multiple de novo mutations in autism/developmental delay risk genes suggest a multifactorial model. <i>Molecular Autism</i> , 2018, 9, 64.	4.9	114
6	Care for the Psychological Status of Frontline Medical Staff Fighting Against Coronavirus Disease 2019 (COVID-19). <i>Clinical Infectious Diseases</i> , 2020, 71, 3268-3269.	5.8	89
7	Minocycline alleviates behavioral deficits and inhibits microglial activation in the offspring of pregnant mice after administration of polyriboinosinic polyribocytidilic acid. <i>Psychiatry Research</i> , 2014, 219, 680-686.	3.3	87
8	Disrupted intrinsic functional brain topology in patients with major depressive disorder. <i>Molecular Psychiatry</i> , 2021, 26, 7363-7371.	7.9	82
9	Abnormal regional homogeneity as a potential imaging biomarker for adolescent-onset schizophrenia: A resting-state fMRI study and support vector machine analysis. <i>Schizophrenia Research</i> , 2018, 192, 179-184.	2.0	80
10	Minocycline and Risperidone Prevent Microglia Activation and Rescue Behavioral Deficits Induced by Neonatal Intrahippocampal Injection of Lipopolysaccharide in Rats. <i>PLoS ONE</i> , 2014, 9, e93966.	2.5	77
11	Altered resting-state dynamic functional brain networks in major depressive disorder: Findings from the REST-meta-MDD consortium. <i>NeuroImage: Clinical</i> , 2020, 26, 102163.	2.7	76
12	Association Between Maternal Obesity and Autism Spectrum Disorder in Offspring: A Meta-analysis. <i>Journal of Autism and Developmental Disorders</i> , 2016, 46, 95-102.	2.7	73
13	Abnormal Causal Connectivity by Structural Deficits in First-Episode, Drug-Naive Schizophrenia at Rest. <i>Schizophrenia Bulletin</i> , 2015, 41, 57-65.	4.3	71
14	Hippocampal and orbital inferior frontal gray matter volume abnormalities and cognitive deficit in treatment-naive, first-episode patients with schizophrenia. <i>Schizophrenia Research</i> , 2014, 152, 339-343.	2.0	68
15	Decreased insular connectivity in drug-naive major depressive disorder at rest. <i>Journal of Affective Disorders</i> , 2015, 179, 31-37.	4.1	65
16	Functional and anatomical brain deficits in drug-naive major depressive disorder. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2014, 54, 1-6.	4.8	61
17	Shared atypical default mode and salience network functional connectivity between autism and schizophrenia. <i>Autism Research</i> , 2017, 10, 1776-1786.	3.8	60
18	Employment and financial burden of families with preschool children diagnosed with autism spectrum disorders in urban China: results from a descriptive study. <i>BMC Psychiatry</i> , 2015, 15, 3.	2.6	59

#	ARTICLE	IF	CITATIONS
19	Disrupted structural connectivity network in treatment-naïve depression. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2015, 56, 18-26.	4.8	58
20	The effect of minocycline on amelioration of cognitive deficits and pro-inflammatory cytokines levels in patients with schizophrenia. <i>Schizophrenia Research</i> , 2019, 212, 92-98.	2.0	57
21	Abnormal long- and short-range functional connectivity in adolescent-onset schizophrenia patients: A resting-state fMRI study. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2018, 81, 445-451.	4.8	55
22	Abnormal neural activity as a potential biomarker for drug-naïve first-episode adolescent-onset schizophrenia with coherence regional homogeneity and support vector machine analyses. <i>Schizophrenia Research</i> , 2018, 192, 408-415.	2.0	52
23	Neonatal intrahippocampal injection of lipopolysaccharide induces deficits in social behavior and prepulse inhibition and microglial activation in rats: Implication for a new schizophrenia animal model. <i>Brain, Behavior, and Immunity</i> , 2014, 38, 166-174.	4.1	51
24	Biotypes of major depressive disorder: Neuroimaging evidence from resting-state default mode network patterns. <i>NeuroImage: Clinical</i> , 2020, 28, 102514.	2.7	51
25	Genome-wide copy number variation analysis in a Chinese autism spectrum disorder cohort. <i>Scientific Reports</i> , 2017, 7, 44155.	3.3	50
26	Different Visual Preference Patterns in Response to Simple and Complex Dynamic Social Stimuli in Preschool-Aged Children with Autism Spectrum Disorders. <i>PLoS ONE</i> , 2015, 10, e0122280.	2.5	50
27	Brain structural abnormalities as potential markers for detecting individuals with ultra-high risk for psychosis: A systematic review and meta-analysis. <i>Schizophrenia Research</i> , 2019, 209, 22-31.	2.0	49
28	Increased Cerebellar Functional Connectivity With the Default-Mode Network in Unaffected Siblings of Schizophrenia Patients at Rest. <i>Schizophrenia Bulletin</i> , 2015, 41, 1317-1325.	4.3	48
29	Increased short-range and long-range functional connectivity in first-episode, medication-naïve schizophrenia at rest. <i>Schizophrenia Research</i> , 2015, 166, 144-150.	2.0	48
30	Brain-derived neurotrophic factor is associated with cognitive impairments in first-episode and chronic schizophrenia. <i>Psychiatry Research</i> , 2019, 273, 528-536.	3.3	47
31	Patients with first-episode, drug-naïve schizophrenia and subjects at ultra-high risk of psychosis shared increased cerebellar-default mode network connectivity at rest. <i>Scientific Reports</i> , 2016, 6, 26124.	3.3	46
32	Minocycline adjunctive treatment to risperidone for negative symptoms in schizophrenia: Association with pro-inflammatory cytokine levels. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2018, 85, 69-76.	4.8	46
33	Cerebellar structural and functional abnormalities in first-episode and drug-naïve patients with schizophrenia: A meta-analysis. <i>Psychiatry Research - Neuroimaging</i> , 2019, 283, 24-33.	1.8	46
34	Resting-state cerebellar-cerebral networks are differently affected in first-episode, drug-naïve schizophrenia patients and unaffected siblings. <i>Scientific Reports</i> , 2015, 5, 17275.	3.3	45
35	Duration of Untreated Psychosis Is Associated with Temporal and Occipitotemporal Gray Matter Volume Decrease in Treatment Naïve Schizophrenia. <i>PLoS ONE</i> , 2013, 8, e83679.	2.5	44
36	Decreased gray matter volume in the left middle temporal gyrus as a candidate biomarker for schizophrenia: A study of drug naïve, first-episode schizophrenia patients and unaffected siblings. <i>Schizophrenia Research</i> , 2014, 159, 43-50.	2.0	43

#	ARTICLE	IF	CITATIONS
37	Decreased Resting-State Interhemispheric Functional Connectivity Correlated with Neurocognitive Deficits in Drug-Naive First-Episode Adolescent-Onset Schizophrenia. <i>International Journal of Neuropsychopharmacology</i> , 2018, 21, 33-41.	2.1	43
38	Dysfunction in Serotonergic and Noradrenergic Systems and Somatic Symptoms in Psychiatric Disorders. <i>Frontiers in Psychiatry</i> , 2019, 10, 286.	2.6	43
39	Association of genetic variants of GRIN2B with autism. <i>Scientific Reports</i> , 2015, 5, 8296.	3.3	39
40	Profile of minocycline and its potential in the treatment of schizophrenia. <i>Neuropsychiatric Disease and Treatment</i> , 2014, 10, 1103.	2.2	37
41	Decreased interhemispheric functional connectivity in insula and angular gyrus/supramarginal gyrus: Significant findings in first-episode, drug-naive somatization disorder. <i>Psychiatry Research - Neuroimaging</i> , 2016, 248, 48-54.	1.8	37
42	Frequency-specific alteration of functional connectivity density in antipsychotic-naive adolescents with early-onset schizophrenia. <i>Journal of Psychiatric Research</i> , 2017, 95, 68-75.	3.1	37
43	Olanzapine modulation of long- and short-range functional connectivity in the resting brain in a sample of patients with schizophrenia. <i>European Neuropsychopharmacology</i> , 2017, 27, 48-58.	0.7	37
44	Social impairment of children with autism spectrum disorder affects parental quality of life in different ways. <i>Psychiatry Research</i> , 2018, 266, 168-174.	3.3	37
45	Shared and distinct homotopic connectivity changes in melancholic and non-melancholic depression. <i>Journal of Affective Disorders</i> , 2021, 287, 268-275.	4.1	37
46	Altered plasma levels of chemokines in autism and their association with social behaviors. <i>Psychiatry Research</i> , 2016, 244, 300-305.	3.3	35
47	Disruptive variants of <i>CSDE1</i> associate with autism and interfere with neuronal development and synaptic transmission. <i>Science Advances</i> , 2019, 5, eaax2166.	10.3	35
48	Novel Risk Loci Associated With Genetic Risk for Bipolar Disorder Among Han Chinese Individuals. <i>JAMA Psychiatry</i> , 2021, 78, 320.	11.0	35
49	Decreased regional activity and network homogeneity of the fronto-limbic network at rest in drug-naive major depressive disorder. <i>Australian and New Zealand Journal of Psychiatry</i> , 2015, 49, 550-556.	2.3	34
50	Frequency-specific alterations in functional connectivity in treatment-resistant and -sensitive major depressive disorder. <i>Journal of Psychiatric Research</i> , 2016, 82, 30-39.	3.1	34
51	Disrupted asymmetry of inter- and intra-hemispheric functional connectivity in patients with drug-naive, first-episode schizophrenia and their unaffected siblings. <i>EBioMedicine</i> , 2018, 36, 429-435.	6.1	32
52	Unidirectionally affected causal connectivity of cortico-limbic-cerebellar circuit by structural deficits in drug-naive major depressive disorder. <i>Journal of Affective Disorders</i> , 2015, 172, 410-416.	4.1	31
53	Dysfunctional resting-state connectivities of brain regions with structural deficits in drug-naive first-episode schizophrenia adolescents. <i>Schizophrenia Research</i> , 2015, 168, 353-359.	2.0	31
54	A novel de novo POGZ mutation in a patient with intellectual disability. <i>Journal of Human Genetics</i> , 2016, 61, 357-359.	2.3	31

#	ARTICLE	IF	CITATIONS
55	Abnormal default-mode network homogeneity and its correlations with personality in drug-naive somatization disorder at rest. <i>Journal of Affective Disorders</i> , 2016, 193, 81-88.	4.1	31
56	Treatment effects of olanzapine on homotopic connectivity in drug-free schizophrenia at rest. <i>World Journal of Biological Psychiatry</i> , 2018, 19, S106-S114.	2.6	31
57	Dysregulated Maturation of the Functional Connectome in Antipsychotic-Naïve, First-Episode Patients With Adolescent-Onset Schizophrenia. <i>Schizophrenia Bulletin</i> , 2019, 45, 689-697.	4.3	30
58	Effects of Clozapine and other Atypical Antipsychotics on Infants Development Who Were Exposed to as Fetus: A Post-Hoc Analysis. <i>PLoS ONE</i> , 2015, 10, e0123373.	2.5	30
59	A Selective Review of the Excitatory-Inhibitory Imbalance in Schizophrenia: Underlying Biology, Genetics, Microcircuits, and Symptoms. <i>Frontiers in Cell and Developmental Biology</i> , 2021, 9, 664535.	3.7	30
60	An investigation of economic costs of schizophrenia in two areas of China. <i>International Journal of Mental Health Systems</i> , 2013, 7, 26.	2.7	29
61	Cerebellar abnormalities in first-episode, drug-naive schizophrenia at rest. <i>Psychiatry Research - Neuroimaging</i> , 2018, 276, 73-79.	1.8	29
62	Enhanced Global-Brain Functional Connectivity in the Left Superior Frontal Gyus as a Possible Endophenotype for Schizophrenia. <i>Frontiers in Neuroscience</i> , 2019, 13, 145.	2.8	29
63	Phenotype-to-genotype approach reveals headâ€circumferenceâ€associated genes in an autism spectrum disorder cohort. <i>Clinical Genetics</i> , 2020, 97, 338-346.	2.0	29
64	Dissociation of functional and anatomical brain abnormalities in unaffected siblings of schizophrenia patients. <i>Clinical Neurophysiology</i> , 2015, 126, 927-932.	1.5	28
65	Altered Serum Tumor Necrosis Factor and Interleukin-1Î² in First-Episode Drug-Naive and Chronic Schizophrenia. <i>Frontiers in Neuroscience</i> , 2018, 12, 296.	2.8	28
66	Developments in Biological Mechanisms and Treatments for Negative Symptoms and Cognitive Dysfunction of Schizophrenia. <i>Neuroscience Bulletin</i> , 2021, 37, 1609-1624.	2.9	28
67	Identification of RELN variation p.Thr3192Ser in a Chinese family with schizophrenia. <i>Scientific Reports</i> , 2016, 6, 24327.	3.3	27
68	Tranlycypromine Causes Neurotoxicity and Represses BHC110/LSD1 in Human-Induced Pluripotent Stem Cell-Derived Cerebral Organoids Model. <i>Frontiers in Neurology</i> , 2017, 8, 626.	2.4	27
69	Efficacy, acceptability and tolerability of 8 atypical antipsychotics in Chinese patients with acute schizophrenia: A network meta-analysis. <i>Schizophrenia Research</i> , 2017, 185, 73-79.	2.0	26
70	Regional white matter abnormalities in drug-naive, first-episode schizophrenia patients and their healthy unaffected siblings. <i>Australian and New Zealand Journal of Psychiatry</i> , 2015, 49, 246-254.	2.3	25
71	Abnormal regional homogeneity as potential imaging biomarker for psychosis risk syndrome: a resting-state fMRI study and support vector machine analysis. <i>Scientific Reports</i> , 2016, 6, 27619.	3.3	25
72	Abnormal functional connectivity strength in patients with adolescent-onset schizophrenia: a resting-state fMRI study. <i>European Child and Adolescent Psychiatry</i> , 2017, 26, 839-845.	4.7	25

#	ARTICLE	IF	CITATIONS
73	Voxel-wise brain-wide functional connectivity abnormalities in first-episode, drug-naive patients with major depressive disorder. <i>American Journal of Medical Genetics Part B: Neuropsychiatric Genetics</i> , 2018, 177, 447-453.	1.7	25
74	Altered dynamic global signal topography in antipsychotic-naive adolescents with early-onset schizophrenia. <i>Schizophrenia Research</i> , 2019, 208, 308-316.	2.0	25
75	Increased Appetite Plays a Key Role in Olanzapine-Induced Weight Gain in First-Episode Schizophrenia Patients. <i>Frontiers in Pharmacology</i> , 2020, 11, 739.	3.5	25
76	Mental health and burnout in primary and secondary school teachers in the remote mountain areas of Guangdong Province in the People's Republic of China. <i>Neuropsychiatric Disease and Treatment</i> , 2014, 10, 123.	2.2	24
77	ZNF804A rs1344706 is associated with cortical thickness, surface area, and cortical volume of the unmedicated first episode schizophrenia and healthy controls. <i>American Journal of Medical Genetics Part B: Neuropsychiatric Genetics</i> , 2015, 168, 265-273.	1.7	24
78	Disrupted amplitude of low-frequency fluctuations in antipsychotic-naïve adolescents with early-onset schizophrenia. <i>Psychiatry Research - Neuroimaging</i> , 2016, 249, 20-26.	1.8	24
79	Enhanced baseline activity in the left ventromedial putamen predicts individual treatment response in drug-naive, first-episode schizophrenia: Results from two independent study samples. <i>EBioMedicine</i> , 2019, 46, 248-255.	6.1	24
80	An open-label randomised comparison of aripiprazole, olanzapine and risperidone for the acute treatment of first-episode schizophrenia: Eight-week outcomes. <i>Journal of Psychopharmacology</i> , 2019, 33, 1227-1236.	4.0	24
81	Functional asymmetry of thalamocortical networks in subjects at ultra-high risk for psychosis and first-episode schizophrenia. <i>European Neuropsychopharmacology</i> , 2019, 29, 519-528.	0.7	24
82	Reduced connectivity in anterior cingulate cortex as an early predictor for treatment response in drug-naive, first-episode schizophrenia: A global-brain functional connectivity analysis. <i>Schizophrenia Research</i> , 2020, 215, 337-343.	2.0	24
83	Public behavior change, perceptions, depression, and anxiety in relation to the COVID-19 outbreak. <i>Open Forum Infectious Diseases</i> , 2020, 7, ofaa273.	0.9	24
84	Decreased interhemispheric coordination in the posterior default-mode network and visual regions as trait alterations in first-episode, drug-naive major depressive disorder. <i>Brain Imaging and Behavior</i> , 2018, 12, 1251-1258.	2.1	23
85	The Wnt Signaling Pathway Effector TCF7L2 Mediates Olanzapine-Induced Weight Gain and Insulin Resistance. <i>Frontiers in Pharmacology</i> , 2018, 9, 379.	3.5	23
86	Associations among maternal pre-pregnancy body mass index, gestational weight gain and risk of autism in the Han Chinese population. <i>BMC Psychiatry</i> , 2018, 18, 11.	2.6	23
87	Disrupted hemispheric connectivity specialization in patients with major depressive disorder: Evidence from the REST-meta-MDD Project. <i>Journal of Affective Disorders</i> , 2021, 284, 217-228.	4.1	23
88	Family-based case-control study of homotopic connectivity in first-episode, drug-naive schizophrenia at rest. <i>Scientific Reports</i> , 2017, 7, 43312.	3.3	22
89	Maternal dietary patterns, supplements intake and autism spectrum disorders. <i>Medicine (United Tj ETQq1 1 0.784314 rgBT /Overloc</i>	1.0	22
90	The Effects of Clozapine on Cognitive Function and Regional Cerebral Blood Flow in the Negative Symptom Profile Schizophrenia. <i>International Journal of Psychiatry in Medicine</i> , 2006, 36, 171-181.	1.8	21

#	ARTICLE	IF	CITATIONS
91	Relationship between long-term use of a typical antipsychotic medication by Chinese schizophrenia patients and the bone turnover markers serum osteocalcin and β -CrossLaps. <i>Schizophrenia Research</i> , 2016, 176, 259-263.	2.0	21
92	Decreased long- and short-range functional connectivity at rest in drug-naive major depressive disorder. <i>Australian and New Zealand Journal of Psychiatry</i> , 2016, 50, 763-769.	2.3	21
93	Olanzapine modulates the default-mode network homogeneity in recurrent drug-free schizophrenia at rest. <i>Australian and New Zealand Journal of Psychiatry</i> , 2017, 51, 1000-1009.	2.3	21
94	Altered functional connectivity strength and its correlations with cognitive function in subjects with ultra-high risk for psychosis at rest. <i>CNS Neuroscience and Therapeutics</i> , 2018, 24, 1140-1148.	3.9	21
95	Reduced Brain Activity in the Right Putamen as an Early Predictor for Treatment Response in Drug-Naive, First-Episode Schizophrenia. <i>Frontiers in Psychiatry</i> , 2019, 10, 741.	2.6	20
96	Cognitive deficits in subjects at risk for psychosis, first-episode and chronic schizophrenia patients. <i>Psychiatry Research</i> , 2019, 274, 235-242.	3.3	20
97	Voxel-based global-brain functional connectivity alterations in first-episode drug-naive patients with somatization disorder. <i>Journal of Affective Disorders</i> , 2019, 254, 82-89.	4.1	20
98	Cognitive effects of atypical antipsychotic drugs in first-episode drug-naïve schizophrenic patients. <i>Neural Regeneration Research</i> , 2013, 8, 277-86.	3.0	20
99	Insertion of a knockout-first cassette in <i>Ampd1</i> gene leads to neonatal death by disruption of neighboring genes expression. <i>Scientific Reports</i> , 2016, 6, 35970.	3.3	19
100	Increased anterior default-mode network homogeneity in first-episode, drug-naive major depressive disorder: A replication study. <i>Journal of Affective Disorders</i> , 2018, 225, 767-772.	4.1	19
101	Reduced Hippocampal Volume and Its Relationship With Verbal Memory and Negative Symptoms in Treatment-Naive First-Episode Adolescent-Onset Schizophrenia. <i>Schizophrenia Bulletin</i> , 2021, 47, 64-74.	4.3	19
102	Disrupted Regional Homogeneity in Melancholic and Non-melancholic Major Depressive Disorder at Rest. <i>Frontiers in Psychiatry</i> , 2021, 12, 618805.	2.6	19
103	Associations Between Internet Addiction and Gender, Anxiety, Coping Styles and Acceptance in University Freshmen in South China. <i>Frontiers in Psychiatry</i> , 2021, 12, 558080.	2.6	19
104	Using short-range and long-range functional connectivity to identify schizophrenia with a family-based case-control design. <i>Psychiatry Research - Neuroimaging</i> , 2017, 264, 60-67.	1.8	18
105	Rare inherited missense variants of <i>POGZ</i> associate with autism risk and disrupt neuronal development. <i>Journal of Genetics and Genomics</i> , 2019, 46, 247-257.	3.9	17
106	Abnormal default-mode network homogeneity and its correlations with neurocognitive deficits in drug-naive first-episode adolescent-onset schizophrenia. <i>Schizophrenia Research</i> , 2020, 215, 140-147.	2.0	17
107	Individual-specific functional connectome biomarkers predict schizophrenia positive symptoms during adolescent brain maturation. <i>Human Brain Mapping</i> , 2021, 42, 1475-1484.	3.6	17
108	Abnormal ACTH and prolactin responses to fenfluramine in rats exposed to single and multiple doses of MDMA. <i>Psychopharmacology</i> , 1997, 131, 411-419.	3.1	16

#	ARTICLE	IF	CITATIONS
109	Bidirectional Causal Connectivity in the Cortico-Limbic-Cerebellar Circuit Related to Structural Alterations in First-Episode, Drug-Naive Somatization Disorder. <i>Frontiers in Psychiatry</i> , 2018, 9, 162.	2.6	16
110	Increased frontal gray matter volume in individuals with prodromal psychosis. <i>CNS Neuroscience and Therapeutics</i> , 2019, 25, 987-994.	3.9	16
111	Genome-wide association analysis of autism identified multiple loci that have been reported as strong signals for neuropsychiatric disorders. <i>Autism Research</i> , 2020, 13, 382-396.	3.8	16
112	Mapping Progressive Gray Matter Alterations in Early Childhood Autistic Brain. <i>Cerebral Cortex</i> , 2021, 31, 1500-1510.	2.9	16
113	Probiotics Plus Dietary Fiber Supplements Attenuate Olanzapine-Induced Weight Gain in Drug-Naïve First-Episode Schizophrenia Patients: Two Randomized Clinical Trials. <i>Schizophrenia Bulletin</i> , 2022, 48, 850-859.	4.3	16
114	Gender differences of neurocognitive functioning in patients with first-episode schizophrenia in China. <i>Comprehensive Psychiatry</i> , 2019, 95, 152132.	3.1	15
115	Citalopram in first episode schizophrenia: The DECIFER trial. <i>Schizophrenia Research</i> , 2019, 208, 331-337.	2.0	15
116	Sleep Problems of Children with Autism May Independently Affect Parental Quality of Life. <i>Child Psychiatry and Human Development</i> , 2021, 52, 488-499.	1.9	15
117	Effect of Bifidobacterium on olanzapine-induced body weight and appetite changes in patients with psychosis. <i>Psychopharmacology</i> , 2021, 238, 2449-2457.	3.1	15
118	APOA-I: A Possible Novel Biomarker for Metabolic Side Effects in First Episode Schizophrenia. <i>PLoS ONE</i> , 2014, 9, e93902.	2.5	14
119	Aggressive behaviors and treatable risk factors of preschool children with autism spectrum disorder. <i>Autism Research</i> , 2017, 10, 1155-1162.	3.8	14
120	Autism spectrum disorder and severe social impairment associated with elevated plasma interleukin-8. <i>Pediatric Research</i> , 2021, 89, 591-597.	2.3	14
121	Abnormal Default-Mode Network Homogeneity in Melancholic and Nonmelancholic Major Depressive Disorder at Rest. <i>Neural Plasticity</i> , 2021, 2021, 1-12.	2.2	14
122	Shuganjieyu capsule increases neurotrophic factor expression in a rat model of depression. <i>Neural Regeneration Research</i> , 2014, 9, 489.	3.0	14
123	Case-control association study of ABCB1 gene and major depressive disorder in a local Chinese Han population. <i>Neuropsychiatric Disease and Treatment</i> , 2015, 11, 1967.	2.2	13
124	Broad autism phenotype features of Chinese parents with autistic children and their associations with severity of social impairment in probands. <i>BMC Psychiatry</i> , 2015, 15, 168.	2.6	13
125	POGZ de novo missense variants in neuropsychiatric disorders. <i>Molecular Genetics & Genomic Medicine</i> , 2019, 7, e900.	1.2	13
126	Prevalence, clinical correlates and IQ of suicidal ideation in drug naïve Chinese Han patients with major depressive disorder. <i>Journal of Affective Disorders</i> , 2019, 248, 59-64.	4.1	13

#	ARTICLE	IF	CITATIONS
127	Disrupted Regional Homogeneity in Major Depressive Disorder With Gastrointestinal Symptoms at Rest. <i>Frontiers in Psychiatry</i> , 2021, 12, 636820.	2.6	13
128	Increased coherence-based regional homogeneity in resting-state patients with first-episode, drug-naive somatization disorder. <i>Journal of Affective Disorders</i> , 2018, 235, 150-154.	4.1	12
129	Cost-effectiveness of olanzapine in the first-line treatment of schizophrenia in China. <i>Journal of Medical Economics</i> , 2019, 22, 439-446.	2.1	12
130	Effect of CYP2D6 polymorphisms on plasma concentration and therapeutic effect of risperidone. <i>BMC Psychiatry</i> , 2021, 21, 70.	2.6	12
131	Anatomical distance affects cortical-subcortical connectivity in first-episode, drug-naive somatization disorder. <i>Journal of Affective Disorders</i> , 2017, 217, 153-158.	4.1	11
132	Rates and predictors of one-year antipsychotic treatment discontinuation in first-episode schizophrenia: Results from an open-label, randomized, "real world" clinical trial. <i>Psychiatry Research</i> , 2019, 273, 631-640.	3.3	11
133	Increased regional homogeneity modulated by metacognitive training predicts therapeutic efficacy in patients with schizophrenia. <i>European Archives of Psychiatry and Clinical Neuroscience</i> , 2021, 271, 783-798.	3.2	11
134	Guidelines Disconcordance in Acute Bipolar Depression: Data from the National Bipolar Mania Pathway Survey (BIPAS) in Mainland China. <i>PLoS ONE</i> , 2014, 9, e96096.	2.5	11
135	Altered regional homogeneity and cognitive impairments in first-episode schizophrenia: A resting-state fMRI study. <i>Asian Journal of Psychiatry</i> , 2022, 71, 103055.	2.0	11
136	The effects of probiotics plus dietary fiber on antipsychotic-induced weight gain: a randomized clinical trial. <i>Translational Psychiatry</i> , 2022, 12, 185.	4.8	11
137	Adjunctive treatment for cognitive impairment in patients with chronic schizophrenia: a double-blind, placebo-controlled study. <i>Neuropsychiatric Disease and Treatment</i> , 2014, 10, 1317.	2.2	10
138	The comparison of glucose and lipid metabolism parameters in drug-naïve, antipsychotic-treated, and antipsychotic discontinuation patients with schizophrenia. <i>Neuropsychiatric Disease and Treatment</i> , 2014, 10, 1361.	2.2	10
139	Dissociation of anatomical and functional alterations of the default-mode network in first-episode, drug-naive schizophrenia. <i>Clinical Neurophysiology</i> , 2015, 126, 2276-2281.	1.5	10
140	Forkhead box C1 is targeted by microRNA-133b and promotes cell proliferation and migration in osteosarcoma. <i>Experimental and Therapeutic Medicine</i> , 2017, 14, 2823-2830.	1.8	10
141	Safety and efficacy of paliperidone palmitate 1-month formulation in Chinese patients with schizophrenia: a 25-week, open-label, multicenter, Phase IV study. <i>Neuropsychiatric Disease and Treatment</i> , 2017, Volume 13, 2045-2056.	2.2	10
142	Changes in plasma levels of nitric oxide metabolites and negative symptoms after 16-week minocycline treatment in patients with schizophrenia. <i>Schizophrenia Research</i> , 2018, 199, 390-394.	2.0	10
143	Decreased white matter FA values in the left inferior frontal gyrus is a possible intermediate phenotype of schizophrenia: evidences from a novel group strategy. <i>European Archives of Psychiatry and Clinical Neuroscience</i> , 2018, 268, 89-98.	3.2	10
144	Frequency-specific altered global signal topography in drug-naïve first-episode patients with adolescent-onset schizophrenia. <i>Brain Imaging and Behavior</i> , 2021, 15, 1876-1885.	2.1	10

#	ARTICLE	IF	CITATIONS
145	Application of Artificial Intelligence on Psychological Interventions and Diagnosis: An Overview. <i>Frontiers in Psychiatry</i> , 2022, 13, 811665.	2.6	10
146	A comparative study of magnetic resonance imaging on the gray matter and resting-state function in prodromal and first-episode schizophrenia. <i>American Journal of Medical Genetics Part B: Neuropsychiatric Genetics</i> , 2018, 177, 537-545.	1.7	9
147	Increased Nucleus Accumbens Connectivity in Resting-State Patients With Drug-Naive, First-Episode Somatization Disorder. <i>Frontiers in Psychiatry</i> , 2019, 10, 585.	2.6	9
148	Resting-state functional hypoconnectivity of amygdala in clinical high risk state and first-episode schizophrenia. <i>Brain Imaging and Behavior</i> , 2020, 14, 1840-1849.	2.1	9
149	Genotype and Phenotype Correlations for TBL1XR1 in Neurodevelopmental Disorders. <i>Journal of Molecular Neuroscience</i> , 2020, 70, 2085-2092.	2.3	9
150	Hippocampal functional network: The mediating role between obsession and anxiety in adult patients with obsessive-compulsive disorder. <i>World Journal of Biological Psychiatry</i> , 2020, 21, 685-695.	2.6	9
151	Metacognitive Training Modulates Default-Mode Network Homogeneity During 8-Week Olanzapine Treatment in Patients With Schizophrenia. <i>Frontiers in Psychiatry</i> , 2020, 11, 234.	2.6	9
152	The similar eye movement dysfunction between major depressive disorder, bipolar depression and bipolar mania. <i>World Journal of Biological Psychiatry</i> , 2022, 23, 689-702.	2.6	9
153	Paliperidone extended-release tablets in Chinese patients with schizophrenia: meta-analysis of randomized controlled trials. <i>Neuropsychiatric Disease and Treatment</i> , 2015, 11, 1817.	2.2	8
154	Metformin attenuates antipsychotic-induced metabolic dysfunctions in MK801-induced schizophrenia-like rats. <i>Psychopharmacology</i> , 2020, 237, 2257-2277.	3.1	8
155	Covariation between Childhood-Trauma Related Resting-State Functional Connectivity and Affective Temperaments is Impaired in Individuals with Major Depressive Disorder. <i>Neuroscience</i> , 2021, 453, 102-112.	2.3	8
156	Optimizing and Individualizing the Pharmacological Treatment of First-Episode Schizophrenic Patients: Study Protocol for a Multicenter Clinical Trial. <i>Frontiers in Psychiatry</i> , 2021, 12, 611070.	2.6	8
157	Increased Homotopic Connectivity in the Prefrontal Cortex Modulated by Olanzapine Predicts Therapeutic Efficacy in Patients with Schizophrenia. <i>Neural Plasticity</i> , 2021, 2021, 1-11.	2.2	8
158	Abnormal interhemispheric homotopic functional connectivity is correlated with gastrointestinal symptoms in patients with major depressive disorder. <i>Journal of Psychiatric Research</i> , 2021, 144, 234-240.	3.1	8
159	Region-specific insular volumetric decreases in drug-naive, first-episode schizophrenia and their unaffected siblings. <i>American Journal of Medical Genetics Part B: Neuropsychiatric Genetics</i> , 2020, 183, 106-112.	1.7	7
160	Frontal white matter abnormalities reveal the pathological basis underlying negative symptoms in antipsychotic-naïve, first-episode patients with adolescent-onset schizophrenia: Evidence from multimodal brain imaging. <i>Schizophrenia Research</i> , 2020, 222, 258-266.	2.0	7
161	Air pollution and hippocampal atrophy in first episode schizophrenia. <i>Schizophrenia Research</i> , 2020, 218, 63-69.	2.0	7
162	Effects of Sleep Disturbances on Behavioral Problems in Preschool Children With Autism Spectrum Disorder. <i>Frontiers in Psychiatry</i> , 2020, 11, 559694.	2.6	7

#	ARTICLE	IF	CITATIONS
163	Gene expression of methylation cycle and related genes in lymphocytes and brain of patients with schizophrenia and non-psychotic controls. <i>Biomarkers in Neuropsychiatry</i> , 2021, 5, 100038.	1.0	7
164	Abnormal global-brain functional connectivity and its relationship with cognitive deficits in drug-naive first-episode adolescent-onset schizophrenia. <i>Brain Imaging and Behavior</i> , 2022, , 1.	2.1	7
165	Guidelines concordance of maintenance treatment in euthymic patients with bipolar disorder: Data from the national bipolar mania pathway survey (BIPAS) in mainland China. <i>Journal of Affective Disorders</i> , 2015, 182, 101-105.	4.1	6
166	Regional white matter volume abnormalities in first-episode somatization disorder. <i>International Journal of Psychophysiology</i> , 2018, 133, 12-16.	1.0	6
167	Effects of Aspirin in Rats With Ouabain Intracerebral Treatment—Possible Involvement of Inflammatory Modulation?. <i>Frontiers in Psychiatry</i> , 2019, 10, 497.	2.6	6
168	<p>Pleiotropic action of genetic variation in ZNF804A on brain structure: a meta-analysis of magnetic resonance imaging studies</p>. <i>Neuropsychiatric Disease and Treatment</i> , 2019, Volume 15, 721-729.	2.2	6
169	Less reduced gray matter volume in the subregions of superior temporal gyrus predicts better treatment efficacy in drug-naive, first-episode schizophrenia. <i>Brain Imaging and Behavior</i> , 2021, 15, 1997-2004.	2.1	6
170	Anesthesia, sex and miscarriage history may influence the association between cesarean delivery and autism spectrum disorder. <i>BMC Pediatrics</i> , 2021, 21, 62.	1.7	6
171	Dietary fiber and probiotics for the treatment of atypical antipsychotic-induced metabolic side effects: study protocol for a randomized, double-blind, placebo-controlled trial. <i>Trials</i> , 2021, 22, 159.	1.6	6
172	Shared and Distinct Fractional Amplitude of Low-Frequency Fluctuation Patterns in Major Depressive Disorders With and Without Gastrointestinal Symptoms. <i>Frontiers in Psychiatry</i> , 2021, 12, 744898.	2.6	6
173	PCSK9 mediates dyslipidemia induced by olanzapine treatment in schizophrenia patients. <i>Psychopharmacology</i> , 2022, 239, 83-91.	3.1	6
174	Clinical utility of orally disintegrating olanzapine in Chinese patients with schizophrenia: a review of effectiveness, patient preference, adherence, and other properties. <i>Neuropsychiatric Disease and Treatment</i> , 2014, 10, 355.	2.2	5
175	Chinese Taoist Cognitive Therapy for Symptoms of Depression and Anxiety in Adults in China: A Systematic Review and Meta-Analysis. <i>Frontiers in Psychology</i> , 2020, 11, 769.	2.1	5
176	The Association Between Cognitive Deficits and Clinical Characteristic in First-Episode Drug Naïve Patients With Schizophrenia. <i>Frontiers in Psychiatry</i> , 2021, 12, 638773.	2.6	5
177	An Overview of Genetic and Environmental Risk of Autism Spectrum Disorder. <i>Global Clinical and Translational Research</i> , 2019, , 37-44.	0.3	5
178	Abnormal Default Mode Network Homogeneity in Major Depressive Disorder With Gastrointestinal Symptoms at Rest. <i>Frontiers in Aging Neuroscience</i> , 2022, 14, 804621.	3.4	5
179	Odor identification impairment in autism spectrum disorder might be associated with mitochondrial dysfunction. <i>Asian Journal of Psychiatry</i> , 2022, 72, 103072.	2.0	5
180	Enhanced Connectivity of Thalamo-Cortical Networks in First-Episode, Treatment-Naive Somatization Disorder. <i>Frontiers in Psychiatry</i> , 2020, 11, 555836.	2.6	4

#	ARTICLE	IF	CITATIONS
181	The Association Between Metabolic Disturbance and Cognitive Impairments in Early-Stage Schizophrenia. <i>Frontiers in Human Neuroscience</i> , 2020, 14, 599720.	2.0	4
182	Reduction of phonetic mismatch negativity may depict illness course and predict functional outcomes in schizophrenia. <i>Journal of Psychiatric Research</i> , 2021, 137, 290-297.	3.1	4
183	Differentiating Melancholic and Non-melancholic Major Depressive Disorder Using Fractional Amplitude of Low-Frequency Fluctuations. <i>Frontiers in Psychiatry</i> , 2021, 12, 763770.	2.6	4
184	Differences among first-episode schizophrenia patients, healthy siblings, and controls at the individual level. <i>International Journal of Psychophysiology</i> , 2016, 104, 24-32.	1.0	3
185	Predictors of menstruation restoration during metformin administration for treatment of antipsychotic drug-induced amenorrhea: A post hoc analysis. <i>Schizophrenia Research</i> , 2017, 190, 121-122.	2.0	3
186	Which Subgroup of First-Episode Schizophrenia Patients Can Remit During the First Year of Antipsychotic Treatment?. <i>Frontiers in Psychiatry</i> , 2020, 11, 566.	2.6	3
187	Face and object visual working memory deficits in first-episode schizophrenia correlate with multiple neurocognitive performances. <i>Annals of General Psychiatry</i> , 2021, 34, e100338.	3.1	3
188	Association of Aripiprazole With Reduced Hippocampal Atrophy During Maintenance Treatment of First-Episode Schizophrenia. <i>Journal of Clinical Psychopharmacology</i> , 2021, 41, 244-249.	1.4	3
189	Abnormal white matter functional connectivity density in antipsychotic-naïve adolescents with schizophrenia. <i>Clinical Neurophysiology</i> , 2021, 132, 1025-1032.	1.5	3
190	Effect of citalopram on hippocampal volume in first-episode schizophrenia: Structural MRI results from the DECIFER trial. <i>Psychiatry Research - Neuroimaging</i> , 2021, 312, 111286.	1.8	3
191	Increased subcortical region volume induced by electroconvulsive therapy in patients with schizophrenia. <i>European Archives of Psychiatry and Clinical Neuroscience</i> , 2021, 271, 1285-1295.	3.2	3
192	Prenatal Environment and Perinatal Factors Associated with Autism Spectrum Disorder. <i>Global Clinical and Translational Research</i> , 2019, , 100-108.	0.3	3
193	Sulforaphane Effects on Cognition and Symptoms in First and Early Episode Schizophrenia: A Randomized Double-Blind Trial. <i>Schizophrenia Bulletin Open</i> , 2022, 3, .	1.7	3
194	Altered Brain Functional Asymmetry in Patients With Major Depressive Disorder Related to Gastrointestinal Symptoms. <i>Frontiers in Neuroscience</i> , 2021, 15, 797598.	2.8	3
195	Aberrant corticostriatal connectivity predict positive symptoms of antipsychotic-naïve patients with adolescent-onset schizophrenia during brain maturation. <i>Schizophrenia Research</i> , 2018, 195, 564-566.	2.0	2
196	Clinical characteristics of familial schizophrenia. <i>Asia-Pacific Psychiatry</i> , 2021, 13, e12422.	2.2	2
197	Serum Interleukin-1 β and tumor necrosis factor- α in first-episode drug-naïve and chronic schizophrenia patients: Associated with cognitive deficits. <i>Asian Journal of Psychiatry</i> , 2021, 58, 102605.	2.0	2
198	Relationship between brain-derived neurotrophic factor gene C270T polymorphisms and the psychotic symptoms and cognitive functioning of patients with schizophrenia. <i>Shanghai Archives of Psychiatry</i> , 2012, 24, 328-34.	0.7	2

#	ARTICLE	IF	CITATIONS
199	Previous exposure to antipsychotic drug treatment is an effective predictor of metabolic disturbances experienced with current antipsychotic drug treatments. <i>BMC Psychiatry</i> , 2022, 22, 210.	2.6	2
200	Cortical Pathways or Mechanism in the Face Inversion Effect in Patients with First-Episode Schizophrenia. <i>Neuropsychiatric Disease and Treatment</i> , 2021, Volume 17, 1893-1906.	2.2	1
201	Amplitude of Low-Frequency Oscillations in First-Episode Drug-Naive Patients with Major Depressive Disorder: A Resting State Functional Magnetic Resonance Imaging Study. <i>Neuropsychiatric Disease and Treatment</i> , 2022, Volume 18, 555-561.	2.2	1
202	Challenges in the early detection and intervention of the psychosis-risk syndrome. <i>Shanghai Archives of Psychiatry</i> , 2015, 27, 45-7.	0.7	0
203	Research advances in add-on treatment for negative symptoms and cognitive dysfunction in schizophrenia. <i>Journal of Central South University (Medical Sciences)</i> , 2020, 45, 1457-1463.	0.1	0