

Jing-Ping Zhao

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/450132/publications.pdf>

Version: 2024-02-01

203
papers

5,623
citations

94433

37
h-index

144013

57
g-index

207
all docs

207
docs citations

207
times ranked

7548
citing authors

#	ARTICLE	IF	CITATIONS
1	Abnormal global-brain functional connectivity and its relationship with cognitive deficits in drug-naive first-episode adolescent-onset schizophrenia. <i>Brain Imaging and Behavior</i> , 2022, , 1.	2.1	7
2	PCSK9 mediates dyslipidemia induced by olanzapine treatment in schizophrenia patients. <i>Psychopharmacology</i> , 2022, 239, 83-91.	3.1	6
3	The similar eye movement dysfunction between major depressive disorder, bipolar depression and bipolar mania. <i>World Journal of Biological Psychiatry</i> , 2022, 23, 689-702.	2.6	9
4	Sulforaphane Effects on Cognition and Symptoms in First and Early Episode Schizophrenia: A Randomized Double-Blind Trial. <i>Schizophrenia Bulletin Open</i> , 2022, 3, .	1.7	3
5	Previous exposure to antipsychotic drug treatment is an effective predictor of metabolic disturbances experienced with current antipsychotic drug treatments. <i>BMC Psychiatry</i> , 2022, 22, 210.	2.6	2
6	Amplitude of Low-Frequency Oscillations in First-Episode Drug-Naive Patients with Major Depressive Disorder: A Resting State Functional Magnetic Resonance Imaging Study. <i>Neuropsychiatric Disease and Treatment</i> , 2022, Volume 18, 555-561.	2.2	1
7	Application of Artificial Intelligence on Psychological Interventions and Diagnosis: An Overview. <i>Frontiers in Psychiatry</i> , 2022, 13, 811665.	2.6	10
8	Abnormal Default Mode Network Homogeneity in Major Depressive Disorder With Gastrointestinal Symptoms at Rest. <i>Frontiers in Aging Neuroscience</i> , 2022, 14, 804621.	3.4	5
9	Altered regional homogeneity and cognitive impairments in first-episode schizophrenia: A resting-state fMRI study. <i>Asian Journal of Psychiatry</i> , 2022, 71, 103055.	2.0	11
10	Odor identification impairment in autism spectrum disorder might be associated with mitochondrial dysfunction. <i>Asian Journal of Psychiatry</i> , 2022, 72, 103072.	2.0	5
11	Probiotics Plus Dietary Fiber Supplements Attenuate Olanzapine-Induced Weight Gain in Drug-Naive First-Episode Schizophrenia Patients: Two Randomized Clinical Trials. <i>Schizophrenia Bulletin</i> , 2022, 48, 850-859.	4.3	16
12	The effects of probiotics plus dietary fiber on antipsychotic-induced weight gain: a randomized clinical trial. <i>Translational Psychiatry</i> , 2022, 12, 185.	4.8	11
13	Frequency-specific altered global signal topography in drug-naive first-episode patients with adolescent-onset schizophrenia. <i>Brain Imaging and Behavior</i> , 2021, 15, 1876-1885.	2.1	10
14	Less reduced gray matter volume in the subregions of superior temporal gyrus predicts better treatment efficacy in drug-naive, first-episode schizophrenia. <i>Brain Imaging and Behavior</i> , 2021, 15, 1997-2004.	2.1	6
15	Increased regional homogeneity modulated by metacognitive training predicts therapeutic efficacy in patients with schizophrenia. <i>European Archives of Psychiatry and Clinical Neuroscience</i> , 2021, 271, 783-798.	3.2	11
16	Autism spectrum disorder and severe social impairment associated with elevated plasma interleukin-8. <i>Pediatric Research</i> , 2021, 89, 591-597.	2.3	14
17	Covariation between Childhood-Trauma Related Resting-State Functional Connectivity and Affective Temperaments is Impaired in Individuals with Major Depressive Disorder. <i>Neuroscience</i> , 2021, 453, 102-112.	2.3	8
18	Reduced Hippocampal Volume and Its Relationship With Verbal Memory and Negative Symptoms in Treatment-Naive First-Episode Adolescent-Onset Schizophrenia. <i>Schizophrenia Bulletin</i> , 2021, 47, 64-74.	4.3	19

#	ARTICLE	IF	CITATIONS
19	Sleep Problems of Children with Autism May Independently Affect Parental Quality of Life. <i>Child Psychiatry and Human Development</i> , 2021, 52, 488-499.	1.9	15
20	Clinical characteristics of familial schizophrenia. <i>Asia-Pacific Psychiatry</i> , 2021, 13, e12422.	2.2	2
21	Novel Risk Loci Associated With Genetic Risk for Bipolar Disorder Among Han Chinese Individuals. <i>JAMA Psychiatry</i> , 2021, 78, 320.	11.0	35
22	Individual-specific functional connectome biomarkers predict schizophrenia positive symptoms during adolescent brain maturation. <i>Human Brain Mapping</i> , 2021, 42, 1475-1484.	3.6	17
23	Mapping Progressive Gray Matter Alterations in Early Childhood Autistic Brain. <i>Cerebral Cortex</i> , 2021, 31, 1500-1510.	2.9	16
24	Optimizing and Individualizing the Pharmacological Treatment of First-Episode Schizophrenic Patients: Study Protocol for a Multicenter Clinical Trial. <i>Frontiers in Psychiatry</i> , 2021, 12, 611070.	2.6	8
25	The Association Between Cognitive Deficits and Clinical Characteristic in First-Episode Drug Naïve Patients With Schizophrenia. <i>Frontiers in Psychiatry</i> , 2021, 12, 638773.	2.6	5
26	Disrupted Regional Homogeneity in Melancholic and Non-melancholic Major Depressive Disorder at Rest. <i>Frontiers in Psychiatry</i> , 2021, 12, 618805.	2.6	19
27	Anesthesia, sex and miscarriage history may influence the association between cesarean delivery and autism spectrum disorder. <i>BMC Pediatrics</i> , 2021, 21, 62.	1.7	6
28	Dietary fiber and probiotics for the treatment of atypical antipsychotic-induced metabolic side effects: study protocol for a randomized, double-blind, placebo-controlled trial. <i>Trials</i> , 2021, 22, 159.	1.6	6
29	Face and object visual working memory deficits in first-episode schizophrenia correlate with multiple neurocognitive performances. <i>Annals of General Psychiatry</i> , 2021, 34, e100338.	3.1	3
30	Effect of CYP2D6 polymorphisms on plasma concentration and therapeutic effect of risperidone. <i>BMC Psychiatry</i> , 2021, 21, 70.	2.6	12
31	Disrupted hemispheric connectivity specialization in patients with major depressive disorder: Evidence from the REST-meta-MDD Project. <i>Journal of Affective Disorders</i> , 2021, 284, 217-228.	4.1	23
32	Abnormal Default-Mode Network Homogeneity in Melancholic and Nonmelancholic Major Depressive Disorder at Rest. <i>Neural Plasticity</i> , 2021, 2021, 1-12.	2.2	14
33	Association of Aripiprazole With Reduced Hippocampal Atrophy During Maintenance Treatment of First-Episode Schizophrenia. <i>Journal of Clinical Psychopharmacology</i> , 2021, 41, 244-249.	1.4	3
34	Serum Interleukin-1 β and tumor necrosis factor- α in first-episode drug-naive and chronic schizophrenia patients: Associated with cognitive deficits. <i>Asian Journal of Psychiatry</i> , 2021, 58, 102605.	2.0	2
35	Shared and distinct homotopic connectivity changes in melancholic and non-melancholic depression. <i>Journal of Affective Disorders</i> , 2021, 287, 268-275.	4.1	37
36	Abnormal white matter functional connectivity density in antipsychotic-naive adolescents with schizophrenia. <i>Clinical Neurophysiology</i> , 2021, 132, 1025-1032.	1.5	3

#	ARTICLE	IF	CITATIONS
37	Reduction of phonetic mismatch negativity may depict illness course and predict functional outcomes in schizophrenia. <i>Journal of Psychiatric Research</i> , 2021, 137, 290-297.	3.1	4
38	Associations Between Internet Addiction and Gender, Anxiety, Coping Styles and Acceptance in University Freshmen in South China. <i>Frontiers in Psychiatry</i> , 2021, 12, 558080.	2.6	19
39	Disrupted Regional Homogeneity in Major Depressive Disorder With Gastrointestinal Symptoms at Rest. <i>Frontiers in Psychiatry</i> , 2021, 12, 636820.	2.6	13
40	Effect of Bifidobacterium on olanzapine-induced body weight and appetite changes in patients with psychosis. <i>Psychopharmacology</i> , 2021, 238, 2449-2457.	3.1	15
41	Cortical Pathways or Mechanism in the Face Inversion Effect in Patients with First-Episode Schizophrenia. <i>Neuropsychiatric Disease and Treatment</i> , 2021, Volume 17, 1893-1906.	2.2	1
42	Effect of citalopram on hippocampal volume in first-episode schizophrenia: Structural MRI results from the DECIFER trial. <i>Psychiatry Research - Neuroimaging</i> , 2021, 312, 111286.	1.8	3
43	Increased subcortical region volume induced by electroconvulsive therapy in patients with schizophrenia. <i>European Archives of Psychiatry and Clinical Neuroscience</i> , 2021, 271, 1285-1295.	3.2	3
44	Developments in Biological Mechanisms and Treatments for Negative Symptoms and Cognitive Dysfunction of Schizophrenia. <i>Neuroscience Bulletin</i> , 2021, 37, 1609-1624.	2.9	28
45	Disrupted intrinsic functional brain topology in patients with major depressive disorder. <i>Molecular Psychiatry</i> , 2021, 26, 7363-7371.	7.9	82
46	Increased Homotopic Connectivity in the Prefrontal Cortex Modulated by Olanzapine Predicts Therapeutic Efficacy in Patients with Schizophrenia. <i>Neural Plasticity</i> , 2021, 2021, 1-11.	2.2	8
47	Gene expression of methylation cycle and related genes in lymphocytes and brain of patients with schizophrenia and non-psychotic controls. <i>Biomarkers in Neuropsychiatry</i> , 2021, 5, 100038.	1.0	7
48	A Selective Review of the Excitatory-Inhibitory Imbalance in Schizophrenia: Underlying Biology, Genetics, Microcircuits, and Symptoms. <i>Frontiers in Cell and Developmental Biology</i> , 2021, 9, 664535.	3.7	30
49	Abnormal interhemispheric homotopic functional connectivity is correlated with gastrointestinal symptoms in patients with major depressive disorder. <i>Journal of Psychiatric Research</i> , 2021, 144, 234-240.	3.1	8
50	Shared and Distinct Fractional Amplitude of Low-Frequency Fluctuation Patterns in Major Depressive Disorders With and Without Gastrointestinal Symptoms. <i>Frontiers in Psychiatry</i> , 2021, 12, 744898.	2.6	6
51	Differentiating Melancholic and Non-melancholic Major Depressive Disorder Using Fractional Amplitude of Low-Frequency Fluctuations. <i>Frontiers in Psychiatry</i> , 2021, 12, 763770.	2.6	4
52	Altered Brain Functional Asymmetry in Patients With Major Depressive Disorder Related to Gastrointestinal Symptoms. <i>Frontiers in Neuroscience</i> , 2021, 15, 797598.	2.8	3
53	Resting-state functional hypoconnectivity of amygdala in clinical high risk state and first-episode schizophrenia. <i>Brain Imaging and Behavior</i> , 2020, 14, 1840-1849.	2.1	9
54	Reduced connectivity in anterior cingulate cortex as an early predictor for treatment response in drug-naive, first-episode schizophrenia: A global-brain functional connectivity analysis. <i>Schizophrenia Research</i> , 2020, 215, 337-343.	2.0	24

#	ARTICLE	IF	CITATIONS
55	Phenotype-to-genotype approach reveals head circumference-associated genes in an autism spectrum disorder cohort. <i>Clinical Genetics</i> , 2020, 97, 338-346.	2.0	29
56	Region-specific insular volumetric decreases in drug-naïve, first-episode schizophrenia and their unaffected siblings. <i>American Journal of Medical Genetics Part B: Neuropsychiatric Genetics</i> , 2020, 183, 106-112.	1.7	7
57	Genome-wide association analysis of autism identified multiple loci that have been reported as strong signals for neuropsychiatric disorders. <i>Autism Research</i> , 2020, 13, 382-396.	3.8	16
58	Altered resting-state dynamic functional brain networks in major depressive disorder: Findings from the REST-meta-MDD consortium. <i>NeuroImage: Clinical</i> , 2020, 26, 102163.	2.7	76
59	Abnormal default-mode network homogeneity and its correlations with neurocognitive deficits in drug-naïve first-episode adolescent-onset schizophrenia. <i>Schizophrenia Research</i> , 2020, 215, 140-147.	2.0	17
60	Enhanced Connectivity of Thalamo-Cortical Networks in First-Episode, Treatment-Naïve Somatization Disorder. <i>Frontiers in Psychiatry</i> , 2020, 11, 555836.	2.6	4
61	Public behavior change, perceptions, depression, and anxiety in relation to the COVID-19 outbreak. <i>Open Forum Infectious Diseases</i> , 2020, 7, ofaa273.	0.9	24
62	Biotypes of major depressive disorder: Neuroimaging evidence from resting-state default mode network patterns. <i>NeuroImage: Clinical</i> , 2020, 28, 102514.	2.7	51
63	Chinese Taoist Cognitive Therapy for Symptoms of Depression and Anxiety in Adults in China: A Systematic Review and Meta-Analysis. <i>Frontiers in Psychology</i> , 2020, 11, 769.	2.1	5
64	Increased Appetite Plays a Key Role in Olanzapine-Induced Weight Gain in First-Episode Schizophrenia Patients. <i>Frontiers in Pharmacology</i> , 2020, 11, 739.	3.5	25
65	Frontal white matter abnormalities reveal the pathological basis underlying negative symptoms in antipsychotic-naïve, first-episode patients with adolescent-onset schizophrenia: Evidence from multimodal brain imaging. <i>Schizophrenia Research</i> , 2020, 222, 258-266.	2.0	7
66	Which Subgroup of First-Episode Schizophrenia Patients Can Remit During the First Year of Antipsychotic Treatment?. <i>Frontiers in Psychiatry</i> , 2020, 11, 566.	2.6	3
67	Genotype and Phenotype Correlations for TBL1XR1 in Neurodevelopmental Disorders. <i>Journal of Molecular Neuroscience</i> , 2020, 70, 2085-2092.	2.3	9
68	Air pollution and hippocampal atrophy in first episode schizophrenia. <i>Schizophrenia Research</i> , 2020, 218, 63-69.	2.0	7
69	Hippocampal functional network: The mediating role between obsession and anxiety in adult patients with obsessive-compulsive disorder. <i>World Journal of Biological Psychiatry</i> , 2020, 21, 685-695.	2.6	9
70	Metformin attenuates antipsychotic-induced metabolic dysfunctions in MK801-induced schizophrenia-like rats. <i>Psychopharmacology</i> , 2020, 237, 2257-2277.	3.1	8
71	Care for the Psychological Status of Frontline Medical Staff Fighting Against Coronavirus Disease 2019 (COVID-19). <i>Clinical Infectious Diseases</i> , 2020, 71, 3268-3269.	5.8	89
72	Metacognitive Training Modulates Default-Mode Network Homogeneity During 8-Week Olanzapine Treatment in Patients With Schizophrenia. <i>Frontiers in Psychiatry</i> , 2020, 11, 234.	2.6	9

#	ARTICLE	IF	CITATIONS
73	Effects of Sleep Disturbances on Behavioral Problems in Preschool Children With Autism Spectrum Disorder. <i>Frontiers in Psychiatry</i> , 2020, 11, 559694.	2.6	7
74	The Association Between Metabolic Disturbance and Cognitive Impairments in Early-Stage Schizophrenia. <i>Frontiers in Human Neuroscience</i> , 2020, 14, 599720.	2.0	4
75	Research advances in add-on treatment for negative symptoms and cognitive dysfunction in schizophrenia. <i>Journal of Central South University (Medical Sciences)</i> , 2020, 45, 1457-1463.	0.1	0
76	Dysregulated Maturation of the Functional Connectome in Antipsychotic-Naïve, First-Episode Patients With Adolescent-Onset Schizophrenia. <i>Schizophrenia Bulletin</i> , 2019, 45, 689-697.	4.3	30
77	Effects of Aspirin in Rats With Ouabain Intracerebral Treatment—Possible Involvement of Inflammatory Modulation?. <i>Frontiers in Psychiatry</i> , 2019, 10, 497.	2.6	6
78	Enhanced baseline activity in the left ventromedial putamen predicts individual treatment response in drug-naive, first-episode schizophrenia: Results from two independent study samples. <i>EBioMedicine</i> , 2019, 46, 248-255.	6.1	24
79	POGZ de novo missense variants in neuropsychiatric disorders. <i>Molecular Genetics & Genomic Medicine</i> , 2019, 7, e900.	1.2	13
80	Gender differences of neurocognitive functioning in patients with first-episode schizophrenia in China. <i>Comprehensive Psychiatry</i> , 2019, 95, 152132.	3.1	15
81	Reduced Brain Activity in the Right Putamen as an Early Predictor for Treatment Response in Drug-Naive, First-Episode Schizophrenia. <i>Frontiers in Psychiatry</i> , 2019, 10, 741.	2.6	20
82	The effect of minocycline on amelioration of cognitive deficits and pro-inflammatory cytokines levels in patients with schizophrenia. <i>Schizophrenia Research</i> , 2019, 212, 92-98.	2.0	57
83	An open-label randomised comparison of aripiprazole, olanzapine and risperidone for the acute treatment of first-episode schizophrenia: Eight-week outcomes. <i>Journal of Psychopharmacology</i> , 2019, 33, 1227-1236.	4.0	24
84	Increased Nucleus Accumbens Connectivity in Resting-State Patients With Drug-Naive, First-Episode Somatization Disorder. <i>Frontiers in Psychiatry</i> , 2019, 10, 585.	2.6	9
85	Disruptive variants of <i>CSDE1</i> associate with autism and interfere with neuronal development and synaptic transmission. <i>Science Advances</i> , 2019, 5, eaax2166.	10.3	35
86	Citalopram in first episode schizophrenia: The DECIFER trial. <i>Schizophrenia Research</i> , 2019, 208, 331-337.	2.0	15
87	Brain-derived neurotrophic factor is associated with cognitive impairments in first-episode and chronic schizophrenia. <i>Psychiatry Research</i> , 2019, 273, 528-536.	3.3	47
88	Cognitive deficits in subjects at risk for psychosis, first-episode and chronic schizophrenia patients. <i>Psychiatry Research</i> , 2019, 274, 235-242.	3.3	20
89	Voxel-based global-brain functional connectivity alterations in first-episode drug-naive patients with somatization disorder. <i>Journal of Affective Disorders</i> , 2019, 254, 82-89.	4.1	20
90	Dysfunction in Serotonergic and Noradrenergic Systems and Somatic Symptoms in Psychiatric Disorders. <i>Frontiers in Psychiatry</i> , 2019, 10, 286.	2.6	43

#	ARTICLE	IF	CITATIONS
91	Increased frontal gray matter volume in individuals with prodromal psychosis. <i>CNS Neuroscience and Therapeutics</i> , 2019, 25, 987-994.	3.9	16
92	Brain structural abnormalities as potential markers for detecting individuals with ultra-high risk for psychosis: A systematic review and meta-analysis. <i>Schizophrenia Research</i> , 2019, 209, 22-31.	2.0	49
93	Rare inherited missense variants of POGZ associate with autism risk and disrupt neuronal development. <i>Journal of Genetics and Genomics</i> , 2019, 46, 247-257.	3.9	17
94	Enhanced Global-Brain Functional Connectivity in the Left Superior Frontal Gyrus as a Possible Endophenotype for Schizophrenia. <i>Frontiers in Neuroscience</i> , 2019, 13, 145.	2.8	29
95	Reduced default mode network functional connectivity in patients with recurrent major depressive disorder. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2019, 116, 9078-9083.	7.1	441
96	<p>Pleiotropic action of genetic variation in ZNF804A on brain structure: a meta-analysis of magnetic resonance imaging studies</p>. <i>Neuropsychiatric Disease and Treatment</i> , 2019, Volume 15, 721-729.	2.2	6
97	Functional asymmetry of thalamocortical networks in subjects at ultra-high risk for psychosis and first-episode schizophrenia. <i>European Neuropsychopharmacology</i> , 2019, 29, 519-528.	0.7	24
98	Rates and predictors of one-year antipsychotic treatment discontinuation in first-episode schizophrenia: Results from an open-label, randomized, â€œreal worldâ€œ clinical trial. <i>Psychiatry Research</i> , 2019, 273, 631-640.	3.3	11
99	Cost-effectiveness of olanzapine in the first-line treatment of schizophrenia in China. <i>Journal of Medical Economics</i> , 2019, 22, 439-446.	2.1	12
100	Altered dynamic global signal topography in antipsychotic-naïve adolescents with early-onset schizophrenia. <i>Schizophrenia Research</i> , 2019, 208, 308-316.	2.0	25
101	Prevalence, clinical correlates and IQ of suicidal ideation in drug naïve Chinese Han patients with major depressive disorder. <i>Journal of Affective Disorders</i> , 2019, 248, 59-64.	4.1	13
102	Cerebellar structural and functional abnormalities in first-episode and drug-naïve patients with schizophrenia: A meta-analysis. <i>Psychiatry Research - Neuroimaging</i> , 2019, 283, 24-33.	1.8	46
103	An Overview of Genetic and Environmental Risk of Autism Spectrum Disorder. <i>Global Clinical and Translational Research</i> , 2019, , 37-44.	0.3	5
104	Prenatal Environment and Perinatal Factors Associated with Autism Spectrum Disorder. <i>Global Clinical and Translational Research</i> , 2019, , 100-108.	0.3	3
105	Minocycline adjunctive treatment to risperidone for negative symptoms in schizophrenia: Association with pro-inflammatory cytokine levels. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2018, 85, 69-76.	4.8	46
106	Cerebellar abnormalities in first-episode, drug-naïve schizophrenia at rest. <i>Psychiatry Research - Neuroimaging</i> , 2018, 276, 73-79.	1.8	29
107	Increased coherence-based regional homogeneity in resting-state patients with first-episode, drug-naïve somatization disorder. <i>Journal of Affective Disorders</i> , 2018, 235, 150-154.	4.1	12
108	Voxelâ€wise brainâ€wide functional connectivity abnormalities in firstâ€episode, drugâ€naïve patients with major depressive disorder. <i>American Journal of Medical Genetics Part B: Neuropsychiatric Genetics</i> , 2018, 177, 447-453.	1.7	25

#	ARTICLE	IF	CITATIONS
109	Altered functional connectivity strength and its correlations with cognitive function in subjects with ultra-high risk for psychosis at rest. <i>CNS Neuroscience and Therapeutics</i> , 2018, 24, 1140-1148.	3.9	21
110	Decreased Resting-State Interhemispheric Functional Connectivity Correlated with Neurocognitive Deficits in Drug-Naive First-Episode Adolescent-Onset Schizophrenia. <i>International Journal of Neuropsychopharmacology</i> , 2018, 21, 33-41.	2.1	43
111	Changes in plasma levels of nitric oxide metabolites and negative symptoms after 16-week minocycline treatment in patients with schizophrenia. <i>Schizophrenia Research</i> , 2018, 199, 390-394.	2.0	10
112	Decreased white matter FA values in the left inferior frontal gyrus is a possible intermediate phenotype of schizophrenia: evidences from a novel group strategy. <i>European Archives of Psychiatry and Clinical Neuroscience</i> , 2018, 268, 89-98.	3.2	10
113	Abnormal neural activity as a potential biomarker for drug-naive first-episode adolescent-onset schizophrenia with coherence regional homogeneity and support vector machine analyses. <i>Schizophrenia Research</i> , 2018, 192, 408-415.	2.0	52
114	Abnormal regional homogeneity as a potential imaging biomarker for adolescent-onset schizophrenia: A resting-state fMRI study and support vector machine analysis. <i>Schizophrenia Research</i> , 2018, 192, 179-184.	2.0	80
115	Treatment effects of olanzapine on homotopic connectivity in drug-free schizophrenia at rest. <i>World Journal of Biological Psychiatry</i> , 2018, 19, S106-S114.	2.6	31
116	Increased anterior default-mode network homogeneity in first-episode, drug-naive major depressive disorder: A replication study. <i>Journal of Affective Disorders</i> , 2018, 225, 767-772.	4.1	19
117	Aberrant corticostriatal connectivity predict positive symptoms of antipsychotic-naïve patients with adolescent-onset schizophrenia during brain maturation. <i>Schizophrenia Research</i> , 2018, 195, 564-566.	2.0	2
118	Decreased interhemispheric coordination in the posterior default-mode network and visual regions as trait alterations in first-episode, drug-naive major depressive disorder. <i>Brain Imaging and Behavior</i> , 2018, 12, 1251-1258.	2.1	23
119	Abnormal long- and short-range functional connectivity in adolescent-onset schizophrenia patients: A resting-state fMRI study. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2018, 81, 445-451.	4.8	55
120	Maternal dietary patterns, supplements intake and autism spectrum disorders. <i>Medicine (United States)</i> , 2018, 97, 1010-1017.	1.0	22
121	Inherited and multiple de novo mutations in autism/developmental delay risk genes suggest a multifactorial model. <i>Molecular Autism</i> , 2018, 9, 64.	4.9	114
122	Emotional Roles of Mono-Aminergic Neurotransmitters in Major Depressive Disorder and Anxiety Disorders. <i>Frontiers in Psychology</i> , 2018, 9, 2201.	2.1	126
123	Disrupted asymmetry of inter- and intra-hemispheric functional connectivity in patients with drug-naive, first-episode schizophrenia and their unaffected siblings. <i>EBioMedicine</i> , 2018, 36, 429-435.	6.1	32
124	Regional white matter volume abnormalities in first-episode somatization disorder. <i>International Journal of Psychophysiology</i> , 2018, 133, 12-16.	1.0	6
125	Social impairment of children with autism spectrum disorder affects parental quality of life in different ways. <i>Psychiatry Research</i> , 2018, 266, 168-174.	3.3	37
126	The Wnt Signaling Pathway Effector TCF7L2 Mediates Olanzapine-Induced Weight Gain and Insulin Resistance. <i>Frontiers in Pharmacology</i> , 2018, 9, 379.	3.5	23

#	ARTICLE	IF	CITATIONS
127	Bidirectional Causal Connectivity in the Cortico-Limbic-Cerebellar Circuit Related to Structural Alterations in First-Episode, Drug-Naive Somatization Disorder. <i>Frontiers in Psychiatry</i> , 2018, 9, 162.	2.6	16
128	Altered Serum Tumor Necrosis Factor and Interleukin-1 β in First-Episode Drug-Naive and Chronic Schizophrenia. <i>Frontiers in Neuroscience</i> , 2018, 12, 296.	2.8	28
129	Associations among maternal pre-pregnancy body mass index, gestational weight gain and risk of autism in the Han Chinese population. <i>BMC Psychiatry</i> , 2018, 18, 11.	2.6	23
130	A comparative study of magnetic resonance imaging on the gray matter and resting-state function in prodromal and first-episode schizophrenia. <i>American Journal of Medical Genetics Part B: Neuropsychiatric Genetics</i> , 2018, 177, 537-545.	1.7	9
131	Efficacy, acceptability and tolerability of 8 atypical antipsychotics in Chinese patients with acute schizophrenia: A network meta-analysis. <i>Schizophrenia Research</i> , 2017, 185, 73-79.	2.0	26
132	Aggressive behaviors and treatable risk factors of preschool children with autism spectrum disorder. <i>Autism Research</i> , 2017, 10, 1155-1162.	3.8	14
133	Genome-wide copy number variation analysis in a Chinese autism spectrum disorder cohort. <i>Scientific Reports</i> , 2017, 7, 44155.	3.3	50
134	Abnormal functional connectivity strength in patients with adolescent-onset schizophrenia: a resting-state fMRI study. <i>European Child and Adolescent Psychiatry</i> , 2017, 26, 839-845.	4.7	25
135	Anatomical distance affects cortical-subcortical connectivity in first-episode, drug-naive somatization disorder. <i>Journal of Affective Disorders</i> , 2017, 217, 153-158.	4.1	11
136	Using short-range and long-range functional connectivity to identify schizophrenia with a family-based case-control design. <i>Psychiatry Research - Neuroimaging</i> , 2017, 264, 60-67.	1.8	18
137	Olanzapine modulates the default-mode network homogeneity in recurrent drug-free schizophrenia at rest. <i>Australian and New Zealand Journal of Psychiatry</i> , 2017, 51, 1000-1009.	2.3	21
138	Family-based case-control study of homotopic connectivity in first-episode, drug-naive schizophrenia at rest. <i>Scientific Reports</i> , 2017, 7, 43312.	3.3	22
139	Predictors of menstruation restoration during metformin administration for treatment of antipsychotic drug-induced amenorrhea: A post hoc analysis. <i>Schizophrenia Research</i> , 2017, 190, 121-122.	2.0	3
140	Forkhead box C1 is targeted by microRNA-133b and promotes cell proliferation and migration in osteosarcoma. <i>Experimental and Therapeutic Medicine</i> , 2017, 14, 2823-2830.	1.8	10
141	Shared atypical default mode and salience network functional connectivity between autism and schizophrenia. <i>Autism Research</i> , 2017, 10, 1776-1786.	3.8	60
142	Frequency-specific alteration of functional connectivity density in antipsychotic-naive adolescents with early-onset schizophrenia. <i>Journal of Psychiatric Research</i> , 2017, 95, 68-75.	3.1	37
143	Olanzapine modulation of long- and short-range functional connectivity in the resting brain in a sample of patients with schizophrenia. <i>European Neuropsychopharmacology</i> , 2017, 27, 48-58.	0.7	37
144	Safety and efficacy of paliperidone palmitate 1-month formulation in Chinese patients with schizophrenia: a 25-week, open-label, multicenter, Phase IV study. <i>Neuropsychiatric Disease and Treatment</i> , 2017, Volume 13, 2045-2056.	2.2	10

#	ARTICLE	IF	CITATIONS
145	Tranlycypromine Causes Neurotoxicity and Represses BHC110/LSD1 in Human-Induced Pluripotent Stem Cell-Derived Cerebral Organoids Model. <i>Frontiers in Neurology</i> , 2017, 8, 626.	2.4	27
146	Relationship between long-term use of a typical antipsychotic medication by Chinese schizophrenia patients and the bone turnover markers serum osteocalcin and β -CrossLaps. <i>Schizophrenia Research</i> , 2016, 176, 259-263.	2.0	21
147	Abnormal regional homogeneity as potential imaging biomarker for psychosis risk syndrome: a resting-state fMRI study and support vector machine analysis. <i>Scientific Reports</i> , 2016, 6, 27619.	3.3	25
148	Patients with first-episode, drug-naive schizophrenia and subjects at ultra-high risk of psychosis shared increased cerebellar-default mode network connectivity at rest. <i>Scientific Reports</i> , 2016, 6, 26124.	3.3	46
149	Identification of RELN variation p.Thr3192Ser in a Chinese family with schizophrenia. <i>Scientific Reports</i> , 2016, 6, 24327.	3.3	27
150	Altered plasma levels of chemokines in autism and their association with social behaviors. <i>Psychiatry Research</i> , 2016, 244, 300-305.	3.3	35
151	Frequency-specific alterations in functional connectivity in treatment-resistant and -sensitive major depressive disorder. <i>Journal of Psychiatric Research</i> , 2016, 82, 30-39.	3.1	34
152	De novo genic mutations among a Chinese autism spectrum disorder cohort. <i>Nature Communications</i> , 2016, 7, 13316.	12.8	293
153	Insertion of a knockout-first cassette in <i>Ampd1</i> gene leads to neonatal death by disruption of neighboring genes expression. <i>Scientific Reports</i> , 2016, 6, 35970.	3.3	19
154	Differences among first-episode schizophrenia patients, healthy siblings, and controls at the individual level. <i>International Journal of Psychophysiology</i> , 2016, 104, 24-32.	1.0	3
155	A novel de novo POGZ mutation in a patient with intellectual disability. <i>Journal of Human Genetics</i> , 2016, 61, 357-359.	2.3	31
156	Decreased interhemispheric functional connectivity in insula and angular gyrus/supramarginal gyrus: Significant findings in first-episode, drug-naive somatization disorder. <i>Psychiatry Research - Neuroimaging</i> , 2016, 248, 48-54.	1.8	37
157	Abnormal default-mode network homogeneity and its correlations with personality in drug-naive somatization disorder at rest. <i>Journal of Affective Disorders</i> , 2016, 193, 81-88.	4.1	31
158	Disrupted amplitude of low-frequency fluctuations in antipsychotic-naïve adolescents with early-onset schizophrenia. <i>Psychiatry Research - Neuroimaging</i> , 2016, 249, 20-26.	1.8	24
159	Decreased long- and short-range functional connectivity at rest in drug-naive major depressive disorder. <i>Australian and New Zealand Journal of Psychiatry</i> , 2016, 50, 763-769.	2.3	21
160	Association Between Maternal Obesity and Autism Spectrum Disorder in Offspring: A Meta-analysis. <i>Journal of Autism and Developmental Disorders</i> , 2016, 46, 95-102.	2.7	73
161	Resting-state cerebellar-cerebral networks are differently affected in first-episode, drug-naive schizophrenia patients and unaffected siblings. <i>Scientific Reports</i> , 2015, 5, 17275.	3.3	45
162	Association of genetic variants of GRIN2B with autism. <i>Scientific Reports</i> , 2015, 5, 8296.	3.3	39

#	ARTICLE	IF	CITATIONS
163	ZNF804A rs1344706 is associated with cortical thickness, surface area, and cortical volume of the unmedicated first episode schizophrenia and healthy controls. <i>American Journal of Medical Genetics Part B: Neuropsychiatric Genetics</i> , 2015, 168, 265-273.	1.7	24
164	Case-control association study of ABCB1 gene and major depressive disorder in a local Chinese Han population. <i>Neuropsychiatric Disease and Treatment</i> , 2015, 11, 1967.	2.2	13
165	Paliperidone extended-release tablets in Chinese patients with schizophrenia: meta-analysis of randomized controlled trials. <i>Neuropsychiatric Disease and Treatment</i> , 2015, 11, 1817.	2.2	8
166	Increased Cerebellar Functional Connectivity With the Default-Mode Network in Unaffected Siblings of Schizophrenia Patients at Rest. <i>Schizophrenia Bulletin</i> , 2015, 41, 1317-1325.	4.3	48
167	Dissociation of anatomical and functional alterations of the default-mode network in first-episode, drug-naive schizophrenia. <i>Clinical Neurophysiology</i> , 2015, 126, 2276-2281.	1.5	10
168	Unidirectionally affected causal connectivity of cortico-limbic-cerebellar circuit by structural deficits in drug-naive major depressive disorder. <i>Journal of Affective Disorders</i> , 2015, 172, 410-416.	4.1	31
169	Guidelines concordance of maintenance treatment in euthymic patients with bipolar disorder: Data from the national bipolar mania pathway survey (BIPAS) in mainland China. <i>Journal of Affective Disorders</i> , 2015, 182, 101-105.	4.1	6
170	Dissociation of functional and anatomical brain abnormalities in unaffected siblings of schizophrenia patients. <i>Clinical Neurophysiology</i> , 2015, 126, 927-932.	1.5	28
171	Decreased insular connectivity in drug-naive major depressive disorder at rest. <i>Journal of Affective Disorders</i> , 2015, 179, 31-37.	4.1	65
172	Employment and financial burden of families with preschool children diagnosed with autism spectrum disorders in urban China: results from a descriptive study. <i>BMC Psychiatry</i> , 2015, 15, 3.	2.6	59
173	Regional white matter abnormalities in drug-naive, first-episode schizophrenia patients and their healthy unaffected siblings. <i>Australian and New Zealand Journal of Psychiatry</i> , 2015, 49, 246-254.	2.3	25
174	Decreased regional activity and network homogeneity of the fronto-limbic network at rest in drug-naive major depressive disorder. <i>Australian and New Zealand Journal of Psychiatry</i> , 2015, 49, 550-556.	2.3	34
175	Dysfunctional resting-state connectivities of brain regions with structural deficits in drug-naive first-episode schizophrenia adolescents. <i>Schizophrenia Research</i> , 2015, 168, 353-359.	2.0	31
176	Increased short-range and long-range functional connectivity in first-episode, medication-naive schizophrenia at rest. <i>Schizophrenia Research</i> , 2015, 166, 144-150.	2.0	48
177	Broad autism phenotype features of Chinese parents with autistic children and their associations with severity of social impairment in probands. <i>BMC Psychiatry</i> , 2015, 15, 168.	2.6	13
178	Abnormal Causal Connectivity by Structural Deficits in First-Episode, Drug-Naive Schizophrenia at Rest. <i>Schizophrenia Bulletin</i> , 2015, 41, 57-65.	4.3	71
179	Disrupted structural connectivity network in treatment-naive depression. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2015, 56, 18-26.	4.8	58
180	Different Visual Preference Patterns in Response to Simple and Complex Dynamic Social Stimuli in Preschool-Aged Children with Autism Spectrum Disorders. <i>PLoS ONE</i> , 2015, 10, e0122280.	2.5	50

#	ARTICLE	IF	CITATIONS
181	Effects of Clozapine and other Atypical Antipsychotics on Infants Development Who Were Exposed to as Fetus: A Post-Hoc Analysis. PLoS ONE, 2015, 10, e0123373.	2.5	30
182	Challenges in the early detection and intervention of the psychosis-risk syndrome. Shanghai Archives of Psychiatry, 2015, 27, 45-7.	0.7	0
183	Clinical utility of orally disintegrating olanzapine in Chinese patients with schizophrenia: a review of effectiveness, patient preference, adherence, and other properties. Neuropsychiatric Disease and Treatment, 2014, 10, 355.	2.2	5
184	APOA-I: A Possible Novel Biomarker for Metabolic Side Effects in First Episode Schizophrenia. PLoS ONE, 2014, 9, e93902.	2.5	14
185	Adjunctive treatment for cognitive impairment in patients with chronic schizophrenia: a double-blind, placebo-controlled study. Neuropsychiatric Disease and Treatment, 2014, 10, 1317.	2.2	10
186	Profile of minocycline and its potential in the treatment of schizophrenia. Neuropsychiatric Disease and Treatment, 2014, 10, 1103.	2.2	37
187	Mental health and burnout in primary and secondary school teachers in the remote mountain areas of Guangdong Province in the People's Republic of China. Neuropsychiatric Disease and Treatment, 2014, 10, 123.	2.2	24
188	The comparison of glucose and lipid metabolism parameters in drug-naïve, antipsychotic-treated, and antipsychotic discontinuation patients with schizophrenia. Neuropsychiatric Disease and Treatment, 2014, 10, 1361.	2.2	10
189	Minocycline supplementation for treatment of negative symptoms in early-phase schizophrenia: A double blind, randomized, controlled trial. Schizophrenia Research, 2014, 153, 169-176.	2.0	119
190	Neonatal intrahippocampal injection of lipopolysaccharide induces deficits in social behavior and prepulse inhibition and microglial activation in rats: Implication for a new schizophrenia animal model. Brain, Behavior, and Immunity, 2014, 38, 166-174.	4.1	51
191	Minocycline alleviates behavioral deficits and inhibits microglial activation in the offspring of pregnant mice after administration of polyriboinosinic-polyribocytidilic acid. Psychiatry Research, 2014, 219, 680-686.	3.3	87
192	Decreased gray matter volume in the left middle temporal gyrus as a candidate biomarker for schizophrenia: A study of drug naive, first-episode schizophrenia patients and unaffected siblings. Schizophrenia Research, 2014, 159, 43-50.	2.0	43
193	Hippocampal and orbital inferior frontal gray matter volume abnormalities and cognitive deficit in treatment-naïve, first-episode patients with schizophrenia. Schizophrenia Research, 2014, 152, 339-343.	2.0	68
194	Functional and anatomical brain deficits in drug-naïve major depressive disorder. Progress in Neuro-Psychopharmacology and Biological Psychiatry, 2014, 54, 1-6.	4.8	61
195	Minocycline and Risperidone Prevent Microglia Activation and Rescue Behavioral Deficits Induced by Neonatal Intrahippocampal Injection of Lipopolysaccharide in Rats. PLoS ONE, 2014, 9, e93966.	2.5	77
196	Guidelines Disconcordance in Acute Bipolar Depression: Data from the National Bipolar Mania Pathway Survey (BIPAS) in Mainland China. PLoS ONE, 2014, 9, e96096.	2.5	11
197	Shuganjieyu capsule increases neurotrophic factor expression in a rat model of depression. Neural Regeneration Research, 2014, 9, 489.	3.0	14
198	An investigation of economic costs of schizophrenia in two areas of China. International Journal of Mental Health Systems, 2013, 7, 26.	2.7	29

#	ARTICLE	IF	CITATIONS
199	Duration of Untreated Psychosis Is Associated with Temporal and Occipitotemporal Gray Matter Volume Decrease in Treatment Na ⁺ -ve Schizophrenia. PLoS ONE, 2013, 8, e83679.	2.5	44
200	Cognitive effects of atypical antipsychotic drugs in first-episode drug-na ⁺ -ve schizophrenic patients. Neural Regeneration Research, 2013, 8, 277-86.	3.0	20
201	Relationship between brain-derived neurotrophic factor gene C270T polymorphisms and the psychotic symptoms and cognitive functioning of patients with schizophrenia. Shanghai Archives of Psychiatry, 2012, 24, 328-34.	0.7	2
202	The Effects of Clozapine on Cognitive Function and Regional Cerebral Blood Flow in the Negative Symptom Profile Schizophrenia. International Journal of Psychiatry in Medicine, 2006, 36, 171-181.	1.8	21
203	Abnormal ACTH and prolactin responses to fenfluramine in rats exposed to single and multiple doses of MDMA. Psychopharmacology, 1997, 131, 411-419.	3.1	16