Pascal Mansuelle

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/4552171/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Lebetin Peptides, A New Class of Potent Platelet Aggregation Inhibitors: Chemical Synthesis, Biological Activity and NMR Spectroscopic Study. International Journal of Peptide Research and Therapeutics, 2020, 26, 21-31.	1.9	3
2	Characterization of all the lipolytic activities in pancreatin and comparison with porcine and human pancreatic juices. Biochimie, 2020, 169, 106-120.	2.6	23
3	The Ig-like domain of Punctin/MADD-4 is the primary determinant for interaction with the ectodomain of neuroligin NLG-1. Journal of Biological Chemistry, 2020, 295, 16267-16279.	3.4	11
4	lsolation of an Anti–tumour Disintegrin: Dabmaurin–1, a Peptide Lebein–1–like, from Daboia mauritanica Venom. Toxins, 2020, 12, 102.	3.4	7
5	The Hunt for the Closed Conformation of the Fruitâ€Ripening Enzyme 1â€Aminocyclopropaneâ€1â€carboxylic Oxidase: A Combined Electron Paramagnetic Resonance and Molecular Dynamics Study. Chemistry - A European Journal, 2019, 25, 13766-13776.	3.3	4
6	Identification of a new natural gastric lipase inhibitor from star anise. Food and Function, 2019, 10, 469-478.	4.6	17
7	Chemical Modification of 1-Aminocyclopropane Carboxylic Acid (ACC) Oxidase: Cysteine Mutational Analysis, Characterization, and Bioconjugation with a Nitroxide Spin Label. Molecular Biotechnology, 2019, 61, 650-662.	2.4	4
8	Functional characterization and FTIR-based 3D modeling of full length and truncated forms of Scorpio maurus venom phospholipase A 2. Biochimica Et Biophysica Acta - General Subjects, 2018, 1862, 1247-1261.	2.4	11
9	Electron transfer in an acidophilic bacterium: interaction between a diheme cytochrome and a cupredoxin. Chemical Science, 2018, 9, 4879-4891.	7.4	17
10	Glycosylate and move! The glycosyltransferase Maf is involved in bacterial flagella formation. Environmental Microbiology, 2018, 20, 228-240.	3.8	20
11	Characterization of pepsin from rabbit gastric extract, its action on β-casein and the effects of lipids on proteolysis. Food and Function, 2018, 9, 5975-5988.	4.6	11
12	A Low Molecular Weight Protein from the Sea Anemone Anemonia viridis with an Anti-Angiogenic Activity. Marine Drugs, 2018, 16, 134.	4.6	8
13	Two New Secreted Proteases Generate a Casein-Derived Antimicrobial Peptide in Bacillus cereus Food Born Isolate Leading to Bacterial Competition in Milk. Frontiers in Microbiology, 2018, 9, 1148.	3.5	29
14	Responses of the marine diatom Thalassiosira pseudonana to changes in CO2 concentration: a proteomic approach. Scientific Reports, 2017, 7, 42333.	3.3	36
15	Mechanisms involved in xyloglucan catabolism by the cellulosome-producing bacterium Ruminiclostridium cellulolyticum. Scientific Reports, 2016, 6, 22770.	3.3	62
16	The scorpion toxin Bot <scp>IX</scp> is a potent member of the αâ€like family and has a unique Nâ€terminal sequence extension. FEBS Letters, 2016, 590, 3221-3232.	2.8	5
17	Biochemical characterization of Yarrowia lipolytica LIP8, a secreted lipase with a cleavable C-terminal region. Biochimica Et Biophysica Acta - Molecular and Cell Biology of Lipids, 2015, 1851, 129-140.	2.4	6
18	Conformational modulation and hydrodynamic radii of <scp>CP</scp> 12 protein and its complexes probed by fluorescence correlation spectroscopy. FEBS Journal, 2014, 281, 3206-3217.	4.7	14

PASCAL MANSUELLE

#	Article	IF	CITATIONS
19	Glyceraldehydeâ€3â€phosphate dehydrogenase is regulated by ferredoxinâ€ <scp>NADP</scp> reductase in the diatom <i><scp>A</scp>sterionella formosa</i> . New Phytologist, 2014, 203, 414-423.	7.3	32
20	Lacticin LC14, a New Bacteriocin Produced by Lactococcus lactis BMG6.14: Isolation, Purification and Partial Characterization. Infectious Disorders - Drug Targets, 2012, 12, 316-325.	0.8	3
21	The mimivirus R355 gene product: preliminary crystallographic analysis of a putative ubiquitin-like protein-specific protease. Acta Crystallographica Section F: Structural Biology Communications, 2011, 67, 169-172.	0.7	Ο
22	Preliminary crystallographic analysis of a possible transcription factor encoded by the mimivirus L544 gene. Acta Crystallographica Section F: Structural Biology Communications, 2011, 67, 922-925.	0.7	1
23	Chemical Synthesis and Characterization of J46 Peptide, an Atypical Class IIa Bacteriocin from Lactococcus lactis subsp. cremoris J46 Strain. Journal of Antibiotics, 2008, 61, 89-93.	2.0	8
24	Nanobacteria Are Mineralo Fetuin Complexes. PLoS Pathogens, 2008, 4, e41.	4.7	88
25	Phoneutria nigriventer Toxin 1: A Novel, State-Dependent Inhibitor of Neuronal Sodium Channels That Interacts with 1¼ Conotoxin Binding Sites. Molecular Pharmacology, 2006, 69, 1931-1937.	2.3	47
26	Chromatographic Characterization and Phytotoxic Activity of Fusarium oxysporum f. sp. albedinis and Saprophytic Strain Toxins. Journal of Phytopathology, 2005, 153, 203-208.	1.0	7
27	Increasing the molecular contacts between maurotoxin and Kv1.2 channel augments ligand affinity. Proteins: Structure, Function and Bioinformatics, 2005, 60, 401-411.	2.6	10
28	The impact of the fourth disulfide bridge in scorpion toxins of the α-KTx6 subfamily. Proteins: Structure, Function and Bioinformatics, 2005, 61, 1010-1023.	2.6	21
29	First chemical synthesis of a scorpion α-toxin affecting sodium channels: The Aah I toxin ofAndroctonus australis hector. Journal of Peptide Science, 2004, 10, 666-677.	1.4	12
30	Synthesis and characterization of Pi4, a scorpion toxin from Pandinus imperator that acts on K+ channels. FEBS Journal, 2003, 270, 3583-3592.	0.2	41
31	Covalent structure and some pharmacological features of native and cleaved ?-KTx12?1, a four disulfide-bridged toxin fromTityus serrulatus venom. Journal of Peptide Science, 2003, 9, 132-140.	1.4	17
32	PnTx4-3, a new insect toxin from Phoneutria nigriventer venom elicits the glutamate uptake inhibition exhibited by PhTx4 toxic fraction. Toxicon, 2003, 42, 793-800.	1.6	27
33	Novel structural class of four disulfide-bridged peptides from Tityus serrulatus venom. Biochemical and Biophysical Research Communications, 2003, 301, 1086-1092.	2.1	30
34	Phoneutria nigriventer ω-Phonetoxin IIA Blocks the Cav2 Family of Calcium Channels and Interacts with ω-Conotoxin-binding Sites. Journal of Biological Chemistry, 2002, 277, 13856-13862.	3.4	57
35	Chemical Synthesis, Molecular Modeling, and Antimicrobial Activity of a Novel Bacteriocin, MMFII. Biochemical and Biophysical Research Communications, 2001, 289, 13-18.	2.1	8
36	Disulfide bridge reorganization induced by proline mutations in maurotoxin. FEBS Letters, 2001, 489, 202-207.	2.8	19

PASCAL MANSUELLE

#	Article	IF	CITATIONS
37	Parameters affecting in vitro oxidation/folding of maurotoxin, a four-disulphide-bridged scorpion toxin. Biochemical Journal, 2001, 358, 681-692.	3.7	21
38	Parameters affecting in vitro oxidation/folding of maurotoxin, a four-disulphide-bridged scorpion toxin. Biochemical Journal, 2001, 358, 681.	3.7	14
39	Purification, characterization and molecular modelling of two toxin-like proteins from the Androctonus australis Hector venom. FEBS Journal, 2000, 267, 5614-5620.	0.2	22
40	Maurotoxin Versus Pi1/HsTx1 Scorpion Toxins. Journal of Biological Chemistry, 2000, 275, 39394-39402.	3.4	38
41	KTX3, the kaliotoxin from Buthus occitanus tunetanus scorpion venom: one of an extensive family of peptidyl ligands of potassium channels. Toxicon, 2000, 38, 105-111.	1.6	13
42	A Novel Anti-Ep-CAM Antibody to Analyze the Organization of Thymic Medulla in Autoimmunity. Current Topics in Microbiology and Immunology, 2000, 251, 109-117.	1.1	4
43	Scorpion α-like toxins, toxic to both mammals and insects, differentially interact with receptor site 3 on voltage-gated sodium channels in mammals and insects. European Journal of Neuroscience, 1999, 11, 975-985.	2.6	46
44	Role of lysine and tryptophan residues in the biological activity of toxin VII (Ts gamma) from the scorpion Tityus serrulatus. FEBS Journal, 1999, 260, 76-86.	0.2	36
45	Aah VI, a novel, N -glycosylated anti-insect toxin from Androctonus australis hector scorpion venom: isolation, characterisation, and glycan structure determination. FEBS Letters, 1999, 443, 175-180.	2.8	30
46	A novel immunoglobulin superfamily junctional molecule expressed by antigen presenting cells, endothelial cells and platelets. Molecular Immunology, 1998, 35, 1111-1119.	2.2	90
47	Maurotoxin, a four disulfide bridges scorpion toxin acting on K+ channels. Toxicon, 1998, 36, 1609-1611.	1.6	18
48	Purification, structure and activity of three insect toxins from Buthus occitanus tunetanus venom. Toxicon, 1997, 35, 365-382.	1.6	38
49	Maurotoxin, a four disulfide bridge toxin fromScorpio maurusvenom: purification, structure and action on potassium channels. FEBS Letters, 1997, 406, 284-290.	2.8	95
50	1H-NMR-Derived Secondary Structure and Overall Fold of a Natural Anatoxin from the Scorpion Androctonus Australis Hector. FEBS Journal, 1997, 247, 1118-1126.	0.2	23
51	Biochemical and Pharmacological Characterization of a Depressant Insect Toxin from the Venom of the Scorpion Buthacus arenicola. FEBS Journal, 1997, 243, 93-99.	0.2	38
52	Two neurotoxins (BmK I and BmK II) from the venom of the scorpion Buthus martensi Karsch: purification, amino acid sequences and assessment of specific activity. Toxicon, 1996, 34, 987-1001.	1.6	107
53	Novel anti-platelet aggregation polypeptides fromVipera lebetinavenom: Isolation and characterization. FEBS Letters, 1996, 392, 6-10.	2.8	40
54	A New Scorpion Venom Toxin Paralytic to Insects that Affects Na+ Channel Activation FEBS Journal, 1996, 241, 525-532.	0.2	21

PASCAL MANSUELLE

#	Article	IF	CITATIONS
55	Characterization of PO ₁ , a new peptide ligand of the apaminâ€sensitive Ca ²⁺ activated K ⁺ channel. International Journal of Peptide and Protein Research, 1996, 48, 514-521.	0.1	40
56	Characterization and ultrastructural localization of annexin VI from mitochondria. FEBS Letters, 1995, 360, 80-84.	2.8	20
57	Positively charged amino acid residues located similarly in sea anemone and scorpion toxins. Journal of Biological Chemistry, 1994, 269, 16785-16788.	3.4	59
58	The kaliotoxin family enlarged. Purification, characterization, and precursor nucleotide sequence of KTX2 from Androctonus australis venom. Journal of Biological Chemistry, 1994, 269, 32835-32843.	3.4	72
59	The kaliotoxin family enlarged. Purification, characterization, and precursor nucleotide sequence of KTX2 from Androctonus australis venom. Journal of Biological Chemistry, 1994, 269, 32835-43.	3.4	62
60	Positively charged amino acid residues located similarly in sea anemone and scorpion toxins. Journal of Biological Chemistry, 1994, 269, 16785-8.	3.4	52
61	Characterization of toxin III of the scorpionLeiurus quinquestriatus quinquestriatus: A new type of alpha-toxin highly toxic both to mammals and insects. Natural Toxins, 1993, 1, 308-312.	1.0	60
62	Characterization of a new leiurotoxin I-like scorpion toxin. FEBS Letters, 1993, 320, 189-192.	2.8	69
63	Fine molecular analysis of the antigenicity of the Androctonus australis hector scorpion neurotoxin II: A new antigenic epitope disclosed by the pepscan method. Molecular Immunology, 1993, 30, 1061-1068.	2.2	21
64	Synthesis and characterization of kaliotoxin. Is the 26-32 sequence essential for potassium channel recognition?. Journal of Biological Chemistry, 1993, 268, 26302-26309.	3.4	70
65	Synthesis and characterization of kaliotoxin. Is the 26-32 sequence essential for potassium channel recognition?. Journal of Biological Chemistry, 1993, 268, 26302-9.	3.4	56
66	Monoclonal antibodies to toxin II from the scorpion Androctonus australis Hector: Further characterization of epitope specificities and neutralizing capacities. Toxicon, 1992, 30, 723-731.	1.6	10
67	The amino acid sequence of toxin IV from theAndroctonus australis scorpion: Differing effects of natural mutations in scorpion α-toxins on their antigenic and toxic properties. Natural Toxins, 1992, 1, 61-69.	1.0	19
68	The β-type toxin Ts II from the scorpionTityus serrulatus: Amino acid sequence determination and assessment of biological and antigenic properties. Natural Toxins, 1992, 1, 119-125.	1.0	26
69	Kaliotoxin, a novel peptidyl inhibitor of neuronal BK-type Ca(2+)-activated K+ channels characterized from Androctonus mauretanicus mauretanicus venom Journal of Biological Chemistry, 1992, 267, 1640-1647.	3.4	175
70	Kaliotoxin, a novel peptidyl inhibitor of neuronal BK-type Ca(2+)-activated K+ channels characterized from Androctonus mauretanicus mauretanicus venom. Journal of Biological Chemistry, 1992, 267, 1640-7.	3.4	137
71	An anti-insect toxin purified from the scorpion Androctonus australis Hector also acts on the .alpha and .betasites of the mammalian sodium channel: sequence and circular dichroism study. Biochemistry, 1991, 30, 633-640.	2.5	94
72	Neurotoxins active on insects: amino acid sequences, chemical modifications, and secondary structure estimation by circular dichroism of toxins from the scorpion Androctonus australis Hector. Biochemistry, 1990, 29, 1492-1501.	2.5	59

#	Article	IF	CITATIONS
73	Delineation of the functional site of a snake venom cardiotoxin: preparation, structure, and function of monoacetylated derivatives. Biochemistry, 1990, 29, 6480-6489.	2.5	49
74	Primary structure of scorpion anti-insect toxins isolated from the venom of Leiurus quinquestriatus quinquestriatus. FEBS Letters, 1990, 261, 423-426.	2.8	66