

Bernardo L Trigatti

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4576754/publications.pdf>

Version: 2024-02-01

65
papers

5,213
citations

136950

32
h-index

106344

65
g-index

66
all docs

66
docs citations

66
times ranked

4411
citing authors

#	ARTICLE	IF	CITATIONS
1	Good Cholesterol Gone Bad? HDL and COVID-19. <i>International Journal of Molecular Sciences</i> , 2021, 22, 10182.	4.1	20
2	Salsalate reduces atherosclerosis through AMPK β 1 in mice. <i>Molecular Metabolism</i> , 2021, 53, 101321.	6.5	8
3	Pcpe2: A New Partner for the Scavenger Receptor Class B Type I in High-Density Lipoprotein Selective Lipid Uptake. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2021, 41, 2726-2729.	2.4	0
4	High Density Lipoprotein and Its Precursor Protein Apolipoprotein A1 as Potential Therapeutics to Prevent Anthracycline Associated Cardiotoxicity. <i>Frontiers in Cardiovascular Medicine</i> , 2020, 7, 65.	2.4	5
5	Un-JAMming atherosclerotic arteries: JAM-L as a target to attenuate plaque development. <i>Clinical Science</i> , 2019, 133, 1581-1585.	4.3	3
6	Treatment with apolipoprotein A1 protects mice against doxorubicin-induced cardiotoxicity in a scavenger receptor class B, type I-dependent manner. <i>American Journal of Physiology - Heart and Circulatory Physiology</i> , 2019, 316, H1447-H1457.	3.2	17
7	High-density lipoprotein protects cardiomyocytes against necrosis induced by oxygen and glucose deprivation through SR-B1, PI3K, and AKT1 and 2. <i>Biochemical Journal</i> , 2018, 475, 1253-1265.	3.7	26
8	HDL protects against doxorubicin-induced cardiotoxicity in a scavenger receptor class B type 1-, PI3K-, and Akt-dependent manner. <i>American Journal of Physiology - Heart and Circulatory Physiology</i> , 2018, 314, H31-H44.	3.2	18
9	Rosuvastatin Reduces Aortic Sinus and Coronary Artery Atherosclerosis in SR-B1 (Scavenger Receptor) Tj ETQq1 1 0.784314 rgBT /Over Lowering. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2018, 38, 26-39.	2.4	24
10	Hyperglycemia Aggravates Diet-Induced Coronary Artery Disease and Myocardial Infarction in SR-B1-Knockout/ApoE-Hypomorphic Mice. <i>Frontiers in Physiology</i> , 2018, 9, 1398.	2.8	12
11	Data on leukocyte PDZK1 deficiency affecting macrophage apoptosis but not monocyte recruitment, cell proliferation, macrophage abundance or ER stress in atherosclerotic plaques of LDLR deficient mice. <i>Data in Brief</i> , 2018, 19, 1148-1161.	1.0	3
12	PDZK1 in leukocytes protects against cellular apoptosis and necrotic core development in atherosclerotic plaques in high fat diet fed ldl receptor deficient mice. <i>Atherosclerosis</i> , 2018, 276, 171-181.	0.8	9
13	Low-density lipoprotein (LDL)-dependent uptake of Gram-positive lipoteichoic acid and Gram-negative lipopolysaccharide occurs through LDL receptor. <i>Scientific Reports</i> , 2018, 8, 10496.	3.3	47
14	Sialidase down-regulation reduces non-HDL cholesterol, inhibits leukocyte transmigration, and attenuates atherosclerosis in ApoE knockout mice. <i>Journal of Biological Chemistry</i> , 2018, 293, 14689-14706.	3.4	42
15	SR-B1 and PDZK1. <i>Current Opinion in Lipidology</i> , 2017, 28, 201-208.	2.7	30
16	Suppression of NK and CD8+ T cells reduces astrogliosis but accelerates cerebellar dysfunction and shortens life span in a mouse model of Sandhoff disease. <i>Journal of Neuroimmunology</i> , 2017, 306, 55-67.	2.3	4
17	Sphingosine-1-Phosphate Receptor 1, Expressed in Myeloid Cells, Slows Diet-Induced Atherosclerosis and Protects against Macrophage Apoptosis in Ldlr KO Mice. <i>International Journal of Molecular Sciences</i> , 2017, 18, 2721.	4.1	22
18	HDL signaling and protection against coronary artery atherosclerosis in mice. <i>Journal of Biomedical Research</i> , 2016, 30, 94-100.	1.6	16

#	ARTICLE	IF	CITATIONS
19	Rare Genetic Variants and High-Density Lipoprotein. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2016, 36, e53-5.	2.4	11
20	Characterization of Proliferating Lesion-Resident Cells During All Stages of Atherosclerotic Growth. <i>Journal of the American Heart Association</i> , 2016, 5, .	3.7	28
21	Characterization of mice harboring a variant of EPCR with impaired ability to bind protein C: novel role of EPCR in hematopoiesis. <i>Blood</i> , 2015, 126, 673-682.	1.4	24
22	The Effects of Diet on Occlusive Coronary Artery Atherosclerosis and Myocardial Infarction in Scavenger Receptor Class B, Type 1/Low-Density Lipoprotein Receptor Double Knockout Mice. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2014, 34, 2394-2403.	2.4	52
23	High Density Lipoprotein Stimulated Migration of Macrophages Depends on the Scavenger Receptor Class B, Type I, PDZK1 and Akt1 and Is Blocked by Sphingosine 1 Phosphate Receptor Antagonists. <i>PLoS ONE</i> , 2014, 9, e106487.	2.5	43
24	Interleukin-15 Modulates Adipose Tissue by Altering Mitochondrial Mass and Activity. <i>PLoS ONE</i> , 2014, 9, e114799.	2.5	31
25	The effect of pomegranate extract on coronary artery atherosclerosis in SR-BI/APOE double knockout mice. <i>Atherosclerosis</i> , 2013, 228, 80-89.	0.8	54
26	Deficiency of TDAG51 Protects Against Atherosclerosis by Modulating Apoptosis, Cholesterol Efflux, and Peroxiredoxin-1 Expression. <i>Journal of the American Heart Association</i> , 2013, 2, e000134.	3.7	27
27	Deletion of tumor necrosis factor- α ameliorates neurodegeneration in Sandhoff disease mice. <i>Human Molecular Genetics</i> , 2013, 22, 3960-3975.	2.9	43
28	SR-BI in Bone Marrow Derived Cells Protects Mice from Diet Induced Coronary Artery Atherosclerosis and Myocardial Infarction. <i>PLoS ONE</i> , 2013, 8, e72492.	2.5	24
29	Hypomorphic sialidase expression decreases serum cholesterol by downregulation of VLDL production in mice. <i>Journal of Lipid Research</i> , 2012, 53, 2573-2585.	4.2	37
30	Modifications in Perfringolysin O Domain 4 Alter the Cholesterol Concentration Threshold Required for Binding. <i>Biochemistry</i> , 2012, 51, 3373-3382.	2.5	82
31	Modulators of Protein Kinase C Affect SR-BI-Dependent HDL Lipid Uptake in Transfected HepG2 Cells. <i>Cholesterol</i> , 2011, 2011, 1-11.	1.6	10
32	Interleukin-15 Contributes to the Regulation of Murine Adipose Tissue and Human Adipocytes. <i>Obesity</i> , 2010, 18, 1601-1607.	3.0	95
33	A role for the scavenger receptor, class B type I in high density lipoprotein dependent activation of cellular signaling pathways. <i>Biochimica Et Biophysica Acta - Molecular and Cell Biology of Lipids</i> , 2010, 1801, 1239-1248.	2.4	55
34	A point mutation in the <i>neu1</i> promoter recruits an ectopic repressor, <i>Nkx3.2</i> and results in a mouse model of sialidase deficiency. <i>Molecular Genetics and Metabolism</i> , 2009, 97, 43-52.	1.1	10
35	Enhanced Cellular Uptake of Remnant High-Density Lipoprotein Particles. <i>Circulation Research</i> , 2008, 103, 159-166.	4.5	32
36	Hyperhomocysteinemia induced by methionine supplementation does not independently cause atherosclerosis in C57BL/6J mice. <i>FASEB Journal</i> , 2008, 22, 2569-2578.	0.5	44

#	ARTICLE	IF	CITATIONS
37	Conversion of Low Density Lipoprotein-associated Phosphatidylcholine to Triacylglycerol by Primary Hepatocytes. <i>Journal of Biological Chemistry</i> , 2008, 283, 6449-6458.	3.4	33
38	Myocardial Infarction Following Atherosclerosis in Murine Models. <i>Current Drug Targets</i> , 2008, 9, 217-223.	2.1	9
39	The inhibition of endocytosis affects HDL-lipid uptake mediated by the human scavenger receptor class B type I. <i>Molecular Membrane Biology</i> , 2007, 24, 442-454.	2.0	21
40	Liver X Receptor Stimulates Cholesterol Efflux and Inhibits Expression of Proinflammatory Mediators in Human Airway Smooth Muscle Cells. <i>Molecular Endocrinology</i> , 2007, 21, 1324-1334.	3.7	46
41	Regulation of SR-BI-mediated selective lipid uptake in Chinese hamster ovary-derived cells by protein kinase signaling pathways. <i>Journal of Lipid Research</i> , 2007, 48, 405-416.	4.2	28
42	Cholesterol depletion inhibits fatty acid uptake without affecting CD36 or caveolin-1 distribution in adipocytes. <i>Biochemical and Biophysical Research Communications</i> , 2007, 355, 67-71.	2.1	19
43	Hepatic Lipase Deficiency Delays Atherosclerosis, Myocardial Infarction, and Cardiac Dysfunction and Extends Lifespan in SR-BI/Apolipoprotein E Double Knockout Mice. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2006, 26, 548-554.	2.4	37
44	Hepatic high-density lipoprotein receptors: Roles in lipoprotein metabolism and potential for therapeutic modulation. <i>Current Atherosclerosis Reports</i> , 2005, 7, 344-350.	4.8	12
45	Biochemical Demonstration of the Involvement of Fatty Acyl-CoA Synthetase in Fatty Acid Translocation across the Plasma Membrane. <i>Journal of Biological Chemistry</i> , 2004, 279, 24163-24170.	3.4	19
46	Scavenger receptor class B type I in high-density lipoprotein metabolism, atherosclerosis and heart disease: lessons from gene-targeted mice. <i>Biochemical Society Transactions</i> , 2004, 32, 116-120.	3.4	34
47	Influence of the HDL Receptor SR-BI on Lipoprotein Metabolism and Atherosclerosis. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2003, 23, 1732-1738.	2.4	229
48	Scavenger Receptor Class B Type I-Mediated Protection Against Atherosclerosis in LDL Receptor-Negative Mice Involves Its Expression in Bone Marrow-Derived Cells. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2003, 23, 1589-1594.	2.4	205
49	Loss of SR-BI Expression Leads to the Early Onset of Occlusive Atherosclerotic Coronary Artery Disease, Spontaneous Myocardial Infarctions, Severe Cardiac Dysfunction, and Premature Death in Apolipoprotein E-Deficient Mice. <i>Circulation Research</i> , 2002, 90, 270-276.	4.5	461
50	Failure of red blood cell maturation in mice with defects in the high-density lipoprotein receptor SR-BI. <i>Blood</i> , 2002, 99, 1817-1824.	1.4	115
51	Failure of red blood cell maturation in mice with defects in the high-density lipoprotein receptor SR-BI. <i>Blood</i> , 2002, 99, 1817-1824.	1.4	111
52	The role of the high-density lipoprotein receptor SR-BI in cholesterol metabolism. <i>Current Opinion in Lipidology</i> , 2000, 11, 123-131.	2.7	172
53	Cellular and physiological roles of SR-BI, a lipoprotein receptor which mediates selective lipid uptake. <i>Biochimica Et Biophysica Acta - Molecular and Cell Biology of Lipids</i> , 2000, 1529, 276-286.	2.4	60
54	Influence of the high density lipoprotein receptor SR-BI on reproductive and cardiovascular pathophysiology. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1999, 96, 9322-9327.	7.1	475

#	ARTICLE	IF	CITATIONS
55	Identification of Caveolin-1 as a Fatty Acid Binding Protein. <i>Biochemical and Biophysical Research Communications</i> , 1999, 255, 34-39.	2.1	193
56	The Efficient Cellular Uptake of High Density Lipoprotein Lipids via Scavenger Receptor Class B Type I Requires Not Only Receptor-mediated Surface Binding but Also Receptor-specific Lipid Transfer Mediated by Its Extracellular Domain. <i>Journal of Biological Chemistry</i> , 1998, 273, 26338-26348.	3.4	198
57	Murine SR-BI, a High Density Lipoprotein Receptor That Mediates Selective Lipid Uptake, Is N-Glycosylated and Fatty Acylated and Colocalizes with Plasma Membrane Caveolae. <i>Journal of Biological Chemistry</i> , 1997, 272, 13242-13249.	3.4	330
58	Scavenger receptor class B, type I (SR-BI) is the major route for the delivery of high density lipoprotein cholesterol to the steroidogenic pathway in cultured mouse adrenocortical cells. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1997, 94, 13600-13605.	7.1	234
59	Scavenger receptor BI - a cell surface receptor for high density lipoprotein. <i>Current Opinion in Lipidology</i> , 1997, 8, 181-188.	2.7	185
60	A targeted mutation in the murine gene encoding the high density lipoprotein (HDL) receptor scavenger receptor class B type I reveals its key role in HDL metabolism. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1997, 94, 12610-12615.	7.1	797
61	The effect of intracellular pH on long-chain fatty acid uptake in 3T3-L1 adipocytes: evidence that uptake involves the passive diffusion of protonated long-chain fatty acids across the plasma membrane. <i>Biochemical Journal</i> , 1996, 313, 487-494.	3.7	35
62	A Single Point Mutation in μ -COP Results in Temperature-sensitive, Lethal Defects in Membrane Transport in a Chinese Hamster Ovary Cell Mutant. <i>Journal of Biological Chemistry</i> , 1996, 271, 11191-11196.	3.4	67
63	Membrane permeation and intracellular trafficking of long chain fatty acids: insights from <i>Escherichia coli</i> and 3T3-L1 adipocytes. <i>Biochemistry and Cell Biology</i> , 1995, 73, 223-234.	2.0	24
64	Identification of high affinity membrane-bound fatty acid-binding proteins using a photoreactive fatty acid. <i>Molecular and Cellular Biochemistry</i> , 1993, 123, 39-44.	3.1	35
65	Fatty acid uptake in <i>Candida tropicalis</i> : induction of a saturable process. <i>Biochemistry and Cell Biology</i> , 1992, 70, 76-80.	2.0	14