

Davy Vancampfort

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4594280/publications.pdf>

Version: 2024-02-01

418
papers

29,789
citations

4960

84
h-index

7348

152
g-index

440
all docs

440
docs citations

440
times ranked

22598
citing authors

#	ARTICLE	IF	CITATIONS
1	Prevalence, incidence and mortality from cardiovascular disease in patients with pooled and specific severe mental illness: a large-scale meta-analysis of 3,211,768 patients and 113,383,368 controls. <i>World Psychiatry</i> , 2017, 16, 163-180.	10.4	1,082
2	Exercise as a treatment for depression: A meta-analysis adjusting for publication bias. <i>Journal of Psychiatric Research</i> , 2016, 77, 42-51.	3.1	950
3	Physical Activity and Incident Depression: A Meta-Analysis of Prospective Cohort Studies. <i>American Journal of Psychiatry</i> , 2018, 175, 631-648.	7.2	933
4	Risk of metabolic syndrome and its components in people with schizophrenia and related psychotic disorders, bipolar disorder and major depressive disorder: a systematic review and meta-analysis. <i>World Psychiatry</i> , 2015, 14, 339-347.	10.4	858
5	The Lancet Psychiatry Commission: a blueprint for protecting physical health in people with mental illness. <i>Lancet Psychiatry</i> , 2019, 6, 675-712.	7.4	815
6	Prevalence of Metabolic Syndrome and Metabolic Abnormalities in Schizophrenia and Related Disorders—A Systematic Review and Meta-Analysis. <i>Schizophrenia Bulletin</i> , 2013, 39, 306-318.	4.3	813
7	Sedentary behavior and physical activity levels in people with schizophrenia, bipolar disorder and major depressive disorder: a global systematic review and meta-analysis. <i>World Psychiatry</i> , 2017, 16, 308-315.	10.4	600
8	Exercise for the Treatment of Depression. <i>The Open Complementary Medicine Journal</i> , 2009, 1, 78-83.	1.5	595
9	Metabolic syndrome in people with schizophrenia: a review. <i>World Psychiatry</i> , 2009, 8, 15-22.	10.4	495
10	Diabetes mellitus in people with schizophrenia, bipolar disorder and major depressive disorder: a systematic review and large scale meta-analysis. <i>World Psychiatry</i> , 2016, 15, 166-174.	10.4	487
11	A meta-review of "lifestyle psychiatry": the role of exercise, smoking, diet and sleep in the prevention and treatment of mental disorders. <i>World Psychiatry</i> , 2020, 19, 360-380.	10.4	424
12	Physical activity and sedentary behavior in people with major depressive disorder: A systematic review and meta-analysis. <i>Journal of Affective Disorders</i> , 2017, 210, 139-150.	4.1	411
13	An examination of the anxiolytic effects of exercise for people with anxiety and stress-related disorders: A meta-analysis. <i>Psychiatry Research</i> , 2017, 249, 102-108.	3.3	402
14	EPA guidance on physical activity as a treatment for severe mental illness: a meta-review of the evidence and Position Statement from the European Psychiatric Association (EPA), supported by the International Organization of Physical Therapists in Mental Health (IOPTMH). <i>European Psychiatry</i> , 2018, 54, 124-144.	0.2	377
15	Motivating factors and barriers towards exercise in severe mental illness: a systematic review and meta-analysis. <i>Psychological Medicine</i> , 2016, 46, 2869-2881.	4.5	345
16	Metabolic Syndrome and Metabolic Abnormalities in Bipolar Disorder: A Meta-Analysis of Prevalence Rates and Moderators. <i>American Journal of Psychiatry</i> , 2013, 170, 265-274.	7.2	336
17	Effect of aerobic exercise on hippocampal volume in humans: A systematic review and meta-analysis. <i>NeuroImage</i> , 2018, 166, 230-238.	4.2	334
18	How much physical activity do people with schizophrenia engage in? A systematic review, comparative meta-analysis and meta-regression. <i>Schizophrenia Research</i> , 2016, 176, 431-440.	2.0	284

#	ARTICLE	IF	CITATIONS
19	The intriguing relationship between coronary heart disease and mental disorders. <i>Dialogues in Clinical Neuroscience</i> , 2018, 20, 31-40.	3.7	282
20	Guideline concordant monitoring of metabolic risk in people treated with antipsychotic medication: systematic review and meta-analysis of screening practices. <i>Psychological Medicine</i> , 2012, 42, 125-147.	4.5	278
21	The effect of active video games on cognitive functioning in clinical and non-clinical populations: A meta-analysis of randomized controlled trials. <i>Neuroscience and Biobehavioral Reviews</i> , 2017, 78, 34-43.	6.1	273
22	Aerobic Exercise Improves Cognitive Functioning in People With Schizophrenia: A Systematic Review and Meta-Analysis. <i>Schizophrenia Bulletin</i> , 2017, 43, sbw115.	4.3	270
23	Is the Prevalence of Metabolic Syndrome and Metabolic Abnormalities Increased in Early Schizophrenia? A Comparative Meta-Analysis of First Episode, Untreated and Treated Patients. <i>Schizophrenia Bulletin</i> , 2013, 39, 295-305.	4.3	259
24	A systematic review of correlates of physical activity in patients with schizophrenia. <i>Acta Psychiatrica Scandinavica</i> , 2012, 125, 352-362.	4.5	255
25	The "online brain" how the Internet may be changing our cognition. <i>World Psychiatry</i> , 2019, 18, 119-129.	10.4	248
26	Exercise therapy improves both mental and physical health in patients with major depression. <i>Disability and Rehabilitation</i> , 2015, 37, 1490-1495.	1.8	246
27	Physical activity protects from incident anxiety: A meta-analysis of prospective cohort studies. <i>Depression and Anxiety</i> , 2019, 36, 846-858.	4.1	226
28	Physical activity in the treatment of Post-traumatic stress disorder: A systematic review and meta-analysis. <i>Psychiatry Research</i> , 2015, 230, 130-136.	3.3	224
29	Metabolic syndrome and metabolic abnormalities in patients with major depressive disorder: a meta-analysis of prevalences and moderating variables. <i>Psychological Medicine</i> , 2014, 44, 2017-2028.	4.5	223
30	Exercise as Medicine for Mental and Substance Use Disorders: A Meta-review of the Benefits for Neuropsychiatric and Cognitive Outcomes. <i>Sports Medicine</i> , 2020, 50, 151-170.	6.5	222
31	A meta-analysis of cardio-metabolic abnormalities in drug naïve, first-episode and multi-episode patients with schizophrenia versus general population controls. <i>World Psychiatry</i> , 2013, 12, 240-250.	10.4	220
32	Weight gain and obesity in schizophrenia: epidemiology, pathobiology, and management. <i>Acta Psychiatrica Scandinavica</i> , 2015, 132, 97-108.	4.5	216
33	Physical activity and anxiety: A perspective from the World Health Survey. <i>Journal of Affective Disorders</i> , 2017, 208, 545-552.	4.1	211
34	Dropout from exercise randomized controlled trials among people with depression: A meta-analysis and meta regression. <i>Journal of Affective Disorders</i> , 2016, 190, 457-466.	4.1	202
35	A Clinical Review of the Treatment of Catatonia. <i>Frontiers in Psychiatry</i> , 2014, 5, 181.	2.6	195
36	Type 2 Diabetes Mellitus in Youth Exposed to Antipsychotics. <i>JAMA Psychiatry</i> , 2016, 73, 247.	11.0	189

#	ARTICLE	IF	CITATIONS
37	The prevalence and predictors of type two diabetes mellitus in people with schizophrenia: a systematic review and comparative meta-analysis. <i>Acta Psychiatrica Scandinavica</i> , 2015, 132, 144-157.	4.5	183
38	Associations of moderate to vigorous physical activity and sedentary behavior with depressive and anxiety symptoms in self-isolating people during the COVID-19 pandemic: A cross-sectional survey in Brazil. <i>Psychiatry Research</i> , 2020, 292, 113339.	3.3	176
39	Prevalence and predictors of treatment dropout from physical activity interventions in schizophrenia: a meta-analysis. <i>General Hospital Psychiatry</i> , 2016, 39, 15-23.	2.4	172
40	Cardiorespiratory Fitness in Severe Mental Illness: A Systematic Review and Meta-analysis. <i>Sports Medicine</i> , 2017, 47, 343-352.	6.5	170
41	Guidelines for screening and monitoring of cardiometabolic risk in schizophrenia: systematic evaluation. <i>British Journal of Psychiatry</i> , 2011, 199, 99-105.	2.8	169
42	The prevalence and risk of metabolic syndrome and its components among people with posttraumatic stress disorder: a systematic review and meta-analysis. <i>Metabolism: Clinical and Experimental</i> , 2015, 64, 926-933.	3.4	167
43	Does loneliness contribute to mild cognitive impairment and dementia? A systematic review and meta-analysis of longitudinal studies. <i>Ageing Research Reviews</i> , 2019, 52, 7-16.	10.9	167
44	Exercise for depression in older adults: a meta-analysis of randomized controlled trials adjusting for publication bias. <i>Revista Brasileira De Psiquiatria</i> , 2016, 38, 247-254.	1.7	160
45	The impact of pharmacological and non-pharmacological interventions to improve physical health outcomes in people with schizophrenia: a meta-review of meta-analyses of randomized controlled trials. <i>World Psychiatry</i> , 2019, 18, 53-66.	10.4	153
46	The association of depression and all-cause and cause-specific mortality: an umbrella review of systematic reviews and meta-analyses. <i>BMC Medicine</i> , 2018, 16, 112.	5.5	143
47	Physical activity and suicidal ideation: A systematic review and meta-analysis. <i>Journal of Affective Disorders</i> , 2018, 225, 438-448.	4.1	140
48	Bullying Victimization and Suicide Attempt Among Adolescents Aged 12-15 Years From 48 Countries. <i>Journal of the American Academy of Child and Adolescent Psychiatry</i> , 2019, 58, 907-918.e4.	0.5	140
49	The efficacy and safety of nutrient supplements in the treatment of mental disorders: a meta-review of meta-analyses of randomized-controlled trials. <i>World Psychiatry</i> , 2019, 18, 308-324.	10.4	139
50	TYPE 2 DIABETES IN PATIENTS WITH MAJOR DEPRESSIVE DISORDER: A META-ANALYSIS OF PREVALENCE ESTIMATES AND PREDICTORS. <i>Depression and Anxiety</i> , 2015, 32, 763-773.	4.1	138
51	Exercise improves depressive symptoms in older adults: An umbrella review of systematic reviews and meta-analyses. <i>Psychiatry Research</i> , 2016, 244, 202-209.	3.3	137
52	A large-scale meta-analytic atlas of mental health problems prevalence during the COVID-19 early pandemic. <i>Journal of Medical Virology</i> , 2022, 94, 1935-1949.	5.0	134
53	The importance of self-determined motivation towards physical activity in patients with schizophrenia. <i>Psychiatry Research</i> , 2013, 210, 812-818.	3.3	133
54	Exercise improves cardiorespiratory fitness in people with depression: A meta-analysis of randomized control trials. <i>Journal of Affective Disorders</i> , 2016, 190, 249-253.	4.1	132

#	ARTICLE	IF	CITATIONS
55	Relationships between physical fitness, physical activity, smoking and metabolic and mental health parameters in people with schizophrenia. <i>Psychiatry Research</i> , 2013, 207, 25-32.	3.3	131
56	Moving to Beat Anxiety: Epidemiology and Therapeutic Issues with Physical Activity for Anxiety. <i>Current Psychiatry Reports</i> , 2018, 20, 63.	4.5	127
57	Selection, Use and Psychometric Properties of Physical Activity Measures to Assess Individuals with Severe Mental Illness: A Narrative Synthesis. <i>Archives of Psychiatric Nursing</i> , 2014, 28, 135-151.	1.4	126
58	Considering a frame of reference for physical activity research related to the cardiometabolic risk profile in schizophrenia. <i>Psychiatry Research</i> , 2010, 177, 271-279.	3.3	125
59	Association Between Gait Speed With Mortality, Cardiovascular Disease and Cancer: A Systematic Review and Meta-analysis of Prospective Cohort Studies. <i>Journal of the American Medical Directors Association</i> , 2018, 19, 981-988.e7.	2.5	123
60	International Organization of Physical Therapy in Mental Health consensus on physical activity within multidisciplinary rehabilitation programmes for minimising cardio-metabolic risk in patients with schizophrenia. <i>Disability and Rehabilitation</i> , 2012, 34, 1-12.	1.8	122
61	The epidemiology of back pain and its relationship with depression, psychosis, anxiety, sleep disturbances, and stress sensitivity: Data from 43 low- and middle-income countries. <i>General Hospital Psychiatry</i> , 2016, 43, 63-70.	2.4	122
62	Associations between sedentary behaviour and metabolic parameters in patients with schizophrenia. <i>Psychiatry Research</i> , 2012, 200, 73-78.	3.3	120
63	The prevalence and predictors of obstructive sleep apnea in major depressive disorder, bipolar disorder and schizophrenia: A systematic review and meta-analysis. <i>Journal of Affective Disorders</i> , 2016, 197, 259-267.	4.1	120
64	Exercise as treatment for alcohol use disorders: systematic review and meta-analysis. <i>British Journal of Sports Medicine</i> , 2017, 51, 1058-1064.	6.7	120
65	Physical multimorbidity and psychosis: comprehensive cross sectional analysis including 242,952 people across 48 low- and middle-income countries. <i>BMC Medicine</i> , 2016, 14, 189.	5.5	119
66	Exercise improves physical and psychological quality of life in people with depression: A meta-analysis including the evaluation of control group response. <i>Psychiatry Research</i> , 2016, 241, 47-54.	3.3	118
67	Chronic physical conditions, multimorbidity and physical activity across 46 low- and middle-income countries. <i>International Journal of Behavioral Nutrition and Physical Activity</i> , 2017, 14, 6.	4.6	115
68	Systematic Review of the Benefits of Physical Therapy Within a Multidisciplinary Care Approach for People With Schizophrenia. <i>Physical Therapy</i> , 2012, 92, 11-23.	2.4	114
69	Is Pain Perception Altered in People With Depression? A Systematic Review and Meta-Analysis of Experimental Pain Research. <i>Journal of Pain</i> , 2016, 17, 1257-1272.	1.4	114
70	Relationships between obesity, functional exercise capacity, physical activity participation and physical self-perception in people with schizophrenia. <i>Acta Psychiatrica Scandinavica</i> , 2011, 123, 423-430.	4.5	113
71	Physical activity and sedentary behavior in people with bipolar disorder: A systematic review and meta-analysis. <i>Journal of Affective Disorders</i> , 2016, 201, 145-152.	4.1	109
72	A review of physical activity correlates in patients with bipolar disorder. <i>Journal of Affective Disorders</i> , 2013, 145, 285-291.	4.1	108

#	ARTICLE	IF	CITATIONS
73	Promotion of cardiorespiratory fitness in schizophrenia: a clinical overview and meta-analysis. <i>Acta Psychiatrica Scandinavica</i> , 2015, 132, 131-143.	4.5	108
74	Relationship between sedentary behavior and depression: A mediation analysis of influential factors across the lifespan among 42,469 people in low- and middle-income countries. <i>Journal of Affective Disorders</i> , 2018, 229, 231-238.	4.1	107
75	Depression and physical health multimorbidity: primary data and country-wide meta-analysis of population data from 190 593 people across 43 low- and middle-income countries. <i>Psychological Medicine</i> , 2017, 47, 2107-2117.	4.5	106
76	Global physical activity levels among people living with HIV: a systematic review and meta-analysis. <i>Disability and Rehabilitation</i> , 2018, 40, 388-397.	1.8	100
77	Neurocognition in clinical high risk young adults who did or did not convert to a first schizophrenic psychosis: A meta-analysis. <i>Schizophrenia Research</i> , 2013, 149, 48-55.	2.0	97
78	Decreased pain sensitivity among people with schizophrenia. <i>Pain</i> , 2015, 156, 2121-2131.	4.2	95
79	Sedentary behavior and depressive symptoms among 67,077 adolescents aged 12-15 years from 30 low- and middle-income countries. <i>International Journal of Behavioral Nutrition and Physical Activity</i> , 2018, 15, 73.	4.6	95
80	Yoga in schizophrenia: a systematic review of randomised controlled trials. <i>Acta Psychiatrica Scandinavica</i> , 2012, 126, 12-20.	4.5	94
81	Bone mineral density, osteoporosis, and fractures among people with eating disorders: a systematic review and meta-analysis. <i>Acta Psychiatrica Scandinavica</i> , 2016, 133, 341-351.	4.5	94
82	Passive and mentally-active sedentary behaviors and incident major depressive disorder: A 13-year cohort study. <i>Journal of Affective Disorders</i> , 2018, 241, 579-585.	4.1	93
83	Exercise improves cardiorespiratory fitness in people with schizophrenia: A systematic review and meta-analysis. <i>Schizophrenia Research</i> , 2015, 169, 453-457.	2.0	92
84	State anxiety, psychological stress and positive well-being responses to yoga and aerobic exercise in people with schizophrenia: a pilot study. <i>Disability and Rehabilitation</i> , 2011, 33, 684-689.	1.8	91
85	Lack of physical activity during leisure time contributes to an impaired health related quality of life in patients with schizophrenia. <i>Schizophrenia Research</i> , 2011, 129, 122-127.	2.0	91
86	Barriers to and Facilitators of Physical Activity Among Persons With Schizophrenia: A Survey of Physical Therapists. <i>Psychiatric Services</i> , 2014, 65, 693-696.	2.0	88
87	Physical activity and depression: a large cross-sectional, population-based study across 36 low- and middle-income countries. <i>Acta Psychiatrica Scandinavica</i> , 2016, 134, 546-556.	4.5	88
88	Chronic Physical Conditions, Multimorbidity, and Mild Cognitive Impairment in Low- and Middle-Income Countries. <i>Journal of the American Geriatrics Society</i> , 2018, 66, 721-727.	2.6	87
89	Adopting and maintaining physical activity behaviours in people with severe mental illness: The importance of autonomous motivation. <i>Preventive Medicine</i> , 2015, 81, 216-220.	3.4	86
90	Are lower levels of cardiorespiratory fitness associated with incident depression? A systematic review of prospective cohort studies. <i>Preventive Medicine</i> , 2016, 93, 159-165.	3.4	85

#	ARTICLE	IF	CITATIONS
91	Prevalence and Predictors of Type 2 Diabetes Mellitus in People With Bipolar Disorder. <i>Journal of Clinical Psychiatry</i> , 2015, 76, 1490-1499.	2.2	85
92	Does exercise improve sleep quality in individuals with mental illness? A systematic review and meta-analysis. <i>Journal of Psychiatric Research</i> , 2019, 109, 96-106.	3.1	83
93	A systematic review of physical therapy interventions for patients with anorexia and bulimia nervosa. <i>Disability and Rehabilitation</i> , 2014, 36, 628-634.	1.8	80
94	The relationship between chronic physical conditions, multimorbidity and anxiety in the general population: A global perspective across 42 countries. <i>General Hospital Psychiatry</i> , 2017, 45, 1-6.	2.4	80
95	Depression comorbid with tuberculosis and its impact on health status: cross-sectional analysis of community-based data from 48 low- and middle-income countries. <i>BMC Medicine</i> , 2017, 15, 209.	5.5	79
96	The prevalence of pain in bipolar disorder: a systematic review and large-scale meta-analysis. <i>Acta Psychiatrica Scandinavica</i> , 2015, 131, 75-88.	4.5	78
97	Negative symptoms are associated with lower autonomous motivation towards physical activity in people with schizophrenia. <i>Comprehensive Psychiatry</i> , 2015, 56, 128-132.	3.1	77
98	The Validity and Value of Self-reported Physical Activity and Accelerometry in People With Schizophrenia: A Population-Scale Study of the UK Biobank. <i>Schizophrenia Bulletin</i> , 2018, 44, 1293-1300.	4.3	77
99	A meta-analysis of prevalence estimates and moderators of low bone mass in people with schizophrenia. <i>Acta Psychiatrica Scandinavica</i> , 2014, 130, 470-486.	4.5	76
100	Schizophrenia and the risk of fractures: a systematic review and comparative meta-analysis. <i>General Hospital Psychiatry</i> , 2015, 37, 126-133.	2.4	76
101	The transcending benefits of physical activity for individuals with schizophrenia: A systematic review and meta-ethnography. <i>Psychiatry Research</i> , 2014, 220, 11-19.	3.3	75
102	Food insecurity (hunger) and suicide attempts among 179,771 adolescents attending school from 9 high-income, 31 middle-income, and 4 low-income countries: A cross-sectional study. <i>Journal of Affective Disorders</i> , 2019, 248, 91-98.	4.1	75
103	The functional exercise capacity is correlated with global functioning in patients with schizophrenia. <i>Acta Psychiatrica Scandinavica</i> , 2012, 125, 382-387.	4.5	74
104	Prevalence and clinical features associated with bipolar disorder polypharmacy: a systematic review. <i>Neuropsychiatric Disease and Treatment</i> , 2016, 12, 719.	2.2	74
105	Type 2 Diabetes Among People With Posttraumatic Stress Disorder. <i>Psychosomatic Medicine</i> , 2016, 78, 465-473.	2.0	73
106	Assessing physical activity in people with mental illness: 23-country reliability and validity of the simple physical activity questionnaire (SIMPAQ). <i>BMC Psychiatry</i> , 2020, 20, 108.	2.6	73
107	The Prevalence of Metabolic Syndrome in Alcohol Use Disorders: A Systematic Review and Meta-analysis. <i>Alcohol and Alcoholism</i> , 2016, 51, 515-521.	1.6	72
108	Do people with mental illness receive adequate smoking cessation advice? A systematic review and meta-analysis. <i>General Hospital Psychiatry</i> , 2015, 37, 14-23.	2.4	71

#	ARTICLE	IF	CITATIONS
109	Cross-sectional and prospective relationships of passive and mentally active sedentary behaviours and physical activity with depression. <i>British Journal of Psychiatry</i> , 2020, 217, 413-419.	2.8	71
110	The prevalence and moderators of clinical pain in people with schizophrenia: A systematic review and large scale meta-analysis. <i>Schizophrenia Research</i> , 2014, 160, 1-8.	2.0	70
111	Depression and pain: primary data and meta-analysis among 237,952 people across 47 low- and middle-income countries. <i>Psychological Medicine</i> , 2017, 47, 2906-2917.	4.5	70
112	Physical chronic conditions, multimorbidity and sedentary behavior amongst middle-aged and older adults in six low- and middle-income countries. <i>International Journal of Behavioral Nutrition and Physical Activity</i> , 2017, 14, 147.	4.6	70
113	Cannabis use and suicide attempts among 86,254 adolescents aged 12-15 years from 21 low- and middle-income countries. <i>European Psychiatry</i> , 2019, 56, 8-13.	0.2	70
114	Impact of Psychotropic Medication Effects on Obesity and the Metabolic Syndrome in People With Serious Mental Illness. <i>Frontiers in Endocrinology</i> , 2020, 11, 573479.	3.5	70
115	Physical activity and sedentary behaviour in outpatients with schizophrenia: A systematic review and meta-analysis. <i>International Journal of Therapy and Rehabilitation</i> , 2013, 20, 588-595.	0.3	69
116	Integrating physical activity as medicine in the care of people with severe mental illness. <i>Australian and New Zealand Journal of Psychiatry</i> , 2015, 49, 681-682.	2.3	69
117	What are the factors associated with physical activity (PA) participation in community dwelling adults with dementia? A systematic review of PA correlates. <i>Archives of Gerontology and Geriatrics</i> , 2014, 59, 195-203.	3.0	67
118	The association between smoking prevalence and eating disorders: a systematic review and meta-analysis. <i>Addiction</i> , 2016, 111, 1914-1922.	3.3	65
119	Physical activity correlates in people living with HIV/AIDS: a systematic review of 45 studies. <i>Disability and Rehabilitation</i> , 2018, 40, 1618-1629.	1.8	65
120	Neurobiological effects of physical exercise in schizophrenia: a systematic review. <i>Disability and Rehabilitation</i> , 2014, 36, 1749-1754.	1.8	63
121	Handgrip strength and depression among 34,129 adults aged 50 years and older in six low- and middle-income countries. <i>Journal of Affective Disorders</i> , 2019, 243, 448-454.	4.1	63
122	Identifying the facilitators and processes which influence recovery in individuals with schizophrenia: a systematic review and thematic synthesis. <i>Journal of Mental Health</i> , 2015, 24, 103-110.	1.9	62
123	Perceived Stress and Its Relationship With Chronic Medical Conditions and Multimorbidity Among 229,293 Community-Dwelling Adults in 44 Low- and Middle-Income Countries. <i>American Journal of Epidemiology</i> , 2017, 186, 979-989.	3.4	62
124	Grip Strength Is Associated With Cognitive Performance in Schizophrenia and the General Population: A UK Biobank Study of 476559 Participants. <i>Schizophrenia Bulletin</i> , 2018, 44, 728-736.	4.3	62
125	Nutritional Deficiencies and Clinical Correlates in First-Episode Psychosis: A Systematic Review and Meta-analysis. <i>Schizophrenia Bulletin</i> , 2018, 44, 1275-1292.	4.3	61
126	Is it possible for people with severe mental illness to sit less and move more? A systematic review of interventions to increase physical activity or reduce sedentary behaviour. <i>Schizophrenia Research</i> , 2018, 202, 3-16.	2.0	60

#	ARTICLE	IF	CITATIONS
127	Are leptin levels increased among people with schizophrenia versus controls? A systematic review and comparative meta-analysis. <i>Psychoneuroendocrinology</i> , 2016, 63, 144-154.	2.7	58
128	A systematic review on physical therapy interventions for patients with binge eating disorder. <i>Disability and Rehabilitation</i> , 2013, 35, 2191-2196.	1.8	55
129	Environmental risk factors and nonpharmacological and nonsurgical interventions for obesity: An umbrella review of meta-analyses of cohort studies and randomized controlled trials. <i>European Journal of Clinical Investigation</i> , 2018, 48, e12982.	3.4	55
130	Markers of inflammation in schizophrenia: association vs. causation. <i>World Psychiatry</i> , 2014, 13, 189-192.	10.4	54
131	Treatment Guidelines for Depression: Greater Emphasis on Physical Activity is Needed. <i>European Psychiatry</i> , 2017, 40, 1-3.	0.2	54
132	Association Between Muscular Strength and Cognition in People With Major Depression or Bipolar Disorder and Healthy Controls. <i>JAMA Psychiatry</i> , 2018, 75, 740.	11.0	54
133	Reliability, minimal detectable changes, practice effects and correlates of the 6-min walk test in patients with schizophrenia. <i>Psychiatry Research</i> , 2011, 187, 62-67.	3.3	53
134	Association between depression and smoking: A global perspective from 48 low- and middle-income countries. <i>Journal of Psychiatric Research</i> , 2018, 103, 142-149.	3.1	53
135	Correlates of sedentary behavior in the general population: A cross-sectional study using nationally representative data from six low- and middle-income countries. <i>PLoS ONE</i> , 2018, 13, e0202222.	2.5	53
136	Association of the metabolic syndrome with physical activity performance in patients with schizophrenia. <i>Diabetes and Metabolism</i> , 2011, 37, 318-323.	2.9	52
137	Physiotherapists can help implement physical activity programmes in clinical practice. <i>British Journal of Psychiatry</i> , 2014, 204, 164-164.	2.8	52
138	State anxiety and subjective well-being responses to acute bouts of aerobic exercise in patients with depressive and anxiety disorders. <i>British Journal of Sports Medicine</i> , 2009, 43, 756-759.	6.7	51
139	What are the factors that influence physical activity participation in individuals with depression? A review of physical activity correlates from 59 studies. <i>Psychiatria Danubina</i> , 2015, 27, 210-24.	0.4	51
140	Effects of progressive muscle relaxation on state anxiety and subjective well-being in people with schizophrenia: a randomized controlled trial. <i>Clinical Rehabilitation</i> , 2011, 25, 567-575.	2.2	50
141	A systematic review of the benefits of physical therapy within a multidisciplinary care approach for people with schizophrenia: An update. <i>Psychiatry Research</i> , 2015, 229, 828-839.	3.3	50
142	Physical activity and generalized anxiety disorder: results from The Irish Longitudinal Study on Ageing (TILDA). <i>International Journal of Epidemiology</i> , 2018, 47, 1443-1453.	1.9	50
143	Physical Activity Levels and Psychosis: A Mediation Analysis of Factors Influencing Physical Activity Target Achievement Among 204 186 People Across 46 Low- and Middle-Income Countries. <i>Schizophrenia Bulletin</i> , 2017, 43, sbw111.	4.3	49
144	Cardiometabolic effects of physical activity interventions for people with schizophrenia. <i>Physical Therapy Reviews</i> , 2009, 14, 388-398.	0.8	48

#	ARTICLE	IF	CITATIONS
145	Physical Activity in People With Posttraumatic Stress Disorder: A Systematic Review of Correlates. <i>Journal of Physical Activity and Health</i> , 2016, 13, 910-918.	2.0	48
146	Progressive muscle relaxation in persons with schizophrenia: a systematic review of randomized controlled trials. <i>Clinical Rehabilitation</i> , 2013, 27, 291-298.	2.2	47
147	Could autonomous motivation hold the key to successfully implementing lifestyle changes in affective disorders? A multicentre cross sectional study. <i>Psychiatry Research</i> , 2015, 228, 100-106.	3.3	47
148	The Association Between Sedentary Behavior and Sarcopenia Among Adults Aged ≥65 Years in Low- and Middle-Income Countries. <i>International Journal of Environmental Research and Public Health</i> , 2020, 17, 1708.	2.6	47
149	Physical activity and mental health. <i>Lancet Psychiatry</i> , 2018, 5, 873.	7.4	46
150	Predictors of Falls and Fractures Leading to Hospitalization in People With Dementia: A Representative Cohort Study. <i>Journal of the American Medical Directors Association</i> , 2018, 19, 607-612.	2.5	46
151	A Systematic Review of Physical Activity Correlates in Alcohol Use Disorders. <i>Archives of Psychiatric Nursing</i> , 2015, 29, 196-201.	1.4	45
152	Leisure-time sedentary behavior and loneliness among 148,045 adolescents aged 12–15 years from 52 low- and middle-income countries. <i>Journal of Affective Disorders</i> , 2019, 251, 149-155.	4.1	44
153	Handgrip Strength Is Associated With Hippocampal Volume and White Matter Hyperintensities in Major Depression and Healthy Controls: A UK Biobank Study. <i>Psychosomatic Medicine</i> , 2020, 82, 39-46.	2.0	44
154	Challenges Establishing the Efficacy of Exercise as an Antidepressant Treatment: A Systematic Review and Meta-Analysis of Control Group Responses in Exercise Randomised Controlled Trials. <i>Sports Medicine</i> , 2016, 46, 699-713.	6.5	43
155	Mild cognitive impairment and physical activity in the general population: Findings from six low- and middle-income countries. <i>Experimental Gerontology</i> , 2017, 100, 100-105.	2.8	43
156	Multimorbidity and perceived stress: a population-based cross-sectional study among older adults across six low- and middle-income countries. <i>Maturitas</i> , 2018, 107, 84-91.	2.4	43
157	Perceived Stress and Mild Cognitive Impairment among 32,715 Community-Dwelling Older Adults across Six Low- and Middle-Income Countries. <i>Gerontology</i> , 2019, 65, 155-163.	2.8	43
158	Why moving more should be promoted for severe mental illness. <i>Lancet Psychiatry</i> , 2015, 2, 295.	7.4	42
159	How can we promote smoking cessation in people with schizophrenia in practice? A clinical overview. <i>Acta Psychiatrica Scandinavica</i> , 2015, 132, 122-130.	4.5	42
160	Dropout from physical activity interventions in people living with HIV: a systematic review and meta-analysis. <i>AIDS Care - Psychological and Socio-Medical Aspects of AIDS/HIV</i> , 2017, 29, 636-643.	1.2	42
161	Associations of sedentary behavior in leisure and occupational contexts with symptoms of depression and anxiety. <i>Preventive Medicine</i> , 2020, 133, 106021.	3.4	42
162	Moderate to vigorous physical activity and sedentary behavior changes in self-isolating adults during the COVID-19 pandemic in Brazil: a cross-sectional survey exploring correlates. <i>Sport Sciences for Health</i> , 2022, 18, 155-163.	1.3	42

#	ARTICLE	IF	CITATIONS
163	The anxiolytic effects of exercise for people with anxiety and related disorders: An update of the available meta-analytic evidence. <i>Psychiatry Research</i> , 2021, 302, 114046.	3.3	42
164	Self-determination and stage of readiness to change physical activity behaviour in schizophrenia. <i>Mental Health and Physical Activity</i> , 2014, 7, 171-176.	1.8	41
165	Low Physical Activity and Cardiorespiratory Fitness in People With Schizophrenia: A Comparison With Matched Healthy Controls and Associations With Mental and Physical Health. <i>Frontiers in Psychiatry</i> , 2019, 10, 87.	2.6	41
166	From impact factors to real impact: translating evidence on lifestyle interventions into routine mental health care. <i>Translational Behavioral Medicine</i> , 2020, 10, 1070-1073.	2.4	41
167	Investigating the benefits of sport participation for individuals with schizophrenia: a systematic review. <i>Psychiatria Danubina</i> , 2015, 27, 2-13.	0.4	41
168	Depressive symptoms, handgrip strength, and weight status in US older adults. <i>Journal of Affective Disorders</i> , 2018, 238, 305-310.	4.1	39
169	Sedentary behavior and anxiety: Association and influential factors among 42,469 community-dwelling adults in six low- and middle-income countries. <i>General Hospital Psychiatry</i> , 2018, 50, 26-32.	2.4	38
170	Food Insecurity Is Associated with Mild Cognitive Impairment among Middle-Aged and Older Adults in South Africa: Findings from a Nationally Representative Survey. <i>Nutrients</i> , 2019, 11, 749.	4.1	38
171	Physical Activity Correlates in Persons with Binge Eating Disorder: A Systematic Review. <i>European Eating Disorders Review</i> , 2014, 22, 1-8.	4.1	37
172	Health related quality of life, physical fitness and physical activity participation in treatment-seeking obese persons with and without binge eating disorder. <i>Psychiatry Research</i> , 2014, 216, 97-102.	3.3	37
173	Associations between physical activity and the built environment in patients with schizophrenia: a multi-centre study. <i>General Hospital Psychiatry</i> , 2013, 35, 653-658.	2.4	36
174	Cardiorespiratory fitness levels and moderators in people with HIV: A systematic review and meta-analysis. <i>Preventive Medicine</i> , 2016, 93, 106-114.	3.4	36
175	A narrative synthesis investigating the use and value of social support to promote physical activity among individuals with schizophrenia. <i>Disability and Rehabilitation</i> , 2016, 38, 123-150.	1.8	36
176	Physical fitness in people with posttraumatic stress disorder: a systematic review. <i>Disability and Rehabilitation</i> , 2017, 39, 2461-2467.	1.8	36
177	Associations between handgrip strength and mild cognitive impairment in middle-aged and older adults in six low- and middle-income countries. <i>International Journal of Geriatric Psychiatry</i> , 2019, 34, 609-616.	2.7	36
178	Health-related quality of life and aerobic fitness in people with schizophrenia. <i>International Journal of Mental Health Nursing</i> , 2015, 24, 394-402.	3.8	35
179	Six minutes walk test for individuals with schizophrenia. <i>Disability and Rehabilitation</i> , 2015, 37, 921-927.	1.8	35
180	Association of leisure-time sedentary behavior with fast food and carbonated soft drink consumption among 133,555 adolescents aged 12-15 years in 44 low- and middle-income countries. <i>International Journal of Behavioral Nutrition and Physical Activity</i> , 2019, 16, 35.	4.6	35

#	ARTICLE	IF	CITATIONS
181	Associations of interruptions to leisure-time sedentary behaviour with symptoms of depression and anxiety. <i>Translational Psychiatry</i> , 2020, 10, 128.	4.8	35
182	The efficacy of meditation-based mind-body interventions for mental disorders: A meta-review of 17 meta-analyses of randomized controlled trials. <i>Journal of Psychiatric Research</i> , 2021, 134, 181-191.	3.1	35
183	Understanding the role of physiotherapists in schizophrenia: an international perspective from members of the International Organisation of Physical Therapists in Mental Health (IOPTMH). <i>Journal of Mental Health</i> , 2014, 23, 125-129.	1.9	34
184	Changes in physical activity, physical fitness, self-perception and quality of life following a 6-month physical activity counseling and cognitive behavioral therapy program in outpatients with binge eating disorder. <i>Psychiatry Research</i> , 2014, 219, 361-366.	3.3	34
185	Sexual behavior and suicide attempts among adolescents aged 12–15 years from 38 countries: A global perspective. <i>Psychiatry Research</i> , 2020, 287, 112564.	3.3	34
186	Autonomous motivation is associated with the maintenance stage of behaviour change in people with affective disorders. <i>Psychiatry Research</i> , 2016, 240, 267-271.	3.3	33
187	Sedentary behaviour and sleep problems among 42,489 community-dwelling adults in six low- and middle-income countries. <i>Journal of Sleep Research</i> , 2018, 27, e12714.	3.2	33
188	Sedentary behavior and anxiety-induced sleep disturbance among 181,093 adolescents from 67 countries: a global perspective. <i>Sleep Medicine</i> , 2019, 58, 19-26.	1.6	33
189	Dropout from exercise randomized controlled trials among people with anxiety and stress-related disorders: A meta-analysis and meta-regression. <i>Journal of Affective Disorders</i> , 2021, 282, 996-1004.	4.1	33
190	Physical and mental health impact of COVID-19 on children, adolescents, and their families: The Collaborative Outcomes study on Health and Functioning during Infection Times - Children and Adolescents (COH-FIT-C&A). <i>Journal of Affective Disorders</i> , 2022, 299, 367-376.	4.1	33
191	Yoga versus standard care for schizophrenia. <i>The Cochrane Library</i> , 2015, 2015, CD010554.	2.8	32
192	Physical activity as a vital sign in patients with schizophrenia: Evidence and clinical recommendations. <i>Schizophrenia Research</i> , 2016, 170, 336-340.	2.0	32
193	Hand grip strength and cognitive function among elderly cancer survivors. <i>PLoS ONE</i> , 2018, 13, e0197909.	2.5	32
194	Leisure-time sedentary behavior and suicide attempt among 126,392 adolescents in 43 countries. <i>Journal of Affective Disorders</i> , 2019, 250, 346-353.	4.1	32
195	Correlates of low physical activity across 46 low- and middle-income countries: A cross-sectional analysis of community-based data. <i>Preventive Medicine</i> , 2018, 106, 107-113.	3.4	31
196	Visual impairment and objectively measured physical activity and sedentary behaviour in US adolescents and adults: a cross-sectional study. <i>BMJ Open</i> , 2019, 9, e027267.	1.9	31
197	Sedentary behavior and depression among community-dwelling adults aged ≥50 years: Results from the Irish Longitudinal Study on Ageing. <i>Journal of Affective Disorders</i> , 2020, 262, 389-396.	4.1	31
198	Eurofit test battery in patients with schizophrenia or schizoaffective disorder: Reliability and clinical correlates. <i>European Psychiatry</i> , 2012, 27, 416-421.	0.2	29

#	ARTICLE	IF	CITATIONS
199	The benefits of walking for individuals with schizophrenia spectrum disorders: A systematic review. <i>International Journal of Therapy and Rehabilitation</i> , 2014, 21, 410-420.	0.3	29
200	How common is bipolar disorder in general primary care attendees? A systematic review and meta-analysis investigating prevalence determined according to structured clinical assessments. <i>Australian and New Zealand Journal of Psychiatry</i> , 2016, 50, 631-639.	2.3	29
201	Leisure time physical activity and future psychological distress: A thirteen year longitudinal population-based study. <i>Journal of Psychiatric Research</i> , 2018, 101, 50-56.	3.1	29
202	Association of bullying victimization with overweight and obesity among adolescents from 41 low- and middle-income countries. <i>Pediatric Obesity</i> , 2020, 15, e12571.	2.8	29
203	An impaired health related muscular fitness contributes to a reduced walking capacity in patients with schizophrenia: a cross-sectional study. <i>BMC Psychiatry</i> , 2013, 13, 5.	2.6	28
204	Diabetes, physical activity participation and exercise capacity in patients with schizophrenia. <i>Psychiatry and Clinical Neurosciences</i> , 2013, 67, 451-456.	1.8	28
205	Lower Bone Mineral Density at the Hip and Lumbar Spine in People with Psychosis Versus Controls: a Comprehensive Review and Skeletal Site-Specific Meta-analysis. <i>Current Osteoporosis Reports</i> , 2016, 14, 249-259.	3.6	28
206	Correlates of physical activity among community-dwelling adults aged 50 or over in six low- and middle-income countries. <i>PLoS ONE</i> , 2017, 12, e0186992.	2.5	28
207	Associations between TV viewing and depressive symptoms among 60,202 Brazilian adults: The Brazilian national health survey. <i>Journal of Affective Disorders</i> , 2018, 236, 23-30.	4.1	28
208	Cardiorespiratory fitness and response to exercise treatment in depression. <i>BJPsych Open</i> , 2018, 4, 346-351.	0.7	28
209	The efficacy of exergaming in people with major neurocognitive disorder residing in long-term care facilities: a pilot randomized controlled trial. <i>Alzheimer's Research and Therapy</i> , 2021, 13, 70.	6.2	28
210	A Cognitive- Behavioral Therapeutic Program for Patients With Obesity and Binge Eating Disorder. <i>Behavior Modification</i> , 2012, 36, 670-686.	1.6	27
211	The prevalence of diabetes mellitus type 2 in people with alcohol use disorders: a systematic review and large scale meta-analysis. <i>Psychiatry Research</i> , 2016, 246, 394-400.	3.3	27
212	What are the top 10 physical activity research questions in schizophrenia?. <i>Disability and Rehabilitation</i> , 2016, 38, 2235-2243.	1.8	27
213	A comparison of physical fitness in patients with bipolar disorder, schizophrenia and healthy controls. <i>Disability and Rehabilitation</i> , 2016, 38, 2047-2051.	1.8	27
214	Is autonomous motivation the key to maintaining an active lifestyle in first-episode psychosis?. <i>Microbial Biotechnology</i> , 2018, 12, 821-827.	1.7	27
215	Predictors of falls and fractures leading to hospitalization in people with schizophrenia spectrum disorder: A large representative cohort study. <i>Schizophrenia Research</i> , 2018, 201, 70-78.	2.0	27
216	Leisure Time Sedentary Behavior and Obesity Among 116,762 Adolescents Aged 12-15 Years from 41 Low- and Middle-Income Countries. <i>Obesity</i> , 2019, 27, 830-836.	3.0	27

#	ARTICLE	IF	CITATIONS
217	Physical activity, exercise, and mental disorders: it is time to move on. Trends in Psychiatry and Psychotherapy, 2021, 43, 177-184.	0.8	27
218	The value of social support to encourage people with schizophrenia to engage in physical activity: an international insight from specialist mental health physiotherapists. Journal of Mental Health, 2014, 23, 256-260.	1.9	26
219	Reliability and clinical correlates of the Astrandâ€“Rhyming sub-maximal exercise test in patients with schizophrenia or schizoaffective disorder. Psychiatry Research, 2014, 220, 778-783.	3.3	26
220	More Reasons to Move: Exercise in the Treatment of Alcohol Use Disorders. Frontiers in Psychiatry, 2017, 8, 160.	2.6	26
221	Correlates of physical activity among 142,118 adolescents aged 12â€“15â€“years from 48 low- and middle-income countries. Preventive Medicine, 2019, 127, 105819.	3.4	26
222	Does physical activity reduce the risk of psychosis? A systematic review and meta-analysis of prospective studies. Psychiatry Research, 2020, 284, 112675.	3.3	26
223	Associations between perceived neighbourhood environmental attributes and self-reported sitting time in patients with schizophrenia: A pilot study. Psychiatry Research, 2014, 215, 33-38.	3.3	25
224	Health-related physical fitness in patients with bipolar disorder vs. healthy controls: An exploratory study. Journal of Affective Disorders, 2015, 177, 22-27.	4.1	25
225	Physical activity is associated with the physical, psychological, social and environmental quality of life in people with mental health problems in a low resource setting. Psychiatry Research, 2017, 258, 250-254.	3.3	25
226	Physical activity correlates among people with psychosis: Data from 47 low- and middle-income countries. Schizophrenia Research, 2018, 193, 412-417.	2.0	25
227	Handgrip strength is associated with suicidal thoughts in men: Crossâ€“sectional analyses from NHANES. Scandinavian Journal of Medicine and Science in Sports, 2020, 30, 92-99.	2.9	25
228	Prospective relationships of mentally passive sedentary behaviors with depression: Mediation by sleep problems. Journal of Affective Disorders, 2020, 265, 538-544.	4.1	25
229	Joint association of ultra-processed food and sedentary behavior with anxiety-induced sleep disturbance among Brazilian adolescents. Journal of Affective Disorders, 2020, 266, 135-142.	4.1	25
230	Is pain sensitivity altered in people with Alzheimer's disease? A systematic review and meta-analysis of experimental pain research. Experimental Gerontology, 2016, 82, 30-38.	2.8	24
231	Handgrip strength, chronic physical conditions and physical multimorbidity in middle-aged and older adults in six low- and middle income countries. European Journal of Internal Medicine, 2019, 61, 96-102.	2.2	24
232	Preferences and motivations for exercise in early psychosis. Acta Psychiatrica Scandinavica, 2016, 134, 83-84.	4.5	23
233	Impact of antipsychotic medication on physical activity and physical fitness in adolescents: An exploratory study. Psychiatry Research, 2016, 242, 192-197.	3.3	23
234	Associations between TV viewing, sitting time, physical activity and insomnia among 100,839 Brazilian adolescents. Psychiatry Research, 2018, 269, 700-706.	3.3	23

#	ARTICLE	IF	CITATIONS
235	Sex difference in the association between physical activity and suicide attempts among adolescents from 48 countries: A global perspective. <i>Journal of Affective Disorders</i> , 2020, 266, 311-318.	4.1	23
236	Can high-intensity interval training improve mental health outcomes in the general population and those with physical illnesses? A systematic review and meta-analysis. <i>British Journal of Sports Medicine</i> , 2022, 56, 279-291.	6.7	23
237	Increased rates of respiratory disease in schizophrenia: A systematic review and meta-analysis including 619,214 individuals with schizophrenia and 52,159,551 controls. <i>Schizophrenia Research</i> , 2021, 237, 131-140.	2.0	23
238	How can we increase physical activity and exercise among youth experiencing first-episode psychosis? A systematic review of intervention variables. <i>Microbial Biotechnology</i> , 2016, 10, 435-440.	1.7	22
239	Sedentary Behavior in People Living With HIV: A Systematic Review and Meta-Analysis. <i>Journal of Physical Activity and Health</i> , 2017, 14, 571-577.	2.0	22
240	Correlates of sedentary behavior in 2,375 people with depression from 6 low- and middle-income countries. <i>Journal of Affective Disorders</i> , 2018, 234, 97-104.	4.1	22
241	Mild cognitive impairment and sedentary behavior: A multinational study. <i>Experimental Gerontology</i> , 2018, 108, 174-180.	2.8	22
242	Psychotic experiences and subjective cognitive complaints among 224 842 people in 48 low- and middle-income countries. <i>Epidemiology and Psychiatric Sciences</i> , 2020, 29, e11.	3.9	22
243	The collaborative outcomes study on health and functioning during infection times in adults (COH-FIT-Adults): Design and methods of an international online survey targeting physical and mental health effects of the COVID-19 pandemic. <i>Journal of Affective Disorders</i> , 2022, 299, 393-407.	4.1	22
244	Quality Assessment of Physical Activity Recommendations Within Clinical Practice Guidelines for the Prevention and Treatment of Cardio-metabolic Risk Factors in People With Schizophrenia. <i>Community Mental Health Journal</i> , 2011, 47, 703-710.	2.0	21
245	Test-retest reliability, feasibility and clinical correlates of the Eurofit test battery in people with bipolar disorder. <i>Psychiatry Research</i> , 2015, 228, 620-625.	3.3	21
246	The functional exercise capacity and its correlates in obese treatment-seeking people with binge eating disorder: an exploratory study. <i>Disability and Rehabilitation</i> , 2015, 37, 777-782.	1.8	21
247	Cardiorespiratory fitness in outpatients with bipolar disorder versus matched controls: An exploratory study. <i>Journal of Affective Disorders</i> , 2016, 199, 1-5.	4.1	21
248	Physical activity correlates among 24,230 people with depression across 46 low- and middle-income countries. <i>Journal of Affective Disorders</i> , 2017, 221, 81-88.	4.1	21
249	Associations between active travel and physical multi-morbidity in six low- and middle-income countries among community-dwelling older adults: A cross-sectional study. <i>PLoS ONE</i> , 2018, 13, e0203277.	2.5	21
250	Sedentary behavior and perceived stress among adults aged ≥ 50 years in six low- and middle-income countries. <i>Maturitas</i> , 2018, 116, 100-107.	2.4	21
251	SOCIAL ANXIETY IN PHYSICAL ACTIVITY PARTICIPATION IN PATIENTS WITH MENTAL ILLNESS: A CROSS-SECTIONAL MULTICENTER STUDY. <i>Depression and Anxiety</i> , 2013, 30, 757-762.	4.1	20
252	A systematic review of physical activity policy recommendations and interventions for people with mental health problems in Sub-Saharan African countries. <i>Pan African Medical Journal</i> , 2017, 26, 104.	0.8	20

#	ARTICLE	IF	CITATIONS
253	Correlates of leisure-time sedentary behavior among 181,793 adolescents aged 12-15 years from 66 low- and middle-income countries. <i>PLoS ONE</i> , 2019, 14, e0224339.	2.5	20
254	Informal Caregiving, Chronic Physical Conditions, and Physical Multimorbidity in 48 Low- and Middle-Income Countries. <i>Journals of Gerontology - Series A Biological Sciences and Medical Sciences</i> , 2020, 75, 1572-1578.	3.6	20
255	Physiotherapy for patients with anorexia nervosa. <i>Advances in Eating Disorders (Abingdon, England)</i> , 2013, 1, 224-238.	0.7	19
256	The Psychosocial Consequences of Sports Participation for Individuals with Severe Mental Illness: A Metasynthesis Review. <i>Advances in Psychiatry</i> , 2015, 2015, 1-8.	0.4	19
257	The functional exercise capacity in patients with bipolar disorder versus healthy controls: A pilot study. <i>Psychiatry Research</i> , 2015, 229, 194-199.	3.3	19
258	Are people with schizophrenia adherent to diabetes medication? A comparative meta-analysis. <i>Psychiatry Research</i> , 2017, 250, 17-24.	3.3	19
259	Physical activity and metabolic disease among people with affective disorders: Prevention, management and implementation. <i>Journal of Affective Disorders</i> , 2017, 224, 87-94.	4.1	19
260	Physical fitness and physical activity levels in people with alcohol use disorder versus matched healthy controls: A pilot study. <i>Alcohol</i> , 2019, 76, 73-79.	1.7	19
261	Physical activity and loneliness among adults aged 50 years or older in six low- and middle-income countries. <i>International Journal of Geriatric Psychiatry</i> , 2019, 34, 1855-1864.	2.7	19
262	Multimorbidity and obesity in older adults from six low- and middle-income countries. <i>Preventive Medicine</i> , 2021, 153, 106816.	3.4	19
263	Aerobic exercise capacity: an important correlate of psychosocial function in first episode psychosis. <i>Acta Psychiatrica Scandinavica</i> , 2015, 131, 234-234.	4.5	18
264	Depression and Reduced Bone Mineral Density at the Hip and Lumbar Spine. <i>Psychosomatic Medicine</i> , 2016, 78, 492-500.	2.0	18
265	Sitting time, physical fitness impairments and metabolic abnormalities in people with bipolar disorder: An exploratory study. <i>Psychiatry Research</i> , 2016, 242, 7-12.	3.3	18
266	Yoga versus non-standard care for schizophrenia. <i>The Cochrane Library</i> , 2017, 2017, CD012052.	2.8	18
267	Prevalence of suicidal ideation and attempt: associations with psychiatric disorders and HIV/AIDS in post-conflict Northern Uganda. <i>African Health Sciences</i> , 2017, 16, 1027.	0.7	18
268	The Impact of Pharmacologic and Nonpharmacologic Interventions to Improve Physical Health Outcomes in People With Dementia: A Meta-Review of Meta-Analyses of Randomized Controlled Trials. <i>Journal of the American Medical Directors Association</i> , 2020, 21, 1410-1414.e2.	2.5	18
269	Walking capacity is associated with health related quality of life and physical activity level in patients with schizophrenia: a preliminary report. <i>Actas Espanolas De Psiquiatria</i> , 2011, 39, 211-6.	0.1	18
270	Aerobic capacity is associated with global functioning in people with schizophrenia. <i>Journal of Mental Health</i> , 2015, 24, 214-218.	1.9	17

#	ARTICLE	IF	CITATIONS
271	Physical activity as a vital sign in patients with bipolar disorder. <i>Psychiatry Research</i> , 2016, 246, 218-222.	3.3	17
272	Risk of Hospitalized Falls and Hip Fractures in 22,103 Older Adults Receiving Mental Health Care vs 161,603 Controls: A Large Cohort Study. <i>Journal of the American Medical Directors Association</i> , 2020, 21, 1893-1899.	2.5	17
273	Cross-sectional associations of leisure and transport related physical activity with depression and anxiety. <i>Journal of Psychiatric Research</i> , 2021, 140, 228-234.	3.1	17
274	Addressing the disparity in physical health provision for people with schizophrenia: an important role for physiotherapists. <i>Physiotherapy</i> , 2014, 100, 185-186.	0.4	16
275	Associations Between Metabolic and Aerobic Fitness Parameters in Patients With Schizophrenia. <i>Journal of Nervous and Mental Disease</i> , 2015, 203, 23-27.	1.0	16
276	Among Inpatients, Posttraumatic Stress Disorder Symptom Severity Is Negatively Associated With Time Spent Walking. <i>Journal of Nervous and Mental Disease</i> , 2016, 204, 15-19.	1.0	16
277	Top 10 research questions to promote physical activity in bipolar disorders: A consensus statement from the International Organization of Physical Therapists in Mental Health. <i>Journal of Affective Disorders</i> , 2016, 195, 82-87.	4.1	16
278	Higher cardio-respiratory fitness is associated with increased mental and physical quality of life in people with bipolar disorder: A controlled pilot study. <i>Psychiatry Research</i> , 2017, 256, 219-224.	3.3	16
279	Physical activity and sleep problems in 38 low- and middle-income countries. <i>Sleep Medicine</i> , 2018, 48, 140-147.	1.6	16
280	Mental health informed physical activity for first responders and their support partner: a protocol for a stepped-wedge evaluation of an online, codesigned intervention. <i>BMJ Open</i> , 2019, 9, e030668.	1.9	16
281	Prospective associations between physical activity and clinician diagnosed major depressive disorder in adults: A 13-year cohort study. <i>Preventive Medicine</i> , 2019, 118, 38-43.	3.4	16
282	Exergames in people with major neurocognitive disorder: a systematic review. <i>Disability and Rehabilitation: Assistive Technology</i> , 2020, , 1-14.	2.2	16
283	Associations between expiratory spirometry parameters and limitations in daily life activities in patients with schizophrenia. <i>General Hospital Psychiatry</i> , 2014, 36, 172-176.	2.4	15
284	Concurrent validity of the international physical activity questionnaire in outpatients with bipolar disorder: Comparison with the Sensewear Armband. <i>Psychiatry Research</i> , 2016, 237, 122-126.	3.3	15
285	Correlates of physical activity among depressed older people in six low-income and middle-income countries: A community-based cross-sectional study. <i>International Journal of Geriatric Psychiatry</i> , 2018, 33, e314-e322.	2.7	15
286	Physical activity and HIV in sub-Saharan Africa: a systematic review of correlates and levels. <i>African Health Sciences</i> , 2018, 18, 394-406.	0.7	15
287	Different patterns of alcohol consumption and the incidence and persistence of depressive and anxiety symptoms among older adults in Ireland: A prospective community-based study. <i>Journal of Affective Disorders</i> , 2018, 238, 651-658.	4.1	15
288	Effects of acute exercise on craving, mood and anxiety in non-treatment seeking adults with alcohol use disorder: An exploratory study. <i>Drug and Alcohol Dependence</i> , 2021, 220, 108506.	3.2	15

#	ARTICLE	IF	CITATIONS
289	Yoga as part of a package of care versus standard care for schizophrenia. The Cochrane Library, 2018, 2018, CD012145.	2.8	15
290	The Assessment, Benefits and Delivery of Physical Activity in People with Schizophrenia: A Survey of Members of the International Organization of Physical Therapists in Mental Health. Physiotherapy Research International, 2014, 19, 248-256.	1.5	14
291	Physical activity correlates in people with anxiety: Data from 46 low- and middle-income countries. General Hospital Psychiatry, 2017, 49, 26-31.	2.4	14
292	Pain and severe sleep disturbance in the general population: Primary data and meta-analysis from 240,820 people across 45 low- and middle-income countries. General Hospital Psychiatry, 2018, 53, 52-58.	2.4	14
293	A Mental Healthâ€œInformed Physical Activity Intervention for First Responders and Their Partners Delivered Using Facebook: Mixed Methods Pilot Study. JMIR Formative Research, 2021, 5, e23432.	1.4	14
294	Exercise is medicine for depression: even when the “pill” is small. Neuropsychiatric Disease and Treatment, 2016, Volume 12, 2715-2721.	2.2	13
295	Depressive symptoms and muscular fitness contribute independently to the ability to perform daily life activities in people with bipolar disorder. Nordic Journal of Psychiatry, 2016, 70, 477-482.	1.3	13
296	Physical health policies and metabolic screening in mental health care systems of sub-Saharan African countries: a systematic review. International Journal of Mental Health Systems, 2017, 11, 31.	2.7	13
297	Physical activity correlates across the lifespan in people with epilepsy: a systematic review. Disability and Rehabilitation, 2021, 43, 1359-1366.	1.8	13
298	Bidirectional longitudinal relationship between leisure-time physical activity and psychotropic medication usage: A register linked follow-up study. Psychiatry Research, 2017, 247, 208-213.	3.3	12
299	Health care professionalsâ€™ perspectives on physical activity within the Ugandan mental health care system. Mental Health and Physical Activity, 2019, 16, 1-7.	1.8	12
300	Test-retest reliability and clinical correlates of the Eurofit test battery in people with alcohol use disorders. Psychiatry Research, 2019, 271, 208-213.	3.3	12
301	Exergaming for people with major neurocognitive disorder: a qualitative study. Disability and Rehabilitation, 2022, 44, 2044-2052.	1.8	12
302	Informal caregiving and physical activity among 204,315 adults in 38 low- and middle-income countries: A cross-sectional study. Preventive Medicine, 2020, 132, 106007.	3.4	12
303	The Relationship between Motor Skills and Intelligence in Children with Autism Spectrum Disorder. Journal of Autism and Developmental Disorders, 2022, 52, 1189-1199.	2.7	12
304	Endothelial dysfunction in people with depressive disorders: A systematic review and meta-analysis. Journal of Psychiatric Research, 2021, 141, 152-159.	3.1	12
305	Evaluation of the Developmental Coordination Questionnaire (DCDQ) as a Screening Instrument for Co-occurring Motor Problems in Children with Autism Spectrum Disorder. Journal of Autism and Developmental Disorders, 2022, 52, 4079-4088.	2.7	12
306	Dropout from physical activity interventions in children and adolescents with attention deficit hyperactivity disorder: A systematic review and meta-analysis. Mental Health and Physical Activity, 2016, 11, 46-52.	1.8	11

#	ARTICLE	IF	CITATIONS
307	Lifetime self-reported arthritis is associated with elevated levels of mental health burden: A multi-national cross sectional study across 46 low- and middle-income countries. <i>Scientific Reports</i> , 2017, 7, 7138.	3.3	11
308	Physical activity correlates in people with mild cognitive impairment: findings from six low- and middle-income countries. <i>Public Health</i> , 2018, 156, 15-25.	2.9	11
309	Adherence to physical activity recommendations and physical and mental health risk in people with severe mental illness in Uganda. <i>Psychiatry Research</i> , 2018, 260, 236-240.	3.3	11
310	The experience of mental health service users in health system strengthening: lessons from Uganda. <i>International Journal of Mental Health Systems</i> , 2019, 13, 60.	2.7	11
311	Associations of the Built Environment With Physical Activity and Sedentary Time in Ugandan Outpatients With Mental Health Problems. <i>Journal of Physical Activity and Health</i> , 2019, 16, 243-250.	2.0	11
312	Validity and correlates of the International Physical Activity Questionnaire in first-episode psychosis. <i>Microbial Biotechnology</i> , 2019, 13, 562-567.	1.7	11
313	The complex and dynamic interplay between self-esteem, belongingness and physical activity in daily life: An experience sampling study in adolescence and young adulthood. <i>Mental Health and Physical Activity</i> , 2021, 21, 100413.	1.8	11
314	Prospective associations between multiple lifestyle behaviors and depressive symptoms. <i>Journal of Affective Disorders</i> , 2022, 301, 233-239.	4.1	11
315	Metabolic syndrome and lung function in schizophrenia: A pilot study. <i>Psychiatry Research</i> , 2014, 220, 58-62.	3.3	10
316	Validity of the 6min walk test in outpatients with bipolar disorder. <i>Psychiatry Research</i> , 2015, 230, 664-667.	3.3	10
317	Lower cardiorespiratory fitness is associated with more time spent sedentary in first episode psychosis: A pilot study. <i>Psychiatry Research</i> , 2017, 253, 13-17.	3.3	10
318	Correlates of physical activity among community-dwelling individuals aged 65 years or older with anxiety in six low- and middle-income countries. <i>International Psychogeriatrics</i> , 2018, 30, 705-714.	1.0	10
319	Perceived barriers, benefits and correlates of physical activity in outpatients with Major Depressive Disorder: A study from Brazil. <i>Psychiatry Research</i> , 2020, 284, 112751.	3.3	10
320	The efficacy of a lay health workers led physical activity counselling program in patients with HIV and mental health problems: a real-world intervention from Uganda. <i>AIDS Care - Psychological and Socio-Medical Aspects of AIDS/HIV</i> , 2021, 33, 1189-1195.	1.2	10
321	Physical Activity, Sedentary Behavior, and Cardiorespiratory Fitness in Hazardous and Non-Hazardous Alcohol Consumers. <i>American Journal of Health Promotion</i> , 2021, 35, 669-678.	1.7	10
322	Motives for physical activity in the adoption and maintenance of physical activity in men with alcohol use disorders. <i>Psychiatry Research</i> , 2018, 261, 522-526.	3.3	9
323	A quantitative assessment of the views of mental health professionals on exercise for people with mental illness: perspectives from a low-resource setting. <i>African Health Sciences</i> , 2019, 19, 2172.	0.7	9
324	Physical activity and sedentary levels among people living with epilepsy: A systematic review and meta-analysis. <i>Epilepsy and Behavior</i> , 2019, 99, 106390.	1.7	9

#	ARTICLE	IF	CITATIONS
325	Metabolic syndrome and its components in people with intellectual disability: a meta-analysis. <i>Journal of Intellectual Disability Research</i> , 2020, 64, 804-815.	2.0	9
326	Food Insecurity With Hunger and Sexual Behavior Among Adolescents From 53 Countries. <i>International Journal of Sexual Health</i> , 2021, 33, 88-98.	2.3	9
327	Changes in craving following acute aerobic exercise in adults with alcohol use disorder. <i>Journal of Psychiatric Research</i> , 2021, 142, 243-249.	3.1	9
328	Independent and combined associations of sugar-sweetened beverage consumption, TV viewing, and physical activity with severe depressive symptoms among 59,402 adults. <i>Revista Brasileira De Psiquiatria</i> , 2021, 43, 574-583.	1.7	9
329	Correlates of sedentary behaviour among adults with hazardous drinking habits in six low- and middle-income countries. <i>Psychiatry Research</i> , 2018, 261, 406-413.	3.3	8
330	Physiotherapy for people with mental health problems in Sub-Saharan African countries: a systematic review. <i>Archives of Physiotherapy</i> , 2018, 8, 2.	1.8	8
331	Do reductions in ghrelin contribute towards antipsychotic-induced weight gain?. <i>Schizophrenia Research</i> , 2019, 210, 301-302.	2.0	8
332	Correlates of sedentary behavior in middle-aged and old age people with mild cognitive impairment: a multinational study. <i>International Psychogeriatrics</i> , 2019, 31, 579-589.	1.0	8
333	Test-retest reliability, validity, and correlates of the 2-min walk test in outpatients with depression. <i>Physiotherapy Research International</i> , 2020, 25, e1821.	1.5	8
334	Physical activity correlates in children and adolescents, adults, and older adults with an intellectual disability: a systematic review. <i>Disability and Rehabilitation</i> , 2022, 44, 4189-4200.	1.8	8
335	The VITAAL Stepping Exergame Prototype for Older Adults With Major Neurocognitive Disorder: A Usability Study. <i>Frontiers in Aging Neuroscience</i> , 2021, 13, 701319.	3.4	8
336	Physical activity participation, functional exercise capacity and self-esteem in patients with schizophrenia. <i>International Journal of Therapy and Rehabilitation</i> , 2011, 18, 222-229.	0.3	7
337	The Functional Exercise Capacity Is Associated With Global Functioning in People With Bipolar Disorder. <i>Journal of Nervous and Mental Disease</i> , 2016, 204, 673-677.	1.0	7
338	Physical activity correlates in heavy episodic drinkers: Data from 46 low- and middle-income countries. <i>Mental Health and Physical Activity</i> , 2017, 13, 163-170.	1.8	7
339	New steps for treating alcohol use disorder: the emerging importance of physical exercise. <i>Psychopharmacology</i> , 2018, 235, 2771-2773.	3.1	7
340	Cannabis use and physical activity among 89,777 adolescents aged 12-15 years from 21 low- and middle-income countries. <i>Drug and Alcohol Dependence</i> , 2019, 205, 107584.	3.2	7
341	Correlates of sedentary behavior among community-dwelling adults with anxiety in six low- and middle-income countries. <i>Psychiatry Research</i> , 2019, 273, 501-508.	3.3	7
342	Functional exercise capacity in inpatients with alcohol use disorder versus healthy controls: A pilot study. <i>Alcohol</i> , 2020, 82, 47-52.	1.7	7

#	ARTICLE	IF	CITATIONS
343	Functioning in schizophrenia: a Delphi study covering the perspective of physiotherapists. <i>Disability and Rehabilitation</i> , 2021, 43, 3733-3740.	1.8	7
344	The Impact of Pharmacological and Non-Pharmacological Interventions to Improve Physical Health Outcomes in People With Schizophrenia: A Meta-Review of Meta-Analyses of Randomized Controlled Trials. <i>Focus (American Psychiatric Publishing)</i> , 2021, 19, 116-128.	0.8	7
345	Considering the Role of Physical Therapists Within the Treatment and Rehabilitation of Individuals With Eating Disorders: An International Survey of Expert Clinicians. <i>Physiotherapy Research International</i> , 2016, 21, 237-246.	1.5	6
346	Interest, competence, appearance, fitness and social relatedness as motives for physical activity in Ugandan outpatients with psychosis. <i>Mental Health and Physical Activity</i> , 2017, 13, 94-99.	1.8	6
347	Psychotic experiences as an independent risk factor for angina pectoris in 48 low- and middle-income countries. <i>World Psychiatry</i> , 2018, 17, 232-234.	10.4	6
348	Physical fitness levels and moderators in people with epilepsy: A systematic review and meta-analysis. <i>Epilepsy and Behavior</i> , 2019, 99, 106448.	1.7	6
349	Associations between physical inactivity, major depressive disorder, and alcohol use disorder in people living with HIV in a Ugandan fishing community. <i>International Journal of STD and AIDS</i> , 2019, 30, 1177-1184.	1.1	6
350	Yoga as part of a package of care versus non-standard care for schizophrenia. <i>The Cochrane Library</i> , 2019, 2019, CD012807.	2.8	6
351	The efficacy of physical activity counseling in Ugandan patients with HIV and a co-morbid mental disorder: a pilot study. <i>AIDS Care - Psychological and Socio-Medical Aspects of AIDS/HIV</i> , 2020, 32, 758-761.	1.2	6
352	Lifestyle behaviors among 4,343 Brazilian adults with severe mental illness and 55,859 general population controls: data from the Brazilian National Health Survey. <i>Revista Brasileira De Psiquiatria</i> , 2020, 42, 245-249.	1.7	6
353	Moderate to severe generalized anxiety disorder symptoms are associated with physical inactivity in people with HIV/AIDS: a study from Uganda. <i>International Journal of STD and AIDS</i> , 2021, 32, 170-175.	1.1	6
354	Top 10 research questions to promote physical activity research in people with binge eating disorder. <i>Eating Disorders</i> , 2016, 24, 326-337.	3.0	5
355	Correlates of Physical Activity Among Middle-Aged and Older Adults With Hazardous Drinking Habits in Six Low- and Middle-Income Countries. <i>Journal of Aging and Physical Activity</i> , 2018, 26, 589-598.	1.0	5
356	Exercise self-efficacy correlates in people with psychosis. <i>Psychiatry Research</i> , 2018, 262, 359-362.	3.3	5
357	Test-retest reliability, concurrent validity and correlates of the two-minute walk test in outpatients with psychosis. <i>Psychiatry Research</i> , 2019, 282, 112619.	3.3	5
358	Motives for physical activity in the adoption and maintenance of physical activity in middle-aged and old age outpatients with a mental disorder: A cross-sectional study from a low-income country. <i>Psychiatry Research</i> , 2019, 282, 112620.	3.3	5
359	Functional exercise capacity is associated with global functioning in patients with alcohol use disorder. <i>Archives of Psychiatric Nursing</i> , 2019, 33, 144-148.	1.4	5
360	Risk of hospitalised falls and hip fractures in working age adults receiving mental health care. <i>General Hospital Psychiatry</i> , 2021, 72, 81-87.	2.4	5

#	ARTICLE	IF	CITATIONS
361	A 10-hours workshop improves physical activity prescription for mental illness knowledge and confidence in health care professionals: a nation-wide multicentre study from Uganda. <i>Disability and Rehabilitation</i> , 2021, , 1-6.	1.8	5
362	Mental contrasting and implementation of physical activity intentions in Ugandan primary care patients with mental health problems: A real-world intervention involving support partners. <i>Psychiatry Research</i> , 2022, 307, 114335.	3.3	5
363	Physical activity participation is associated with higher quality of life scores in men with alcohol use disorders: a study from Uganda. <i>African Health Sciences</i> , 2020, 20, 1407-1415.	0.7	5
364	Clinical correlates of global functioning in obese treatment seeking persons with binge eating disorder. <i>Psychiatria Danubina</i> , 2014, 26, 256-60.	0.4	5
365	Antipsychotic treatment, prolactin, and breast tumorigenesis. <i>Psychiatria Danubina</i> , 2016, 28, 243-254.	0.4	5
366	Mental health outcomes of the Daily Mile in elementary school children: a single-arm pilot study. <i>Child and Adolescent Mental Health</i> , 0, , .	3.5	5
367	No evidence of a control group response in exercise randomised controlled trials in people with schizophrenia: A systematic review and meta-analysis. <i>Psychiatry Research</i> , 2015, 229, 840-843.	3.3	4
368	Moving beyond the weight-loss paradigm of exercise interventions for mental illness. <i>Psychiatry Research</i> , 2016, 246, 392-393.	3.3	4
369	Gender differences in motives for physical activity across the stages of change in Ugandan outpatients with psychosis. <i>Schizophrenia Research</i> , 2018, 197, 568-569.	2.0	4
370	Cultural responses towards the aftermath of suicide among the Acholi in Northern Uganda. <i>International Journal of Social Psychiatry</i> , 2018, 64, 545-553.	3.1	4
371	Promoting physical activity in vulnerable adults at risk of homelessness: a randomised controlled trial protocol. <i>BMJ Open</i> , 2019, 9, e026466.	1.9	4
372	Is there an association between cardiorespiratory fitness and stage of illness in psychotic disorders? A systematic review and meta-analysis. <i>Acta Psychiatrica Scandinavica</i> , 2020, 141, 190-205.	4.5	4
373	Correlates of physical activity stages of change in people living with HIV in a Ugandan community. <i>Disability and Rehabilitation</i> , 2020, , 1-6.	1.8	4
374	Reliability and validity of physical fitness tests in people with mental disorders: A systematic review and meta-analysis. <i>Physiotherapy Research International</i> , 2021, 26, e1904.	1.5	4
375	Physical activity correlates in children and adolescents with autism spectrum disorder: a systematic review. <i>Disability and Rehabilitation</i> , 2022, 44, 6539-6550.	1.8	4
376	Objectively measured physical activity and symptoms of psychopathology in general population adolescents from the SIGMA cohort. <i>Mental Health and Physical Activity</i> , 2021, 21, 100416.	1.8	4
377	Association between physical activity and leisure-time sedentary behavior among 140,808 adolescents aged 12 to 15 from 47 low- and middle-income countries. <i>Public Health</i> , 2021, 199, 1-9.	2.9	4
378	Physical activity, physical fitness and quality of life in outpatients with major depressive disorder versus matched healthy controls: Data from a low-income country. <i>Journal of Affective Disorders</i> , 2021, 294, 802-804.	4.1	4

#	ARTICLE	IF	CITATIONS
379	Barriers, attitudes, confidence, and knowledge of nurses regarding metabolic health screening and intervention in people with mental illness: a pilot study from Uganda. <i>African Health Sciences</i> , 2019, 19, 2546-2554.	0.7	4
380	Internalised HIV-related stigma associated with physical inactivity in people with HIV and AIDS: A cross-sectional study from Uganda. <i>African Journal of AIDS Research</i> , 2021, 20, 238-243.	0.9	4
381	Prevalence of diabetes in people with intellectual disabilities and age- and gender-matched controls: A meta-analysis. <i>Journal of Applied Research in Intellectual Disabilities</i> , 2022, 35, 301-311.	2.0	4
382	Prospective Associations of Leisure-Time Physical Activity With Psychological Distress and Well-Being: A 12-Year Cohort Study. <i>Psychosomatic Medicine</i> , 2022, 84, 116-122.	2.0	4
383	Test-retest reliability and correlates of the Simple Physical Activity Questionnaire in Ugandan out-patients with psychosis. <i>African Health Sciences</i> , 2020, 20, 1438-1445.	0.7	4
384	The Metabolic Syndrome Is Associated with Self-Reported Physical Complaints in Patients with Bipolar Disorder. <i>Psychiatria Danubina</i> , 2016, 28, 139-45.	0.4	4
385	Exercise to spot the differences: a framework for the effect of exercise on hippocampal pattern separation in humans. <i>Reviews in the Neurosciences</i> , 2022, 33, 555-582.	2.9	4
386	Yoga vs Standard Care for Schizophrenia: Table 1.. <i>Schizophrenia Bulletin</i> , 2015, 42, sbv165.	4.3	3
387	Type 2 Diabetes Mellitus. , 2016, , 255-278.		3
388	Is Active Transport and Leisure-Time Physical Activity Associated With Inflammatory Markers in US Adults? A Cross-Sectional Analyses From NHANES. <i>Journal of Physical Activity and Health</i> , 2019, 16, 540-546.	2.0	3
389	Physical activity, physical fitness and quality of life in outpatients with a psychotic disorder versus healthy matched controls in a low-income country. <i>Schizophrenia Research</i> , 2021, 229, 1-2.	2.0	3
390	Response to Bartoli et al.. <i>American Journal of Psychiatry</i> , 2013, 170, 928-929.	7.2	2
391	Yoga as part of a package of care versus non-standard care for schizophrenia. <i>The Cochrane Library</i> , 2017, , .	2.8	2
392	Cannabis use and leisure-time sedentary behavior among 94,035 adolescents aged 12-15 years from 24 low- and middle-income countries. <i>Addictive Behaviors</i> , 2019, 99, 106104.	3.0	2
393	Which clinical and biochemical predictors should be used to screen for diabetes in patients with serious mental illness receiving antipsychotic medication? A large observational study. <i>PLoS ONE</i> , 2019, 14, e0210674.	2.5	2
394	Predictors of physical activity recording in routine mental healthcare. <i>Mental Health and Physical Activity</i> , 2020, 18, 100329.	1.8	2
395	Test-retest reliability, concurrent validity and correlates of the two-minute walk test in outpatients with alcohol use disorder. <i>Alcohol</i> , 2021, 90, 74-79.	1.7	2
396	Evaluating correlations between physical activity, psychological mediators of physical activity, and negative symptoms in individuals living with psychosis and diabetes.. <i>Psychiatric Rehabilitation Journal</i> , 2018, 41, 153-156.	1.1	2

#	ARTICLE	IF	CITATIONS
397	Test-retest study of the six-minute walk test in people with bipolar disorder. <i>Psychiatria Danubina</i> , 2016, 28, 39-44.	0.4	2
398	Predictors of falls and fractures leading to hospitalisation in 36â€™101 people with affective disorders: a large representative cohort study. <i>BMJ Open</i> , 2022, 12, e055070.	1.9	2
399	Exercise and PTSD Symptoms in Emergency Service and Frontline Medical Workers: A Systematic Review. <i>Translational Journal of the American College of Sports Medicine</i> , 2022, 7, .	0.6	2
400	A systematic evaluation and comparison of the guidelines for screening and monitoring of cardiometabolic risk in people with schizophrenia. <i>European Psychiatry</i> , 2011, 26, 2189-2189.	0.2	1
401	Yoga for schizophrenia. <i>The Cochrane Library</i> , 2018, , .	2.8	1
402	Bipolar Disorder and Physical Activity. , 2018, , 53-63.		1
403	Sibship size, birth order and psychotic experiences: Evidence from 43 low- and middle-income countries. <i>Schizophrenia Research</i> , 2018, 201, 406-412.	2.0	1
404	Sedentary Behavior and Quality of Life in People with Psychotic Disorders from a Low Income Country: A Study from Uganda. <i>Community Mental Health Journal</i> , 2019, 55, 714-720.	2.0	1
405	Testâ€™retest reliability and correlates of the 6â€™min walk test in people with alcohol use disorders. <i>Physiotherapy Research International</i> , 2021, 26, e1900.	1.5	1
406	Influence of weekday of admission and level of distress on length of hospital stay in patients with low back pain: a retrospective cohort study. <i>BMC Musculoskeletal Disorders</i> , 2021, 22, 656.	1.9	1
407	Exercise and Posttraumatic Stress Disorder. , 2018, , 375-387.		1
408	Test-retest reliability and correlates of the Simple Physical Activity Questionnaire in Ugandan out-patients with psychosis. <i>African Health Sciences</i> , 2020, 20, 1438-1445.	0.7	1
409	Multimorbidity among Latinx-Americans and Asian American/Pacific Islanders. <i>Journal of Ageing and Longevity</i> , 2022, 2, 26-33.	0.6	1
410	Schizophrenia and Exercise. , 2018, , 65-82.		0
411	Research and Evaluation in Exercise and Mental Health. , 2018, , 301-317.		0
412	Reliability and Convergent Validity of Self-Reported Physical Activity Questionnaires for People With Mental Disorders: A Systematic Review and Meta-Analysis. <i>Journal of Physical Activity and Health</i> , 2021, 18, 109-115.	2.0	0
413	Correlates of missed HIV appointments in low-resource settings: a study from Uganda. <i>African Journal of AIDS Research</i> , 2021, 20, 125-131.	0.9	0
414	Does being overweight play a role in the reduced inhibitory control of patients receiving treatment for substance use disorder?. <i>Physiology and Behavior</i> , 2021, 241, 113587.	2.1	0

#	ARTICLE	IF	CITATIONS
415	Exercise and Posttraumatic Stress Disorder. , 2017, , 1-13.		0
416	Physical activity participation is associated with higher quality of life scores in men with alcohol use disorders: a study from Uganda. African Health Sciences, 2020, 20, 1407-1415.	0.7	0
417	Osteoporosis referral and treatment among people with severe mental illness: A ten-year data linkage study. Journal of Psychiatric Research, 2022, 147, 94-102.	3.1	0
418	Less time spent walking and depressive symptoms are associated with more self-perceived disability in primary care patients: a cross-sectional study from Uganda. Pan African Medical Journal, 2022, 41, 173.	0.8	0