

# Hiroshi Maegawa

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4603396/publications.pdf>

Version: 2024-02-01

329  
papers

12,051  
citations

25034

57  
h-index

42399

92  
g-index

342  
all docs

342  
docs citations

342  
times ranked

16767  
citing authors

#	ARTICLE	IF	CITATIONS
1	A case of central diabetes insipidus due to neurophysin II gene abnormality diagnosed based on a family history of nocturnal enuresis. <i>Endocrine Journal</i> , 2022, 69, 95-100.	1.6	0
2	Patient characteristics associated with improvement in glycemic control following addition of an oral antidiabetic drug to DPP-4 inhibitor monotherapy in Japanese patients with type 2 diabetes mellitus (JDDM 60). <i>Diabetology International</i> , 2022, 13, 132-141.	1.4	0
3	Glycemic control and number of natural teeth: analysis of cross-sectional Japanese employment-based dental insurance claims and medical check-up data. <i>Diabetology International</i> , 2022, 13, 244-252.	1.4	8
4	Relationship between Kidney Function and Subclinical Atherosclerosis Progression Evaluated by Coronary Artery Calcification. <i>Journal of Atherosclerosis and Thrombosis</i> , 2022, 29, 1359-1371.	2.0	9
5	Higher Iron Intake Is Independently Associated with Obesity in Younger Japanese Type-2 Diabetes Mellitus Patients. <i>Nutrients</i> , 2022, 14, 211.	4.1	1
6	Clinical course of different long-acting insulin therapies—glargine U100, U300, degludec, and insulin degludec/insulin aspart—among Japanese patients with type 2 diabetes: a multicenter retrospective observational study (JDDM65 study). <i>Endocrine Journal</i> , 2022, , .	1.6	0
7	A new era of diabetic kidney disease treatment with sodium—glucose cotransporter-2 inhibitors. <i>Journal of Diabetes Investigation</i> , 2022, 13, 765-767.	2.4	4
8	Improvement in Estimated Glomerular Filtration Rate Decline Rate after Febuxostat Treatment in a Fabry Disease Patient with Enzyme Replacement Therapy-resistant Proteinuria. <i>Internal Medicine</i> , 2022, , .	0.7	2
9	Eighteen-year trends in the management of patients with diabetes in the Shiga Diabetes Clinical Survey: overall trends and differences by age group. <i>Diabetology International</i> , 2022, 13, 566-574.	1.4	1
10	Metabolic changes induced by dapagliflozin, an SGLT2 inhibitor, in Japanese patients with type 2 diabetes treated by oral anti-diabetic agents: A randomized, clinical trial. <i>Diabetes Research and Clinical Practice</i> , 2022, 186, 109781.	2.8	15
11	Inhibition of mitochondrial fission protects podocytes from albumin-induced cell damage in diabetic kidney disease. <i>Biochimica Et Biophysica Acta - Molecular Basis of Disease</i> , 2022, 1868, 166368.	3.8	11
12	Glycaemia and body weight are regulated by sodium-glucose cotransporter 1 (SGLT1) expression via O-GlcNAcylation in the intestine. <i>Molecular Metabolism</i> , 2022, 59, 101458.	6.5	8
13	Differential Association of Serum n-3 Polyunsaturated Fatty Acids with Various Cerebrovascular Lesions in Japanese Men. <i>Cerebrovascular Diseases</i> , 2022, 51, 774-780.	1.7	0
14	Trends in glycemic control in patients with insulin therapy compared with non-insulin or no drugs in type 2 diabetes in Japan: a long-term view of real-world treatment between 2002 and 2018 (JDDM 66). <i>BMJ Open Diabetes Research and Care</i> , 2022, 10, e002727.	2.8	3
15	Ketone body 3-hydroxybutyrate enhances adipocyte function. <i>Scientific Reports</i> , 2022, 12, .	3.3	8
16	A Long-term Estimated Glomerular Filtration Rate Plot Analysis Permits the Accurate Assessment of a Decline in the Renal Function by Minimizing the Influence of Estimated Glomerular Filtration Rate Fluctuations. <i>Internal Medicine</i> , 2022, 61, 1823-1833.	0.7	0
17	Limited effects of systemic or renal lipoprotein lipase deficiency on renal physiology and diseases. <i>Biochemical and Biophysical Research Communications</i> , 2022, 620, 15-20.	2.1	2
18	Sodium—glucose cotransporter-2 inhibitors represent a paradigm shift in the prevention of heart failure in type-2 diabetes patients. <i>Journal of Diabetes Investigation</i> , 2021, 12, 6-20.	2.4	17

#	ARTICLE	IF	CITATIONS
19	Role of O-linked N-acetylglucosamine in the homeostasis of metabolic organs, and its potential links with diabetes and its complications. <i>Journal of Diabetes Investigation</i> , 2021, 12, 130-136.	2.4	10
20	Geometry of Sleeve Gastrectomy Measured by 3D CT Versus Weight Loss: Preliminary Analysis. <i>World Journal of Surgery</i> , 2021, 45, 235-242.	1.6	5
21	Real-World Evidence for Long-Term Safety and Effectiveness of Ipragliflozin in Japanese Patients with Type 2 Diabetes Mellitus: final Results of a 3-Year Post-Marketing Surveillance Study (STELLA-LONG TERM). <i>Diabetologia</i> , 2021, 64, 1431-1437.	10.784	11
22	Glucagon-Like Peptide-1 Receptor Agonist Utilization in Type 2 Diabetes in Japan: A Retrospective Database Analysis (JDDM 57). <i>Diabetes Therapy</i> , 2021, 12, 345-361.	2.5	12
23	Safety and effectiveness of ipragliflozin in Japanese patients with type 2 diabetes mellitus and impaired renal function: subgroup analysis of a 3-year post-marketing surveillance study (STELLA-LONG TERM). <i>Diabetologia</i> , 2021, 64, 181-196.	1.4	3
24	Clinical inertia in patients with type 2 diabetes treated with oral antidiabetic drugs: Results from a Japanese cohort study (JDDM53). <i>Journal of Diabetes Investigation</i> , 2021, 12, 374-381.	2.4	15
25	Effect of ipragliflozin on liver function in Japanese type 2 diabetes mellitus patients: subgroup analysis of a 3-year post-marketing surveillance study (STELLA-LONG TERM). <i>Endocrine Journal</i> , 2021, 68, 905-918.	1.6	2
26	Machine Learning Approach to Decision Making for Insulin Initiation in Japanese Patients With Type 2 Diabetes (JDDM 58): Model Development and Validation Study. <i>JMIR Medical Informatics</i> , 2021, 9, e22148.	2.6	7
27	MicroRNA-494-3p inhibits formation of fast oxidative muscle fibres by targeting E1A-binding protein p300 in human-induced pluripotent stem cells. <i>Scientific Reports</i> , 2021, 11, 1161.	3.3	2
28	Alcohol drinking and brain morphometry in apparently healthy community-dwelling Japanese men. <i>Alcohol</i> , 2021, 90, 57-65.	1.7	6
29	Genome-wide association studies identify two novel loci conferring susceptibility to diabetic retinopathy in Japanese patients with type 2 diabetes. <i>Human Molecular Genetics</i> , 2021, 30, 716-726.	2.9	13
30	Roles of mTOR in Diabetic Kidney Disease. <i>Antioxidants</i> , 2021, 10, 321.	5.1	21
31	Targeted deletion of Atg5 in intestinal epithelial cells promotes dextran sodium sulfate-induced colitis. <i>Journal of Clinical Biochemistry and Nutrition</i> , 2021, 68, 156-163.	1.4	4
32	Safety and Effectiveness of Ipragliflozin in Elderly Versus Non-elderly Japanese Patients with Type 2 Diabetes: Subgroup Analysis of STELLA-LONG TERM. <i>Diabetes Therapy</i> , 2021, 12, 1359-1378.	2.5	0
33	Real-world evidence for long-term safety and effectiveness of ipragliflozin in treatment-naïve versus non-naïve Japanese patients with type 2 diabetes mellitus: subgroup analysis of a 3-year post-marketing surveillance study (STELLA-LONG TERM). <i>Diabetologia</i> , 2021, 64, 430-444.	1.4	3
34	Combination of disease duration at diagnosis and hemoglobin A1c-to-serum C-peptide reactivity ratios predicts patient response to glucose-lowering medication in type 2 diabetes: A retrospective cohort study across Japan (JDDM59). <i>Journal of Diabetes Investigation</i> , 2021, 12, 1967-1977.	2.4	1
35	Malfunctioning CD106-positive, short-term hematopoietic stem cells trigger diabetic neuropathy in mice by cell fusion. <i>Communications Biology</i> , 2021, 4, 575.	4.4	6
36	Current status of oral antidiabetic drug prescribing patterns based on the body mass index for Japanese type 2 diabetes mellitus patients and yearly changes in diabetologists' prescribing patterns from 2002 to 2019 (JDDM61). <i>Journal of Diabetes Investigation</i> , 2021, , .	2.4	14

#	ARTICLE	IF	CITATIONS
37	Comparing Associations of Dietary Energy Density and Energy Intake, Macronutrients with Obesity in Patients with Type 2 Diabetes (JDDM 63). <i>Nutrients</i> , 2021, 13, 3167.	4.1	5
38	Cardio- and reno-protective effects of dipeptidyl peptidase III in diabetic mice. <i>Journal of Biological Chemistry</i> , 2021, 296, 100761.	3.4	12
39	Nutrition and Periodontal Health in the Patients with Diabetes Mellitus: a Review from the Viewpoint of Endothelial Function. <i>Current Oral Health Reports</i> , 2021, 8, 67-74.	1.6	2
40	Liver fat accumulation assessed by computed tomography is an independent risk factor for diabetes mellitus in a population-based study: SESSA (Shiga Epidemiological Study of Subclinical) Tj ETQq0 0 0 rgBT /Overlook 2010 Tf 50 617 Td (A	2.8	10
41	Impact of obesity on underreporting of energy intake in type 2 diabetic patients: Clinical Evaluation of Energy Requirements in Patients with Diabetes Mellitus (CLEVER-DM) study. <i>Clinical Nutrition ESPEN</i> , 2020, 39, 251-254.	1.2	5
42	SGLT2 Inhibition Mediates Protection from Diabetic Kidney Disease by Promoting Ketone Body-Induced mTORC1 Inhibition. <i>Cell Metabolism</i> , 2020, 32, 404-419.e6.	16.2	197
43	Family Support for Medical Nutritional Therapy and Dietary Intake among Japanese with Type 2 Diabetes (JDDM 56). <i>Nutrients</i> , 2020, 12, 2649.	4.1	4
44	Validity of the Use of a Triaxial Accelerometer and a Physical Activity Questionnaire for Estimating Total Energy Expenditure and Physical Activity Level among Elderly Patients with Type 2 Diabetes Mellitus: CLEVER-DM Study. <i>Annals of Nutrition and Metabolism</i> , 2020, 76, 62-72.	1.9	10
45	Protective role of podocyte autophagy against glomerular endothelial dysfunction in diabetes. <i>Biochemical and Biophysical Research Communications</i> , 2020, 525, 319-325.	2.1	17
46	Contrast medium-induced severe thrombocytopenia in patient on maintenance hemodialysis: a case report and literature review. <i>CEN Case Reports</i> , 2020, 9, 266-270.	0.9	1
47	The Prognosis of Patients With Type 2 Diabetes and Nonalbuminuric Diabetic Kidney Disease Is Not Always Poor: Implication of the Effects of Coexisting Macrovascular Complications (JDDM 54). <i>Diabetes Care</i> , 2020, 43, 1102-1110.	8.6	40
48	A Real-World Observational Study Evaluating the Probability of Glycemic Control with Basal Insulin or Glucagon-Like Peptide-1 Receptor Agonist in Japanese Patients with Type 2 Diabetes. <i>Diabetes Therapy</i> , 2020, 11, 1481-1496.	2.5	11
49	Lipotoxicity, Nutrient-Sensing Signals, and Autophagy in Diabetic Nephropathy. <i>JMA Journal</i> , 2020, 3, 87-94.	0.8	10
50	Association of blood levels of marine omega-3 fatty acids with coronary calcification and calcium density in Japanese men. <i>European Journal of Clinical Nutrition</i> , 2019, 73, 783-792.	2.9	22
51	Carotid Intima-Media Thickness and Plaque in Apparently Healthy Japanese Individuals with an Estimated 10-Year Absolute Risk of CAD Death According to the Japan Atherosclerosis Society (JAS) Guidelines 2012: The Shiga Epidemiological Study of Subclinical Atherosclerosis (SESSA). <i>Journal of Atherosclerosis and Thrombosis</i> , 2019, 26, 746-746.	2.0	1
52	Safety and effectiveness of ipragliflozin in elderly versus non-elderly Japanese type 2 diabetes mellitus patients: 12 month interim results of the STELLA-LONG TERM study. <i>Current Medical Research and Opinion</i> , 2019, 35, 1901-1910.	1.9	6
53	FO053ROLE OF KETONE BODY METABOLISM IN SGLT2 INHIBITOR-MEDIATED RENOPROTECTION IN HIGH FAT DIET-FED APOE-KNOCKOUT MICE. <i>Nephrology Dialysis Transplantation</i> , 2019, 34, .	0.7	0
54	SP430Potential role for ketone body metabolism in an SGLT2 inhibitor-mediated anti-albuminuric effect in type 2 diabetic db/db mice. <i>Nephrology Dialysis Transplantation</i> , 2019, 34, .	0.7	0

#	ARTICLE	IF	CITATIONS
55	Efficacy of metformin on postprandial plasma triglyceride concentration by administration timing in patients with type 2 diabetes mellitus: A randomized crossover pilot study. <i>Journal of Diabetes Investigation</i> , 2019, 10, 1284-1290.	2.4	6
56	Advanced Technology for Gene Delivery with Homing Peptides to Spinal Cord through Systemic Circulation in Mice. <i>Molecular Therapy - Methods and Clinical Development</i> , 2019, 13, 474-483.	4.1	1
57	Preserving $\beta$ -cell function is the major determinant of diabetes remission following laparoscopic sleeve gastrectomy in Japanese obese diabetic patients. <i>Endocrine Journal</i> , 2019, 66, 817-826.	1.6	6
58	Identification of subgroups of patients with type 2 diabetes with differences in renal function preservation, comparing patients receiving sodium-glucose cotransporter 2 inhibitors with those receiving dipeptidyl peptidase 4 inhibitors, using a supervised machine learning algorithm (PROFILE) <i>Tj ETQq0 04rgBT /Owlock 10 Metabolism</i> , 2019, 21, 1925-1934.		
59	Protein O-GlcNAcylation Is Essential for the Maintenance of Renal Energy Homeostasis and Function via Lipolysis during Fasting and Diabetes. <i>Journal of the American Society of Nephrology: JASN</i> , 2019, 30, 962-978.	6.1	18
60	Efficacy and safety of pemafibrate in people with type 2 diabetes and elevated triglyceride levels: 52-week data from the PROVIDE study. <i>Diabetes, Obesity and Metabolism</i> , 2019, 21, 1737-1744.	4.4	35
61	Microbiome potentiates endurance exercise through intestinal acetate production. <i>American Journal of Physiology - Endocrinology and Metabolism</i> , 2019, 316, E956-E966.	3.5	131
62	Combined Effects of Energy Intake and Physical Activity on Obesity in Japanese Patients with Type 2 Diabetes (JDDM 50): A Cross-Sectional Study. <i>Diabetes Therapy</i> , 2019, 10, 1133-1138.	2.5	1
63	Diabetes management and treatment approaches outside of North America and West Europe in 2006 and 2015. <i>Acta Diabetologica</i> , 2019, 56, 889-897.	2.5	4
64	Impact of body mass index on the efficacy and safety of ipragliflozin in Japanese patients with type 2 diabetes mellitus: A subgroup analysis of 3-month interim results from the Specified Drug Use Results Survey of Ipragliflozin Treatment in Type 2 Diabetic Patients: Long-term Use study. <i>Journal of Diabetes Investigation</i> , 2019, 10, 1262-1271.	2.4	5
65	Safety and Effectiveness of Ipragliflozin for Type 2 Diabetes in Japan: 12-Month Interim Results of the STELLA-LONG TERM Post-Marketing Surveillance Study. <i>Advances in Therapy</i> , 2019, 36, 923-949.	2.9	15
66	Monkeys mutant for PKD1 recapitulate human autosomal dominant polycystic kidney disease. <i>Nature Communications</i> , 2019, 10, 5517.	12.8	33
67	Ipragliflozin, a sodium-glucose cotransporter 2 inhibitor, reduces bodyweight and fat mass, but not muscle mass, in Japanese type 2 diabetes patients treated with insulin: A randomized clinical trial. <i>Journal of Diabetes Investigation</i> , 2019, 10, 1012-1021.	2.4	41
68	Effect of ipragliflozin on liver function in Japanese type 2 diabetes mellitus patients: a subgroup analysis of the STELLA-LONG TERM study (3-month interim results). <i>Endocrine Journal</i> , 2019, 66, 31-41.	1.6	19
69	Secular changes in clinical manifestations of kidney disease among Japanese adults with type 2 diabetes from 1996 to 2014. <i>Journal of Diabetes Investigation</i> , 2019, 10, 1032-1040.	2.4	39
70	A role for bone marrow-derived cells in diabetic nephropathy. <i>FASEB Journal</i> , 2019, 33, 4067-4076.	0.5	10
71	Total energy expenditure is comparable between patients with and without diabetes mellitus: Clinical Evaluation of Energy Requirements in Patients with Diabetes Mellitus (CLEVER-DM) Study. <i>BMJ Open Diabetes Research and Care</i> , 2019, 7, e000648.	2.8	19
72	Gene Therapy for Neuropathic Pain through siRNA-IRF5 Gene Delivery with Homing Peptides to Microglia. <i>Molecular Therapy - Nucleic Acids</i> , 2018, 11, 203-215.	5.1	36

#	ARTICLE	IF	CITATIONS
73	Safety and efficacy of ipragliflozin in elderly versus non-elderly Japanese patients with type 2 diabetes mellitus: a subgroup analysis of the STELLA-LONG TERM study. <i>Expert Opinion on Pharmacotherapy</i> , 2018, 19, 327-336.	1.8	10
74	Lack of O-GlcNAcylation enhances exercise-dependent glucose utilization potentially through AMP-activated protein kinase activation in skeletal muscle. <i>Biochemical and Biophysical Research Communications</i> , 2018, 495, 2098-2104.	2.1	18
75	Effects of Pemafibrate, a Novel Selective PPAR $\alpha$ Modulator, on Lipid and Glucose Metabolism in Patients With Type 2 Diabetes and Hypertriglyceridemia: A Randomized, Double-Blind, Placebo-Controlled, Phase 3 Trial. <i>Diabetes Care</i> , 2018, 41, 538-546.	8.6	122
76	International Comparison of Abdominal Fat Distribution Among Four Populations: The ERA-JUMP Study. <i>Metabolic Syndrome and Related Disorders</i> , 2018, 16, 166-173.	1.3	9
77	Current status of achieving blood pressure target and its clinical correlates in Japanese type 2 diabetes (JDDM45). <i>Journal of Diabetes Investigation</i> , 2018, 9, 594-601.	2.4	5
78	Overexpression of acetyl CoA carboxylase $\beta$ exacerbates podocyte injury in the kidney of streptozotocin-induced diabetic mice. <i>Biochemical and Biophysical Research Communications</i> , 2018, 495, 1115-1121.	2.1	3
79	Safety and efficacy of ipragliflozin in Japanese patients with type 2 diabetes in real-world clinical practice: interim results of the STELLA-LONG TERM post-marketing surveillance study. <i>Expert Opinion on Pharmacotherapy</i> , 2018, 19, 189-201.	1.8	29
80	Impact of obesity on annual medical expenditures and diabetes care in Japanese patients with type 2 diabetes mellitus. <i>Journal of Diabetes Investigation</i> , 2018, 9, 776-781.	2.4	10
81	A variant within the FTO confers susceptibility to diabetic nephropathy in Japanese patients with type 2 diabetes. <i>PLoS ONE</i> , 2018, 13, e0208654.	2.5	30
82	Clinical inertia in basal insulin-treated patients with type 2 diabetes – Results from a retrospective database study in Japan (JDDM 43). <i>PLoS ONE</i> , 2018, 13, e0198160.	2.5	15
83	MiR-494-3p regulates mitochondrial biogenesis and thermogenesis through PGC1- $\beta$ signalling in beige adipocytes. <i>Scientific Reports</i> , 2018, 8, 15096.	3.3	71
84	Definitive diagnosis of mandibular hypoplasia, deafness, progeroid features and lipodystrophy (MDPL) syndrome caused by a recurrent <i>de novo</i> mutation in the <i>POLD1</i> gene. <i>Endocrine Journal</i> , 2018, 65, 227-238.	1.6	42
85	Role of dietary amino acid balance in diet restriction-mediated lifespan extension, renoprotection, and muscle weakness in aged mice. <i>Aging Cell</i> , 2018, 17, e12796.	6.7	45
86	Change in Pericardial Fat Volume and Cardiovascular Risk Factors in a General Population of Japanese Men. <i>Circulation Journal</i> , 2018, 82, 2542-2548.	1.6	11
87	Improved glucose metabolism by Eragrostis tef potentially through beige adipocyte formation and attenuating adipose tissue inflammation. <i>PLoS ONE</i> , 2018, 13, e0201661.	2.5	6
88	Twelve-year trends of increasing overweight and obesity in patients with diabetes: the Shiga Diabetes Clinical Survey. <i>Endocrine Journal</i> , 2018, 65, 527-536.	1.6	21
89	Declining trends of diabetic nephropathy, retinopathy and neuropathy with improving diabetes care indicators in Japanese patients with type 2 and type 1 diabetes (JDDM 46). <i>BMJ Open Diabetes Research and Care</i> , 2018, 6, e000521.	2.8	42
90	MicroRNA-494 plays a role in fiber type-specific skeletal myogenesis by targeting transcriptional coactivator p300 in human induced pluripotent stem cells. <i>Proceedings for Annual Meeting of the Japanese Pharmacological Society</i> , 2018, WCP2018, OR19-3.	0.0	0



#	ARTICLE	IF	CITATIONS
91	A case of bacterial peritonitis complicated by eosinophilic peritonitis in the peritoneal dialysis induction period. <i>Nihon Toseki Igakkai Zasshi</i> , 2018, 51, 463-467.	0.1	0
92	Statin use and all-cause and cancer mortality: BioBank Japan cohort. <i>Journal of Epidemiology</i> , 2017, 27, S84-S91.	2.4	25
93	Metabolic and hemodynamic effects of sodium-dependent glucose cotransporter 2 inhibitors on cardio-renal protection in the treatment of patients with type 2 diabetes mellitus. <i>Journal of Diabetes Investigation</i> , 2017, 8, 416-427.	2.4	59
94	Pivotal Role of $\alpha$ -GlcNAc Modification in Cold-Induced Thermogenesis by Brown Adipose Tissue Through Mitochondrial Biogenesis. <i>Diabetes</i> , 2017, 66, 2351-2362.	0.6	28
95	Cholesterol levels of Japanese dyslipidaemic patients with various comorbidities: BioBank Japan. <i>Journal of Epidemiology</i> , 2017, 27, S77-S83.	2.4	3
96	O-linked $\beta$ -N-acetylglucosamine modification of proteins is essential for foot process maturation and survival in podocytes. <i>Nephrology Dialysis Transplantation</i> , 2017, 32, 1477-1487.	0.7	23
97	Effect of an intensified multifactorial intervention on cardiovascular outcomes and mortality in type 2 diabetes (J-DOIT3): an open-label, randomised controlled trial. <i>Lancet Diabetes and Endocrinology</i> , 2017, 5, 951-964.	11.4	228
98	Diverse metabolic effects of O-GlcNAcylation in the pancreas but limited effects in insulin-sensitive organs in mice. <i>Diabetologia</i> , 2017, 60, 1761-1769.	6.3	25
99	Serum glucose, cholesterol and blood pressure levels in Japanese type 1 and 2 diabetic patients: BioBank Japan. <i>Journal of Epidemiology</i> , 2017, 27, S92-S97.	2.4	12
100	Trends in medical performance in diabetic patients in primary care clinics compared with those in hospitals: Shiga Diabetes Clinical Survey, Japan, 2000-2012. <i>Diabetology International</i> , 2017, 8, 59-68.	1.4	6
101	Comparison of baseline characteristics and clinical course in Japanese patients with type 2 diabetes among whom different types of oral hypoglycemic agents were chosen by diabetes specialists as initial monotherapy (JDDM 42). <i>Medicine (United States)</i> , 2017, 96, e6122.	1.0	21
102	N-3 Polyunsaturated Fatty Acids Decrease the Protein Expression of Soluble Epoxide Hydrolase via Oxidative Stress-Induced P38 Kinase in Rat Endothelial Cells. <i>Nutrients</i> , 2017, 9, 654.	4.1	10
103	Acute Effect of Metformin on Postprandial Hypertriglyceridemia through Delayed Gastric Emptying. <i>International Journal of Molecular Sciences</i> , 2017, 18, 1282.	4.1	17
104	Fiber-rich diet with brown rice improves endothelial function in type 2 diabetes mellitus: A randomized controlled trial. <i>PLoS ONE</i> , 2017, 12, e0179869.	2.5	52
105	Amla Enhances Mitochondrial Spare Respiratory Capacity by Increasing Mitochondrial Biogenesis and Antioxidant Systems in a Murine Skeletal Muscle Cell Line. <i>Oxidative Medicine and Cellular Longevity</i> , 2016, 2016, 1-11.	4.0	49
106	Stearoyl-CoA Desaturase-1 Protects Cells against Lipotoxicity-Mediated Apoptosis in Proximal Tubular Cells. <i>International Journal of Molecular Sciences</i> , 2016, 17, 1868.	4.1	41
107	Evaluation of a Novel Glucose Area Under the Curve (AUC) Monitoring System: Comparison with the AUC by Continuous Glucose Monitoring. <i>Diabetes and Metabolism Journal</i> , 2016, 40, 326.	4.7	7
108	Concentrations of Water-Soluble Vitamins in Blood and Urinary Excretion in Patients with Diabetes Mellitus. <i>Nutrition and Metabolic Insights</i> , 2016, 9, NMI.S40595.	1.9	17

#	ARTICLE	IF	CITATIONS
109	Comparative Effects of Direct Renin Inhibitor and Angiotensin Receptor Blocker on Albuminuria in Hypertensive Patients with Type 2 Diabetes. A Randomized Controlled Trial. PLoS ONE, 2016, 11, e0164936.	2.5	11
110	Mammalian autophagy is essential for hepatic and renal ketogenesis during starvation. Scientific Reports, 2016, 6, 18944.	3.3	58
111	Data set for renal sinus fat volume and visceral adipose tissue volume on computed tomography. Data in Brief, 2016, 7, 1658-1664.	1.0	3
112	Association between symptoms of bilateral numbness and/or paresthesia in the feet and postural instability in Japanese patients with diabetes. Diabetology International, 2016, 7, 69-76.	1.4	7
113	Association between attentional function and postural instability in Japanese older patients with diabetes mellitus. Diabetology International, 2016, 7, 83-88.	1.4	1
114	Renal sinus fat volume on computed tomography in middle-aged patients at risk for cardiovascular disease and its association with coronary artery calcification. Atherosclerosis, 2016, 246, 374-381.	0.8	12
115	Emerging role of mammalian autophagy in ketogenesis to overcome starvation. Autophagy, 2016, 12, 709-710.	9.1	24
116	MicroRNA148b-3p inhibits mTORC1-dependent apoptosis in diabetes by repressing TNFR2 in proximal tubular cells. Kidney International, 2016, 90, 1211-1225.	5.2	27
117	Smoking, Smoking Cessation, and Measures of Subclinical Atherosclerosis in Multiple Vascular Beds in Japanese Men. Journal of the American Heart Association, 2016, 5, .	3.7	39
118	Lifetime cigarette smoking is associated with abdominal obesity in a community-based sample of Japanese men: The Shiga Epidemiological Study of Subclinical Atherosclerosis (SESSA). Preventive Medicine Reports, 2016, 4, 225-232.	1.8	30
119	Baseline characteristics and interim (3-month) efficacy and safety data from STELLA-LONG TERM, a long-term post-marketing surveillance study of ipragliflozin in Japanese patients with type 2 diabetes in real-world clinical practice. Expert Opinion on Pharmacotherapy, 2016, 17, 1985-1994.	1.8	21
120	Hypothalamic AMP-Activated Protein Kinase Regulates Biphasic Insulin Secretion from Pancreatic $\beta^2$ Cells during Fasting and in Type 2 Diabetes. EBioMedicine, 2016, 13, 168-180.	6.1	14
121	Large-scale survey of rates of achieving targets for blood glucose, blood pressure, and lipids and prevalence of complications in type 2 diabetes (JDDM 40). BMJ Open Diabetes Research and Care, 2016, 4, e000294.	2.8	67
122	Smoking status is associated with mild cognitive impairment assessed with the mini-mental state examination in Japanese diabetic patients. Diabetology International, 2016, 7, 361-367.	1.4	7
123	Relationship of Insulin Resistance to Prevalence and Progression of Coronary Artery Calcification Beyond Metabolic Syndrome Components. Arteriosclerosis, Thrombosis, and Vascular Biology, 2016, 36, 1703-1708.	2.4	44
124	Renoprotective effect of DPP-4 inhibitors against free fatty acid-bound albumin-induced renal proximal tubular cell injury. Biochemical and Biophysical Research Communications, 2016, 470, 539-545.	2.1	37
125	Genome-wide association studies in the Japanese population identify seven novel loci for type 2 diabetes. Nature Communications, 2016, 7, 10531.	12.8	149
126	Lipoprotein-associated phospholipase A2 is related to risk of subclinical atherosclerosis but is not supported by Mendelian randomization analysis in a general Japanese population. Atherosclerosis, 2016, 246, 141-147.	0.8	48



#	ARTICLE	IF	CITATIONS
127	A case of local delayed-type allergy to zinc-containing insulin as a cause of diabetic ketoacidosis in a patient with type 1 diabetes mellitus undergoing continuous subcutaneous insulin infusion. <i>Diabetology International</i> , 2016, 7, 447-450.	1.4	3
128	Associations of serum LDL particle concentration with carotid intima-media thickness and coronary artery calcification. <i>Journal of Clinical Lipidology</i> , 2016, 10, 1195-1202.e1.	1.5	12
129	Impaired Podocyte Autophagy Exacerbates Proteinuria in Diabetic Nephropathy. <i>Diabetes</i> , 2016, 65, 755-767.	0.6	243
130	Replication Study in a Japanese Population of Six Susceptibility Loci for Type 2 Diabetes Originally Identified by a Transethnic Meta-Analysis of Genome-Wide Association Studies. <i>PLoS ONE</i> , 2016, 11, e0154093.	2.5	10
131	Hyperglycemia Induces Skin Barrier Dysfunctions with Impairment of Epidermal Integrity in Non-Wounded Skin of Type 1 Diabetic Mice. <i>PLoS ONE</i> , 2016, 11, e0166215.	2.5	47
132	Focal Segmental Glomerular Sclerosis Ameliorated by Long-term Hemodialysis Therapy with Low-density Lipoprotein Apheresis. <i>Internal Medicine</i> , 2015, 54, 2213-2217.	0.7	7
133	Duality of n-3 Polyunsaturated Fatty Acids on Mcp-1 Expression in Vascular Smooth Muscle: A Potential Role of 4-Hydroxy Hexenal. <i>Nutrients</i> , 2015, 7, 8112-8126.	4.1	7
134	Predictors for Mild and Severe Hypoglycemia in Insulin-Treated Japanese Diabetic Patients. <i>PLoS ONE</i> , 2015, 10, e0130584.	2.5	8
135	Epidermis-dermis junction as a novel location for bone marrow-derived cells to reside in response to ionizing radiation. <i>Biochemical and Biophysical Research Communications</i> , 2015, 461, 695-701.	2.1	3
136	Emerging role of podocyte autophagy in the progression of diabetic nephropathy. <i>Autophagy</i> , 2015, 11, 2385-2386.	9.1	87
137	1-Methylnicotinamide ameliorates lipotoxicity-induced oxidative stress and cell death in kidney proximal tubular cells. <i>Free Radical Biology and Medicine</i> , 2015, 89, 831-841.	2.9	41
138	High-density lipoprotein particle concentration and subclinical atherosclerosis of the carotid arteries in Japanese men. <i>Atherosclerosis</i> , 2015, 239, 444-450.	0.8	18
139	Comparison of HOMA-IR, HOMA- $\beta$ % and disposition index between US white men and Japanese men in Japan: the ERA JUMP study. <i>Diabetologia</i> , 2015, 58, 265-271.	6.3	39
140	Use of MRI signal intensity of extraocular muscles to evaluate methylprednisolone pulse therapy in thyroid-associated ophthalmopathy. <i>Japanese Journal of Ophthalmology</i> , 2015, 59, 124-130.	1.9	28
141	Population Pharmacokinetics and Therapeutic Efficacy of Febuxostat in Patients with Severe Renal Impairment. <i>Pharmacology</i> , 2015, 96, 90-98.	2.2	32
142	Replication study of the association of rs7578597 in THADA, rs10886471 in GRK5, and rs7403531 in RASGRP1 with susceptibility to type 2 diabetes among a Japanese population. <i>Diabetology International</i> , 2015, 6, 306-312.	1.4	1
143	Multiple osteoporotic fractures in a patient with CKD stage G3b. <i>Kidney International</i> , 2015, 87, 1077.	5.2	0
144	MicroRNA-494 plays a role in fiber type-specific skeletal myogenesis in human induced pluripotent stem cells. <i>Biochemical and Biophysical Research Communications</i> , 2015, 468, 208-213.	2.1	10

#	ARTICLE	IF	CITATIONS
145	Lamp-2 deficiency prevents high-fat diet-induced obese diabetes via enhancing energy expenditure. Biochemical and Biophysical Research Communications, 2015, 465, 249-255.	2.1	18
146	Urinary Potassium Excretion and Renal and Cardiovascular Complications in Patients with Type 2 Diabetes and Normal Renal Function. Clinical Journal of the American Society of Nephrology: CJASN, 2015, 10, 2152-2158.	4.5	68
147	Replication Study in a Japanese Population to Evaluate the Association between 10 SNP Loci, Identified in European Genome-Wide Association Studies, and Type 2 Diabetes. PLoS ONE, 2015, 10, e0126363.	2.5	14
148	Effect of nutritional counseling and long term isomaltulose based liquid formula (MHN-01) intake on metabolic syndrome. Journal of Clinical Biochemistry and Nutrition, 2015, 57, 140-144.	1.4	8
149	Predictive Properties of Plasma Amino Acid Profile for Cardiovascular Disease in Patients with Type 2 Diabetes. PLoS ONE, 2014, 9, e101219.	2.5	41
150	Long chain n-3 polyunsaturated fatty acids and incidence rate of coronary artery calcification in Japanese men in Japan and white men in the USA: population based prospective cohort study. Heart, 2014, 100, 569-573.	2.9	47
151	Role of Nutrient-Sensing Signals in the Pathogenesis of Diabetic Nephropathy. BioMed Research International, 2014, 2014, 1-9.	1.9	51
152	CCDC3 is specifically upregulated in omental adipose tissue in subjects with abdominal obesity. Obesity, 2014, 22, 1070-1077.	3.0	14
153	Stem cell factor-activated bone marrow ameliorates amyotrophic lateral sclerosis by promoting protective microglial migration. Journal of Neuroscience Research, 2014, 92, 856-869.	2.9	13
154	Oral glucose-stimulated serum C-peptide predicts successful switching from insulin therapy to liraglutide monotherapy in Japanese patients with type 2 diabetes and renal impairment. Journal of Diabetes Investigation, 2014, 5, 435-441.	2.4	7
155	Dapagliflozin as Monotherapy or Combination Therapy in Japanese Patients with Type 2 Diabetes: an Open-Label Study. Diabetes Therapy, 2014, 5, 415-433.	2.5	56
156	A fish-based diet intervention improves endothelial function in postmenopausal women with type 2 diabetes mellitus: A randomized crossover trial. Metabolism: Clinical and Experimental, 2014, 63, 930-940.	3.4	43
157	Hyperglycemia induces abnormal gene expression in hematopoietic stem cells and their progeny in diabetic neuropathy. FEBS Letters, 2014, 588, 1080-1086.	2.8	22
158	Autophagy: Emerging Therapeutic Target for Diabetic Nephropathy. Seminars in Nephrology, 2014, 34, 9-16.	1.6	56
159	Genome-wide association study identifies three novel loci for type 2 diabetes. Human Molecular Genetics, 2014, 23, 239-246.	2.9	158
160	Safety and efficacy of skin patches containing loxoprofen sodium in diabetic patients with overt nephropathy. Clinical and Experimental Nephrology, 2014, 18, 487-491.	1.6	6
161	4-Hydroxy hexenal derived from dietary n-3 polyunsaturated fatty acids induces anti-oxidative enzyme heme oxygenase-1 in multiple organs. Biochemical and Biophysical Research Communications, 2014, 443, 991-996.	2.1	35
162	Glycated Albumin Predicts the Risk of Mortality in Type 2 Diabetic Patients on Hemodialysis: Evaluation of a Target Level for Improving Survival. Therapeutic Apheresis and Dialysis, 2014, 18, 434-442.	0.9	33

#	ARTICLE	IF	CITATIONS
163	A high-fiber, low-fat diet improves periodontal disease markers in high-risk subjects: a pilot study. <i>Nutrition Research</i> , 2014, 34, 491-498.	2.9	59
164	A Mutation of COX6A1 Causes a Recessive Axonal or Mixed Form of Charcot-Marie-Tooth Disease. <i>American Journal of Human Genetics</i> , 2014, 95, 294-300.	6.2	65
165	Fatty acids are novel nutrient factors to regulate mTORC1 lysosomal localization and apoptosis in podocytes. <i>Biochimica Et Biophysica Acta - Molecular Basis of Disease</i> , 2014, 1842, 1097-1108.	3.8	99
166	Gene Therapy for Neuropathic Pain by Silencing of TNF- $\alpha$ Expression with Lentiviral Vectors Targeting the Dorsal Root Ganglion in Mice. <i>PLoS ONE</i> , 2014, 9, e92073.	2.5	54
167	Assessing the Clinical Utility of a Genetic Risk Score Constructed Using 49 Susceptibility Alleles for Type 2 Diabetes in a Japanese Population. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2013, 98, E1667-E1673.	3.6	37
168	Replication study for the association of 3 SNP loci identified in a genome-wide association study for diabetic nephropathy in European type 1 diabetes with diabetic nephropathy in Japanese patients with type 2 diabetes. <i>Clinical and Experimental Nephrology</i> , 2013, 17, 866-871.	1.6	16
169	Effect of sleeve gastrectomy on body weight, food intake, glucose tolerance, and metabolic hormone level in two different rat models: Goto-Kakizaki and diet-induced obese rat. <i>Journal of Surgical Research</i> , 2013, 185, 159-165.	1.6	12
170	Altered Unfolded Protein Response Is Implicated in the Age-Related Exacerbation of Proteinuria-Induced Proximal Tubular Cell Damage. <i>American Journal of Pathology</i> , 2013, 183, 774-785.	3.8	21
171	LOX-1 ligands containing apolipoprotein B and carotid intima-media thickness in middle-aged community-dwelling US Caucasian and Japanese men. <i>Atherosclerosis</i> , 2013, 229, 240-245.	0.8	21
172	Ethnic difference in liver fat content: A cross-sectional observation among Japanese American in Hawaii, Japanese in Japan, and non-Hispanic whites in United States. <i>Obesity Research and Clinical Practice</i> , 2013, 7, e198-e205.	1.8	7
173	Ketogenic essential amino acids replacement diet ameliorated hepatosteatosis with altering autophagy-associated molecules. <i>Biochimica Et Biophysica Acta - Molecular Basis of Disease</i> , 2013, 1832, 1605-1612.	3.8	28
174	Anti-aging molecule, Sirt1: a novel therapeutic target for diabetic nephropathy. <i>Archives of Pharmacal Research</i> , 2013, 36, 230-236.	6.3	60
175	Genome-Wide Association Study Identifies a Novel Locus Contributing to Type 2 Diabetes Susceptibility in Sikhs of Punjabi Origin From India. <i>Diabetes</i> , 2013, 62, 1746-1755.	0.6	167
176	Haematopoietic cells produce BDNF and regulate appetite upon migration to the hypothalamus. <i>Nature Communications</i> , 2013, 4, 1526.	12.8	32
177	Omega-3 polyunsaturated fatty acid has an anti-oxidant effect via the Nrf-2/HO-1 pathway in 3T3-L1 adipocytes. <i>Biochemical and Biophysical Research Communications</i> , 2013, 430, 225-230.	2.1	81
178	A cross-sectional association of obesity with coronary calcium among Japanese, Koreans, Japanese Americans, and US Whites. <i>European Heart Journal Cardiovascular Imaging</i> , 2013, 14, 921-927.	1.2	21
179	Cilostazol Attenuates Spontaneous Microaggregation of Platelets in Type 2 Diabetic Patients With Insufficient Platelet Response to Aspirin. <i>Diabetes Care</i> , 2013, 36, e92-e93.	8.6	3
180	The Role of Autophagy in the Pathogenesis of Diabetic Nephropathy. <i>Journal of Diabetes Research</i> , 2013, 2013, 1-9.	2.3	64

#	ARTICLE	IF	CITATIONS
181	Predictive Effects of Urinary Liver-Type Fatty Acidâ€‘Binding Protein for Deteriorating Renal Function and Incidence of Cardiovascular Disease in Type 2 Diabetic Patients Without Advanced Nephropathy. <i>Diabetes Care</i> , 2013, 36, 1248-1253.	8.6	86
182	Evaluation of a Minimally Invasive System for Measuring Glucose Area under the Curve during Oral Glucose Tolerance Tests: Usefulness of Sweat Monitoring for Precise Measurement. <i>Journal of Diabetes Science and Technology</i> , 2013, 7, 678-688.	2.2	40
183	Influence of cigarette smoking on coronary artery and aortic calcium among random samples from populations of middle-aged Japanese and Korean men. <i>Journal of Epidemiology and Community Health</i> , 2013, 67, 119-124.	3.7	24
184	Obesity-Mediated Autophagy Insufficiency Exacerbates Proteinuria-induced Tubulointerstitial Lesions. <i>Journal of the American Society of Nephrology: JASN</i> , 2013, 24, 1769-1781.	6.1	185
185	Usefulness of a novel system for measuring glucose area under the curve while screening for glucose intolerance in outpatients. <i>Journal of Diabetes Investigation</i> , 2013, 4, 552-559.	2.4	17
186	Replication study for the association of rs391300 in <i>SRR</i> and rs17584499 in <i>PTPRD</i> with susceptibility to type 2 diabetes in a Japanese population. <i>Journal of Diabetes Investigation</i> , 2013, 4, 168-173.	2.4	11
187	Ezetimibe prevents hepatic steatosis induced by a high-fat but not a high-fructose diet. <i>American Journal of Physiology - Endocrinology and Metabolism</i> , 2013, 305, E293-E304.	3.5	30
188	Replication study for the association of a single-nucleotide polymorphism, rs3746876, within <i>KCNJ15</i> , with susceptibility to type 2 diabetes in a Japanese population. <i>Journal of Human Genetics</i> , 2013, 58, 490-493.	2.3	8
189	Association of myeloperoxidase G-463A gene polymorphism with diabetic nephropathy in Japanese type 2 diabetic subjects. <i>Endocrine Journal</i> , 2013, 60, 457-471.	1.6	7
190	Octreotide improves early dumping syndrome potentially through incretins: a case report. <i>Endocrine Journal</i> , 2013, 60, 847-853.	1.6	13
191	Enhanced Intestinal Motility during Oral Glucose Tolerance Test after Laparoscopic Sleeve Gastrectomy: Preliminary Results Using Cine Magnetic Resonance Imaging. <i>PLoS ONE</i> , 2013, 8, e65739.	2.5	30
192	Replication Study for the Association of 9 East Asian GWAS-Derived Loci with Susceptibility to Type 2 Diabetes in a Japanese Population. <i>PLoS ONE</i> , 2013, 8, e76317.	2.5	43
193	Carotid Intima-Media Thickness and Plaque in Apparently Healthy Japanese Individuals with an Estimated 10-Year Absolute Risk of CAD Death According to the Japan Atherosclerosis Society (JAS) Guidelines 2012: The Shiga Epidemiological Study of Subclinical Atherosclerosis (SESSA). <i>Journal of Atherosclerosis and Thrombosis</i> , 2013, 20, 755-766.	2.0	43
194	The Influence of a Single Nucleotide Polymorphism within <i>CNDP1</i> on Susceptibility to Diabetic Nephropathy in Japanese Women with Type 2 Diabetes. <i>PLoS ONE</i> , 2013, 8, e54064.	2.5	23
195	4-Hydroxy Hexenal Derived from Docosahexaenoic Acid Protects Endothelial Cells via Nrf2 Activation. <i>PLoS ONE</i> , 2013, 8, e69415.	2.5	69
196	Laparoscopic Sleeve Gastrectomy and Dipeptidyl Peptidase-4 Inhibitor which Led to the Improvement of Long-Term Continued Type 2 Diabetes Mellitus. <i>Japanese Journal of Gastroenterological Surgery</i> , 2013, 46, 334-341.	0.1	1
197	High Sodium Intake Is Associated With Masked Hypertension in Japanese Patients With Type 2 Diabetes and Treated Hypertension. <i>American Journal of Hypertension</i> , 2012, 25, 1170-1174.	2.0	18
198	Regulation of Mitochondrial Biogenesis by Lipoprotein Lipase in Muscle of Insulin-Resistant Offspring of Parents With Type 2 Diabetes. <i>Diabetes</i> , 2012, 61, 877-887.	0.6	63

#	ARTICLE	IF	CITATIONS
199	A single-nucleotide polymorphism in ANK1 is associated with susceptibility to type 2 diabetes in Japanese populations. <i>Human Molecular Genetics</i> , 2012, 21, 3042-3049.	2.9	99
200	Efficacy and tolerability of vildagliptin in type 2 diabetic patients on hemodialysis. <i>Journal of Diabetes Investigation</i> , 2012, 3, 298-301.	2.4	12
201	Association between urinary angiotensinogen levels and renal and cardiovascular prognoses in patients with type 2 diabetes mellitus. <i>Journal of Diabetes Investigation</i> , 2012, 3, 318-324.	2.4	41
202	Autophagy as a Therapeutic Target in Diabetic Nephropathy. <i>Experimental Diabetes Research</i> , 2012, 2012, 1-12.	3.8	92
203	Arterial stiffness and renal impairment in nonproteinuric type 2 diabetic patients. <i>Journal of Diabetes Investigation</i> , 2012, 3, 86-91.	2.4	3
204	Autophagy: a novel therapeutic target for kidney diseases. <i>Clinical and Experimental Nephrology</i> , 2012, 16, 827-832.	1.6	18
205	The effects of blood pressure control levels on the renoprotection of type 2 diabetic patients without overt proteinuria. <i>Journal of the American Society of Hypertension</i> , 2012, 6, 124-131.	2.3	11
206	Serum levels of marine-derived n-3 fatty acids in Icelanders, Japanese, Koreans, and Americans—a descriptive epidemiologic study. <i>Prostaglandins Leukotrienes and Essential Fatty Acids</i> , 2012, 87, 11-16.	2.2	31
207	Autophagy regulates inflammation in adipocytes. <i>Biochemical and Biophysical Research Communications</i> , 2012, 417, 352-357.	2.1	91
208	Role of angiotensin II-mediated AMPK inactivation on obesity-related salt-sensitive hypertension. <i>Biochemical and Biophysical Research Communications</i> , 2012, 418, 559-564.	2.1	37
209	Fructose induces tubulointerstitial injury in the kidney of mice. <i>Biochemical and Biophysical Research Communications</i> , 2012, 419, 244-249.	2.1	27
210	MicroRNA-494 regulates mitochondrial biogenesis in skeletal muscle through mitochondrial transcription factor A and Forkhead box j3. <i>American Journal of Physiology - Endocrinology and Metabolism</i> , 2012, 303, E1419-E1427.	3.5	119
211	Factors associated with progression of diabetic nephropathy in Japanese elderly patients with type 2 diabetes: Subanalysis of the Japanese Elderly Diabetes Intervention Trial. <i>Geriatrics and Gerontology International</i> , 2012, 12, 127-133.	1.5	13
212	A Single Nucleotide Polymorphism within DUSP9 Is Associated with Susceptibility to Type 2 Diabetes in a Japanese Population. <i>PLoS ONE</i> , 2012, 7, e46263.	2.5	38
213	Fenofibrate, a PPAR $\alpha$ agonist, has renoprotective effects in mice by enhancing renal lipolysis. <i>Kidney International</i> , 2011, 79, 871-882.	5.2	145
214	Association between serum soluble TNF $\alpha$ receptors and renal dysfunction in type 2 diabetic patients without proteinuria. <i>Diabetes Research and Clinical Practice</i> , 2011, 92, 174-180.	2.8	25
215	Association of New Loci Identified in European Genome-Wide Association Studies with Susceptibility to Type 2 Diabetes in the Japanese. <i>PLoS ONE</i> , 2011, 6, e26911.	2.5	62
216	SIRT3 attenuates palmitate-induced ROS production and inflammation in proximal tubular cells. <i>Free Radical Biology and Medicine</i> , 2011, 51, 1258-1267.	2.9	121

#	ARTICLE	IF	CITATIONS
217	Elevated serum levels of interleukin-18 in patients with overt diabetic nephropathy: effects of miglitol. <i>Clinical and Experimental Nephrology</i> , 2011, 15, 58-63.	1.6	13
218	Association between single nucleotide polymorphisms within genes encoding sirtuin families and diabetic nephropathy in Japanese subjects with type 2 diabetes. <i>Clinical and Experimental Nephrology</i> , 2011, 15, 381-390.	1.6	63
219	The role of sleep disturbance and depression in patients with type 2 diabetes. <i>Diabetology International</i> , 2011, 2, 79-85.	1.4	3
220	Postprandial activation of protein kinase C $\alpha$ 1/2 regulates the expression of adipocytokines via the transcription factor AP-2 $\beta$ . <i>International Journal of Molecular Medicine</i> , 2011, 28, 95-100.	4.0	6
221	Furosemide-associated nephrocalcinosis and renal cysts. <i>Kidney International</i> , 2011, 80, 556.	5.2	5
222	Inactivation of TNF- $\alpha$ ameliorates diabetic neuropathy in mice. <i>American Journal of Physiology - Endocrinology and Metabolism</i> , 2011, 301, E844-E852.	3.5	109
223	CW501516, a PPAR $\gamma$ Agonist, Ameliorates Tubulointerstitial Inflammation in Proteinuric Kidney Disease via Inhibition of TAK1-NF $\kappa$ B Pathway in Mice. <i>PLoS ONE</i> , 2011, 6, e25271.	2.5	31
224	Effects of hyperglycemia on oxidative stress and antioxidant potential in patients with type 2 diabetes. <i>Diabetology International</i> , 2010, 1, 72-77.	1.4	8
225	Laparoscopic sleeve gastrectomy leads to rapid improvement of glucose tolerance and insulin secretion with enhanced glucagon-like peptide (GLP-1) secretion. <i>Diabetology International</i> , 2010, 1, 99-103.	1.4	7
226	Presence and functional role of the rapidly activating delayed rectifier K $^{+}$ current in left and right atria of adult mice. <i>European Journal of Pharmacology</i> , 2010, 649, 14-22.	3.5	20
227	Relation of the Expression of Transcriptional Factor <i>TFAP2B</i> to That of Adipokines in Subcutaneous and Omental Adipose Tissues. <i>Obesity</i> , 2010, 18, 1277-1282.	3.0	14
228	A genome-wide association study in the Japanese population identifies susceptibility loci for type 2 diabetes at UBE2E2 and C2CD4A-C2CD4B. <i>Nature Genetics</i> , 2010, 42, 864-868.	21.4	245
229	Effects of a Fish-Based Diet on the Serum Adiponectin Concentration in Young, Non-Obese, Healthy Japanese Subjects. <i>Journal of Atherosclerosis and Thrombosis</i> , 2010, 17, 628-637.	2.0	39
230	Association Between Urinary Type IV Collagen Level and Deterioration of Renal Function in Type 2 Diabetic Patients Without Overt Proteinuria. <i>Diabetes Care</i> , 2010, 33, 1805-1810.	8.6	62
231	Circulating Levels of 8 Cytokines and Marine n-3 Fatty Acids and Indices of Obesity in Japanese, White, and Japanese American Middle-Aged Men. <i>Journal of Interferon and Cytokine Research</i> , 2010, 30, 541-548.	1.2	19
232	Transcription factor AP-2 $\beta$ : A negative regulator of IRS-1 gene expression. <i>Biochemical and Biophysical Research Communications</i> , 2010, 392, 526-532.	2.1	24
233	Low concentration of 4-hydroxy hexenal increases heme oxygenase-1 expression through activation of Nrf2 and antioxidative activity in vascular endothelial cells. <i>Biochemical and Biophysical Research Communications</i> , 2010, 402, 99-104.	2.1	65
234	Oleate and eicosapentaenoic acid attenuate palmitate-induced inflammation and apoptosis in renal proximal tubular cell. <i>Biochemical and Biophysical Research Communications</i> , 2010, 402, 265-271.	2.1	72



#	ARTICLE	IF	CITATIONS
235	Effects of blood pressure and the renin-angiotensin system on platelet activation in type 2 diabetes. <i>Journal of Diabetes Investigation</i> , 2010, 1, 196-201.	2.4	1
236	Membrane Localization of Protein-Tyrosine Phosphatase 1B is Essential for its Activation of Sterol Regulatory Element-Binding Protein-1 Gene Expression and Consequent Hypertriglyceridaemia. <i>Journal of Biochemistry</i> , 2009, 146, 541-547.	1.7	12
237	Correlation Between Albuminuria and Spontaneous Platelet Microaggregate Formation in Type 2 Diabetic Patients. <i>Diabetes Care</i> , 2009, 32, 2062-2067.	8.6	14
238	Transcription Factor Activating Protein-2 $\beta$ : A Positive Regulator of Monocyte Chemoattractant Protein-1 Gene Expression. <i>Endocrinology</i> , 2009, 150, 1654-1661.	2.8	14
239	Cholesteryl Ester Transfer Protein, Coronary Calcium, and Intima-Media Thickness of the Carotid Artery in Middle-Age Japanese Men. <i>American Journal of Cardiology</i> , 2009, 104, 818-822.	1.6	19
240	Higher arterial stiffness, greater peripheral vascular resistance and lower blood flow in lower leg arteries are associated with long-term hyperglycaemia in type 2 diabetic patients with normal ankle-brachial index. <i>Diabetes/Metabolism Research and Reviews</i> , 2009, 25, 363-369.	4.0	15
241	Construction of a prediction model for type 2 diabetes mellitus in the Japanese population based on 11 genes with strong evidence of the association. <i>Journal of Human Genetics</i> , 2009, 54, 236-241.	2.3	70
242	Soy phosphatidylcholine inhibited TLR4-mediated MCP-1 expression in vascular cells. <i>Atherosclerosis</i> , 2009, 205, 404-412.	0.8	45
243	Visceral and Subcutaneous Adiposity and Adiponectin in Middle-aged Japanese Men: The ERA JUMP Study. <i>Obesity</i> , 2009, 17, 1269-1273.	3.0	31
244	The Onset of Diabetes in Three out of Four Sisters: A Japanese Family with Type 1 Diabetes. A Case Report. <i>Endocrine Journal</i> , 2009, 56, 767-772.	1.6	5
245	SAFB1, an RBMX-binding protein, is a newly identified regulator of hepatic SREBP-1c gene. <i>BMB Reports</i> , 2009, 42, 232-237.	2.4	16
246	Association of TCF7L2 polymorphisms with susceptibility to type 2 diabetes in 4,087 Japanese subjects. <i>Journal of Human Genetics</i> , 2008, 53, 174-180.	2.3	80
247	Postprandial hyperglycemia after a gastrectomy and the prediabetic state: A comparison between a distal and total gastrectomy. <i>Surgery Today</i> , 2008, 38, 685-690.	1.5	6
248	Variants in KCNQ1 are associated with susceptibility to type 2 diabetes mellitus. <i>Nature Genetics</i> , 2008, 40, 1092-1097.	21.4	694
249	Intima-media thickness of the carotid artery and the distribution of lipoprotein subclasses in men aged 40 to 49 years between whites in the United States and the Japanese in Japan for the ERA JUMP study. <i>Metabolism: Clinical and Experimental</i> , 2008, 57, 177-182.	3.4	20
250	Renal segmental hypoplasia, Ask-Upmark kidney, in a patient with microalbuminuric hypertensive type 2 diabetes mellitus. <i>Diabetes Research and Clinical Practice</i> , 2008, 80, e22-e24.	2.8	1
251	Impaired peripheral circulation in lower-leg arteries caused by higher arterial stiffness and greater vascular resistance associates with nephropathy in type 2 diabetic patients with normal ankle-brachial indices. <i>Diabetes Research and Clinical Practice</i> , 2008, 80, 416-423.	2.8	16
252	Replication of Genome-Wide Association Studies of Type 2 Diabetes Susceptibility in Japan. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2008, 93, 3136-3141.	3.6	130

#	ARTICLE	IF	CITATIONS
253	Impact of Renin-Angiotensin System Inhibition on Microalbuminuria in Type 2 Diabetes: A Post Hoc Analysis of the Shiga Microalbuminuria Reduction Trial (SMART). Hypertension Research, 2008, 31, 1171-1176.	2.7	9
254	Reduction of Microalbuminuria in Patients With Type 2 Diabetes. Diabetes Care, 2007, 30, 1581-1583.	8.6	56
255	Less Subclinical Atherosclerosis in Japanese Men in Japan than in White Men in the United States in the Post-World War II Birth Cohort. American Journal of Epidemiology, 2007, 165, 617-624.	3.4	132
256	Short-term Low-dosage Pioglitazone Treatment Improves Vascular Dysfunction in Patients with Type 2 Diabetes. Endocrine Journal, 2007, 54, 613-618.	1.6	15
257	Massive pericardial effusion secondary to Hashimoto's disease. European Journal of Internal Medicine, 2007, 18, 438-440.	2.2	13
258	RBMX is a novel hepatic transcriptional regulator of SREBP-1c gene response to high-fructose diet. FEBS Letters, 2007, 581, 218-222.	2.8	22
259	Higher levels of adiponectin in American than in Japanese men despite obesity. Metabolism: Clinical and Experimental, 2006, 55, 1561-1563.	3.4	28
260	MafA differentiates rat intestinal cells into insulin-producing cells. Biochemical and Biophysical Research Communications, 2006, 349, 136-143.	2.1	28
261	Enhanced sodium sensitivity and disturbed circadian rhythm of blood pressure in essential hypertension. Journal of Hypertension, 2006, 24, 1627-1632.	0.5	113
262	Alcohol Consumption and Coronary Artery Calcium in Middle-Aged Japanese Men. American Journal of Cardiology, 2006, 98, 141-144.	1.6	21
263	Intronic Polymorphisms within TFAP2B Regulate Transcriptional Activity and Affect Adipocytokine Gene Expression in Differentiated Adipocytes. Molecular Endocrinology, 2006, 20, 1104-1111.	3.7	48
264	The Transcription Factor AP-2 $\hat{1}$ Causes Cell Enlargement and Insulin Resistance in 3T3-L1 Adipocytes. Endocrinology, 2006, 147, 1685-1696.	2.8	38
265	Transcription Factor Activating Enhancer-binding Protein-2 $\hat{1}$ . Journal of Biological Chemistry, 2006, 281, 31245-31253.	3.4	37
266	Low Blood Flow Estimates in Lower-Leg Arteries Predict Cardiovascular Events in Japanese Patients With Type 2 Diabetes With Normal Ankle-Brachial Indexes. Diabetes Care, 2006, 29, 1884-1890.	8.6	12
267	Transcription Factor Activating Enhancer-binding Protein-2 $\hat{1}$ . Journal of Biological Chemistry, 2006, 281, 31245-31253.	3.4	7
268	Genetic variations in the gene encoding TFAP2B are associated with type 2 diabetes mellitus. Journal of Human Genetics, 2005, 50, 283-292.	2.3	68
269	Bidirectional regulation of monocyte chemoattractant protein-1 gene at distinct sites of its promoter by nitric oxide in vascular smooth muscle cells. American Journal of Physiology - Cell Physiology, 2005, 289, C582-C590.	4.6	13
270	Regulation of ATP-sensitive Potassium Channel Subunit Kir6.2 Expression in Rat Intestinal Insulin-producing Progenitor Cells. Journal of Biological Chemistry, 2005, 280, 1893-1900.	3.4	17

#	ARTICLE	IF	CITATIONS
271	Diet high in lipid hydroperoxide by vitamin E deficiency induces insulin resistance and impaired insulin secretion in normal rats. <i>Diabetes Research and Clinical Practice</i> , 2005, 67, 99-109.	2.8	17
272	Abnormal peripheral circulation in type 2 diabetic patients with normal ankle-brachial index associates with coronary atherosclerosis, large artery stiffness, and peripheral vascular resistance. <i>Diabetes Research and Clinical Practice</i> , 2005, 70, 253-262.	2.8	29
273	Single Nucleotide Polymorphism (â€“468 Gly to Ala) at the Promoter Region of Sterol Regulatory Element-binding Protein-1c Associates with Genetic Defect of Fructose-induced Hepatic Lipogenesis. <i>Journal of Biological Chemistry</i> , 2004, 279, 29031-29042.	3.4	55
274	Much lower prevalence of coronary calcium detected by electron-beam computed tomography among men aged 40-49 in Japan than in the US, despite a less favorable profile of major risk factors. <i>International Journal of Epidemiology</i> , 2004, 34, 173-179.	1.9	47
275	Protein Phosphatase 2A Negatively Regulates Insulin's Metabolic Signaling Pathway by Inhibiting Akt (Protein Kinase B) Activity in 3T3-L1 Adipocytes. <i>Molecular and Cellular Biology</i> , 2004, 24, 8778-8789.	2.3	199
276	Protein Phosphatase-2C $\pm$ as a Positive Regulator of Insulin Sensitivity through Direct Activation of Phosphatidylinositol 3-Kinase in 3T3-L1 Adipocytes. <i>Journal of Biological Chemistry</i> , 2004, 279, 22715-22726.	3.4	47
277	Protein-Tyrosine Phosphatase 1B Associates with Insulin Receptor and Negatively Regulates Insulin Signaling without Receptor Internalization. <i>Journal of Biochemistry</i> , 2004, 136, 89-96.	1.7	31
278	Stiffness and Impaired Blood Flow in Lower-Leg Arteries Are Associated With Severity of Coronary Artery Calcification Among Asymptomatic Type 2 Diabetic Patients. <i>Diabetes Care</i> , 2004, 27, 2409-2415.	8.6	37
279	Reduction of insulin-stimulated glucose uptake by peroxynitrite is concurrent with tyrosine nitration of insulin receptor substrate-1. <i>Biochemical and Biophysical Research Communications</i> , 2004, 320, 639-647.	2.1	65
280	A simple and sensitive method for glutamine:fructose-6-phosphate amidotransferase assay. <i>Journal of Proteomics</i> , 2004, 59, 201-208.	2.4	12
281	Prevalence and Major Risk Factors of Reduced Flow Volume in Lower Extremities With Normal Ankle-Brachial Index in Japanese Patients With Type 2 Diabetes. <i>Diabetes Care</i> , 2003, 26, 1764-1769.	8.6	19
282	Protein-tyrosine Phosphatase 1B as New Activator for Hepatic Lipogenesis via Sterol Regulatory Element-binding Protein-1 Gene Expression. <i>Journal of Biological Chemistry</i> , 2003, 278, 43095-43101.	3.4	70
283	Sumoylation of Pdx1 is associated with its nuclear localization and insulin gene activation. <i>American Journal of Physiology - Endocrinology and Metabolism</i> , 2003, 284, E830-E840.	3.5	81
284	Insulin Activates CCAAT/Enhancer Binding Proteins and Proinflammatory Gene Expression through the Phosphatidylinositol 3-Kinase Pathway in Vascular Smooth Muscle Cells. <i>Journal of Biological Chemistry</i> , 2002, 277, 36631-36639.	3.4	67
285	Amelioration of high fructose-induced metabolic derangements by activation of PPAR $\gamma$ . <i>American Journal of Physiology - Endocrinology and Metabolism</i> , 2002, 282, E1180-E1190.	3.5	172
286	Membrane Localization of 3-Phosphoinositide-dependent Protein Kinase-1 Stimulates Activities of Akt and Atypical Protein Kinase C but Does Not Stimulate Glucose Transport and Glycogen Synthesis in 3T3-L1 Adipocytes. <i>Journal of Biological Chemistry</i> , 2002, 277, 38863-38869.	3.4	31
287	Mechanism for Differential Effect of Protein-Tyrosine Phosphatase 1B on AktVersusMitogen-Activated Protein Kinase in 3T3-L1 Adipocytes. <i>Endocrinology</i> , 2002, 143, 4563-4569.	2.8	23
288	Combined Expression of Pancreatic Duodenal Homeobox 1 and Islet Factor 1 Induces Immature Enterocytes to Produce Insulin. <i>Diabetes</i> , 2002, 51, 1398-1408.	0.6	142

#	ARTICLE	IF	CITATIONS
289	Protein-tyrosine Phosphatase-1B Negatively Regulates Insulin Signaling in L6 Myocytes and Fao Hepatoma Cells. Journal of Biological Chemistry, 2001, 276, 10207-10211.	3.4	126
290	Insulin Production in a Neuroectodermal Tumor that Expresses Islet Factor-1, But Not Pancreatic-Duodenal Homeobox 1. Journal of Clinical Endocrinology and Metabolism, 2001, 86, 1795-1800.	3.6	29
291	Insulin-Induced c-JunN-Terminal Kinase Activation Is Negatively Regulated by Protein Kinase C $\beta$ 1. Endocrinology, 2001, 142, 2669-2676.	2.8	10
292	Evaluation of a New Care System Provided to Diabetic Patients in the Outpatient Clinic.. Internal Medicine, 2000, 39, 783-787.	0.7	7
293	Oral Administration of Tetrahydrobiopterin Prevents Endothelial Dysfunction and Vascular Oxidative Stress in the Aortas of Insulin-Resistant Rats. Circulation Research, 2000, 87, 566-573.	4.5	224
294	Peptide and Protein Library Screening Defines Optimal Substrate Motifs for AKT/PKB. Journal of Biological Chemistry, 2000, 275, 36108-36115.	3.4	349
295	Persistent Activation of Phosphatidylinositol 3-Kinase Causes Insulin Resistance Due to Accelerated Insulin-Induced Insulin Receptor Substrate-1 Degradation in 3T3-L1 Adipocytes*. Endocrinology, 2000, 141, 1930-1935.	2.8	65
296	Endothelium-specific activation of NAD(P)H oxidase in aortas of exogenously hyperinsulinemic rats. American Journal of Physiology - Endocrinology and Metabolism, 1999, 277, E976-E983.	3.5	43
297	Expression of a Dominant Negative SHP-2 in Transgenic Mice Induces Insulin Resistance. Journal of Biological Chemistry, 1999, 274, 30236-30243.	3.4	62
298	.BETA-Sitosterolemia with Generalized Eruptive Xanthomatosis.. Endocrine Journal, 1997, 44, 59-64.	1.6	10
299	SHPTP2 Serves Adapter Protein Linking between Janus Kinase 2 and Insulin Receptor Substrates. Biochemical and Biophysical Research Communications, 1996, 228, 122-127.	2.1	18
300	Lysophosphatidylcholine stimulates the expression and production of MCP-1 by human vascular endothelial cells. Metabolism: Clinical and Experimental, 1996, 45, 559-564.	3.4	83
301	Expression of Dominant Negative Mutant SHPTP2 Attenuates Phosphatidylinositol 3-Kinase Activity via Modulation of Phosphorylation of Insulin Receptor Substrate-1. Journal of Biological Chemistry, 1996, 271, 12595-12602.	3.4	43
302	Insulin Signaling and Its Regulation of System A Amino Acid Uptake in Cultured Rat Vascular Smooth Muscle Cells. Circulation Research, 1996, 79, 1167-1176.	4.5	33
303	High Glucose Condition Desensitizes Insulin Action at the Levels of Receptor Kinase.. Endocrine Journal, 1995, 42, 1-8.	1.6	3
304	Localization of the Insulin-like Growth Factor I Receptor Binding Sites for the SH2 Domain Proteins p85, Syp, and GTPase Activating Protein. Journal of Biological Chemistry, 1995, 270, 19151-19157.	3.4	58
305	Thiazolidine Derivatives Ameliorate High Glucose-induced Insulin Resistance via the Normalization of Protein-tyrosine Phosphatase Activities. Journal of Biological Chemistry, 1995, 270, 7724-7730.	3.4	84
306	Clinical and laboratory characteristics in the families with diabetes and a mitochondrial tRNA <sup>Leu</sup> (UUR) gene mutation. Diabetes Research and Clinical Practice, 1995, 29, 75-82.	2.8	12

#	ARTICLE	IF	CITATIONS
307	Insulin-specific activation of S6 kinase and its desensitization in cultured rat vascular smooth muscle cells. <i>Atherosclerosis</i> , 1995, 113, 19-27.	0.8	15
308	Src homology 2 domains of protein tyrosine phosphatase are associated in vitro with both the insulin receptor and insulin receptor substrate-1 via different phosphotyrosine motifs. <i>FEBS Letters</i> , 1994, 340, 216-220.	2.8	36
309	Co-expression of mutant and normal human insulin receptors in COS 7 cells. <i>Biochimica Et Biophysica Acta Gene Regulatory Mechanisms</i> , 1993, 1216, 425-430.	2.4	0
310	Impaired Autophosphorylation of Insulin Receptors From Abdominal Skeletal Muscles in Nonobese Subjects With NIDDM. <i>Diabetes</i> , 1991, 40, 815-819.	0.6	99
311	The ligand binding characteristics of a kinase-defective A/K1018 human insulin receptor expressed in Rat 1 fibroblasts. <i>Biochimica Et Biophysica Acta - Molecular Cell Research</i> , 1990, 1053, 185-188.	4.1	7
312	Long-term in vitro effects of insulin on insulin binding and glucose transport. <i>Diabetes Research and Clinical Practice</i> , 1986, 2, 1-8.	2.8	11
313	Effect of duration of diabetic state on insulin action in isolated rat soleus muscles. <i>Metabolism: Clinical and Experimental</i> , 1986, 35, 499-504.	3.4	19
314	A primary defect in insulin receptor in a young male patient with insulin resistance. <i>Metabolism: Clinical and Experimental</i> , 1986, 35, 950-955.	3.4	48
315	In vivo and in vitro effect of p-chlorophenoxyisobutyrate on insulin binding and glucose transport in isolated rat adipocytes.. <i>Endocrinologia Japonica</i> , 1985, 32, 829-836.	0.5	0
316	Prolonged Disappearance Rate of a Structurally Abnormal Mutant Insulin from the Circulation in Humans*. <i>Journal of Clinical Endocrinology and Metabolism</i> , 1985, 61, 1142-1145.	3.6	8
317	Inhibition of down regulation by chloroquine in cultured lymphocytes (RPMI-1788 line). <i>Diabetes Research and Clinical Practice</i> , 1985, 1, 145-153.	2.8	3
318	<b>CHANGES IN RECEPTOR BINDING, BIOLOGICAL ACTIVITY AND IMMUNOREACTIVITY OF INSULIN CAUSED BY REPLACING THE RESIDUES B23-B26 WITH </b><b>ALANINE </b>. <i>Biomedical Research</i> , 1984, 5, 267-272.	0.9	8
319	Receptor binding and biological activity of [SerB24]-insulin, an abnormal mutant insulin. <i>Biochemical and Biophysical Research Communications</i> , 1984, 119, 49-57.	2.1	22
320	Receptor-mediated degradation by rat adipocytes: Comparison of A-14 with A-19 125I-labelled insulin.. <i>Endocrinologia Japonica</i> , 1984, 31, 321-326.	0.5	0
321	In Vitro Effects of Glucocorticoid on Glucose Transport in Rat Adipocytes: Evidence of a Post-Receptor Coupling Defect in Insulin Action1. <i>Journal of Biochemistry</i> , 1984, 96, 1893-1902.	1.7	12
322	A new potentiator of insulin action. <i>FEBS Letters</i> , 1983, 163, 50-53.	2.8	21
323	Evidence of the lack of receptor-mediated insulin degradation in human cultured lymphocytes (RPMI-1788 line).. <i>Endocrinologia Japonica</i> , 1983, 30, 679-687.	0.5	2
324	<b>INSULIN BINDING AND GLUCOSE UPTAKE IN ISOLATED SOLEUS MUSCLES IN SPONTANEOUSLY NON-OBESE DIABETIC </b><b>MICE </b>. <i>Biomedical Research</i> , 1983, 4, 533-536.	0.9	9

#	ARTICLE	IF	CITATIONS
325	Effect of acute exercise on insulin binding to erythrocytes in type II diabetes.. Endocrinologia Japonica, 1982, 29, 561-566.	0.5	3
326	Supernormal insulin: [D-PheB24]-insulin with increased affinity for insulin receptors. Biochemical and Biophysical Research Communications, 1982, 107, 329-336.	2.1	66
327	Effect of age and sex on insulin binding to human erythrocytes.. Endocrinologia Japonica, 1981, 28, 623-630.	0.5	1
328	Evaluation of the method of insulin binding studies in human erythrocytes.. Endocrinologia Japonica, 1980, 27, 337-342.	0.5	27
329	The renoprotective effect of onceâ€‘weekly <scp>GLP</scp> â€‘1 receptor agonist dulaglutide on progression of nephropathy in Japanese patients with type 2 diabetes and moderate to severe chronic kidney disease ( <scp>JDDM67</scp> ). Journal of Diabetes Investigation, 0, , .	2.4	1