## Carlo Bruno Giorda

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/4650393/publications.pdf

Version: 2024-02-01

114 papers 3,691 citations

34 h-index 54 g-index

115 all docs

115 docs citations

115 times ranked

5218 citing authors

#	Article	IF	CITATIONS
1	Effects on the incidence of cardiovascular events of the addition of pioglitazone versus sulfonylureas in patients with type 2 diabetes inadequately controlled with metformin (TOSCA.IT): a randomised, multicentre trial. Lancet Diabetes and Endocrinology,the, 2017, 5, 887-897.	11.4	231
2	Incidence of Coronary Heart Disease in Type 2 Diabetic Men and Women. Diabetes Care, 2007, 30, 1241-1247.	8.6	144
3	Aspirin and Simvastatin Combination for Cardiovascular Events Prevention Trial in Diabetes (ACCEPT-D): design of a randomized study of the efficacy of low-dose aspirin in the prevention of cardiovascular events in subjects with diabetes mellitus treated with statins. Trials, 2007, 8, 21.	1.6	140
4	Serum Uric Acid and Risk of CKD in Type 2 Diabetes. Clinical Journal of the American Society of Nephrology: CJASN, 2015, 10, 1921-1929.	4.5	136
5	Sex Disparities in the Quality of Diabetes Care: Biological and Cultural Factors May Play a Different Role for Different Outcomes: A cross-sectional observational study from the AMD Annals initiative. Diabetes Care, 2013, 36, 3162-3168.	8.6	102
6	Incidence and Risk Factors for Stroke in Type 2 Diabetic Patients. Stroke, 2007, 38, 1154-1160.	2.0	98
7	Plasma Triglycerides and HDL-C Levels Predict the Development of Diabetic Kidney Disease in Subjects With Type 2 Diabetes: The AMD Annals Initiative. Diabetes Care, 2016, 39, 2278-2287.	8.6	93
8	Rationale, design, and baseline characteristics in Evaluation of LIXisenatide in Acute Coronary Syndrome, a long-term cardiovascular end point trial of lixisenatide versus placebo. American Heart Journal, 2015, 169, 631-638.e7.	2.7	88
9	Sexual Dysfunction at the Onset of Type 2 Diabetes: The Interplay of Depression, Hormonal and Cardiovascular Factors. Journal of Sexual Medicine, 2014, 11, 2065-2073.	0.6	83
10	Incidence and risk factors for severe and symptomatic hypoglycemia in type 1 diabetes. Results of the HYPOS-1 study. Acta Diabetologica, 2015, 52, 845-853.	2.5	79
11	Pharmacokinetics, safety, and efficacy of DPP-4 inhibitors and GLP-1 receptor agonists in patients with type 2 diabetes mellitus and renal or hepatic impairment. A systematic review of the literature. Endocrine, 2014, 46, 406-419.	2.3	73
12	Recurrence of Cardiovascular Events in Patients With Type 2 Diabetes. Diabetes Care, 2008, 31, 2154-2159.	8.6	71
13	Variability in <scp>HbA1c</scp> , blood pressure, lipid parameters and serum uric acid, and risk of development of chronic kidney disease in type 2 diabetes. Diabetes, Obesity and Metabolism, 2017, 19, 1570-1578.	4.4	70
14	The Burden of NAFLD and Its Characteristics in a Nationwide Population with Type 2 Diabetes. Journal of Diabetes Research, 2016, 2016, 1-9.	2.3	68
15	Incretin-based therapies and acute pancreatitis risk: a systematic review and meta-analysis of observational studies. Endocrine, 2015, 48, 461-471.	2.3	67
16	Impact of severe and symptomatic hypoglycemia on quality of life and fear of hypoglycemia in type 1 and type 2 diabetes. Results of the Hypos-1 observational study. Nutrition, Metabolism and Cardiovascular Diseases, 2019, 29, 736-743.	2.6	67
17	Baseline Quality-of-Care Data From a Quality-Improvement Program Implemented by a Network of Diabetes Outpatient Clinics. Diabetes Care, 2008, 31, 2166-2168.	8.6	61
18	Determinants of Quality in Diabetes Care Process. Diabetes Care, 2009, 32, 1986-1992.	8.6	58

#	Article	IF	Citations
19	Fourâ€year impact of a continuous quality improvement effort implemented by a network of diabetes outpatient clinics: the AMDâ€Annals initiative. Diabetic Medicine, 2010, 27, 1041-1048.	2.3	55
20	Polyphenol intake and cardiovascular risk factors in a population with Âtype 2 diabetes: The TOSCA.IT study. Clinical Nutrition, 2017, 36, 1686-1692.	5.0	52
21	Kidney dysfunction and related cardiovascular risk factors among patients with type 2 diabetes. Nephrology Dialysis Transplantation, 2014, 29, 657-662.	0.7	49
22	Predictors of chronic kidney disease in type 2 diabetes. Medicine (United States), 2016, 95, e4007.	1.0	48
23	Diabetic kidney disease in the elderly: prevalence and clinical correlates. BMC Geriatrics, 2018, 18, 38.	2.7	47
24	The Impact of Adherence to Screening Guidelines and of Diabetes Clinics Referral on Morbidity and Mortality in Diabetes. PLoS ONE, 2012, 7, e33839.	2.5	45
25	Sex differences in food choices, adherence to dietary recommendations and plasma lipid profile in type 2 diabetes – The TOSCA.IT study. Nutrition, Metabolism and Cardiovascular Diseases, 2016, 26, 879-885.	2.6	43
26	Addition of either pioglitazone or a sulfonylurea in type 2 diabetic patients inadequately controlled with metformin alone: Impact on cardiovascular events. A randomized controlled trial. Nutrition, Metabolism and Cardiovascular Diseases, 2012, 22, 997-1006.	2.6	42
27	Identifying patients with type 2 diabetes at high risk of microalbuminuria: results of the DEMAND (Developing Education on Microalbuminuria for Awareness of reNal and cardiovascular risk in) Tj ETQq1 1 0.784	31 <b>4</b> ngBT/	Ovæglock 10
28	Achievement of therapeutic targets in patients with diabetes and chronic kidney disease: insights from the Associazione Medici Diabetologi Annals initiative. Nephrology Dialysis Transplantation, 2015, 30, 1526-1533.	0.7	39
29	Achievement of low density lipoprotein (LDL) cholesterol targets in primary and secondary prevention: Analysis of a large real practice database in Italy. Atherosclerosis, 2019, 285, 40-48.	0.8	39
30	Age- and Gender-Related Differences in LDL-Cholesterol Management in Outpatients with Type 2 Diabetes Mellitus. International Journal of Endocrinology, 2015, 2015, 1-8.	1.5	38
31	Ten-year comparative analysis of incidence, prognosis, and associated factors for dialysis and renal transplantation in type 1 and type 2 diabetes versus non-diabetes. Acta Diabetologica, 2018, 55, 733-740.	2.5	38
32	Dietary intake and major food sources of polyphenols in people with type 2 diabetes: The TOSCA.IT Study. European Journal of Nutrition, 2018, 57, 679-688.	3.9	38
33	Factors Associated with Beta-Cell Dysfunction in Type 2 Diabetes: The BETADECLINE Study. PLoS ONE, 2014, 9, e109702.	2.5	37
34	Incretin therapies and risk of hospital admission for acute pancreatitis in an unselected population of European patients with type 2 diabetes: a case-control study. Lancet Diabetes and Endocrinology,the, 2014, 2, 111-115.	11.4	36
35	Trends over 8Âyears in quality of diabetes care: results of the AMD Annals continuous quality improvement initiative. Acta Diabetologica, 2015, 52, 557-571.	2.5	36
36	The impact of diabetes mellitus on healthcare costs in Italy. Expert Review of Pharmacoeconomics and Outcomes Research, 2011, 11, 709-719.	1.4	34

#	Article	IF	CITATIONS
37	Healthcare resource use, direct and indirect costs of hypoglycemia in type 1 and type 2 diabetes, and nationwide projections. Results of the HYPOS-1 study. Nutrition, Metabolism and Cardiovascular Diseases, 2017, 27, 209-216.	2.6	34
38	Persistent platelet activation in patients with type 2 diabetes treated with low doses of aspirin. Journal of Thrombosis and Haemostasis, 2007, 5, 2197-2203.	3.8	32
39	Gender-Disparities in Adults with Type 1 Diabetes: More Than a Quality of Care Issue. A Cross-Sectional Observational Study from the AMD Annals Initiative. PLoS ONE, 2016, 11, e0162960.	2.5	31
40	Natural history and risk factors for diabetic kidney disease in patients with T2D: lessons from the AMD-annals. Journal of Nephrology, 2019, 32, 517-525.	2.0	30
41	$\hat{l}\pm 1$ -blocker doxazosin improves peripheral insulin sensitivity in diabetic hypertensive patients. Metabolism: Clinical and Experimental, 1995, 44, 673-676.	3.4	29
42	The prevalence of coronary heart disease in TypeÂ2 diabetic patients in Italy: the DAI study. Diabetic Medicine, 2004, 21, 738-745.	2.3	29
43	Predictors of early-stage left ventricular dysfunction in type 2 diabetes: results of DYDA study. European Journal of Cardiovascular Prevention and Rehabilitation, 2011, 18, 415-423.	2.8	28
44	Analysis of midwall shortening reveals high prevalence of left ventricular myocardial dysfunction in patients with diabetes mellitus: the DYDA study. European Journal of Preventive Cardiology, 2012, 19, 935-943.	1.8	28
45	Blood pressure status and the incidence of diabetic kidney disease in patients with hypertension and type 2 diabetes. Journal of Hypertension, 2016, 34, 2090-2098.	0.5	28
46	Overall Quality of Care Predicts the Variability of Key Risk Factors for Complications in Type 2 Diabetes: An Observational, Longitudinal Retrospective Study. Diabetes Care, 2019, 42, 514-519.	8.6	28
47	Obesity and changes in urine albumin/creatinine ratio in patients with type 2 diabetes: TheÂDEMAND Study. Nutrition, Metabolism and Cardiovascular Diseases, 2010, 20, 110-116.	2.6	27
48	The SUBITO-DE study: sexual dysfunction in newly diagnosed type 2 diabetes male patients. Journal of Endocrinological Investigation, 2013, 36, 864-8.	3.3	27
49	Epidemiology of diabetic kidney disease in adult patients with type 1 diabetes in Italy: The AMDâ€Annals initiative. Diabetes/Metabolism Research and Reviews, 2017, 33, e2873.	4.0	26
50	Correlates of total homocysteine plasma concentration in type 2 diabetes. European Journal of Clinical Investigation, 2004, 34, 197-204.	3.4	25
51	Metabolic syndrome, serum uric acid and renal risk in patients with T2D. PLoS ONE, 2017, 12, e0176058.	2.5	25
52	Ezetimibe + simvastatin versus doubling the dose of simvastatin in high cardiovascular risk diabetics: a multicenter, randomized trial (the LEAD study). Cardiovascular Diabetology, 2010, 9, 20.	6.8	24
53	Effects of doxazosin, a selective $\hat{l}\pm 1$ -inhibitor, on plasma insulin and blood glucose response to a glucose tolerance test in essential hypertension. Metabolism: Clinical and Experimental, 1993, 42, 1440-1442.	3.4	23
54	Hospitalisation for heart failure and mortality associated with dipeptidyl peptidase 4 (DPP-4) inhibitor use in an unselected population of subjects with type 2 diabetes: a nested case-control study. BMJ Open, 2015, 5, e007959-e007959.	1.9	23

#	Article	IF	CITATIONS
55	Sexual Dysfunction in Type 2 Diabetes at Diagnosis: Progression over Time and Drug and Non-Drug Correlated Factors. PLoS ONE, 2016, 11, e0157915.	2.5	23
56	Predictors of chronic kidney disease in type 1 diabetes: a longitudinal study from the AMD Annals initiative. Scientific Reports, 2017, 7, 3313.	3.3	23
57	Long-term blood pressure variability and development of chronic kidney disease in type 2 diabetes. Journal of Hypertension, 2019, 37, 805-813.	0.5	23
58	Occurrence over time and regression of nonalcoholic fatty liver disease in type 2 diabetes. Diabetes/Metabolism Research and Reviews, 2017, 33, e2878.	4.0	22
59	Association of kidney disease measures with risk of renal function worsening in patients with hypertension and type 2 diabetes. Journal of Diabetes and Its Complications, 2017, 31, 419-426.	2.3	22
60	Mild hyperhomocysteinemia, C677T polymorphism on methylenetetrahydrofolate reductase gene and the risk of macroangiopathy in type 2 diabetes: a prospective study. Acta Diabetologica, 2011, 48, 95-101.	2.5	21
61	Resistant Hypertension, Timeâ€Updated Blood Pressure Values and Renal Outcome in Type 2 Diabetes Mellitus. Journal of the American Heart Association, 2017, 6, .	3.7	21
62	Trend over time in hepatic fibrosis score in a cohort of type 2 diabetes patients. Diabetes Research and Clinical Practice, 2018, 135, 65-72.	2.8	21
63	The impact of second-level specialized care on hospitalization in persons with diabetes: a multilevel population-based study. Diabetic Medicine, 2006, 23, 377-383.	2.3	20
64	Recommendations for the implementation of international standardization of glycated hemoglobin in Italy. Clinical Chemistry and Laboratory Medicine, 2010, 48, 623-626.	2.3	20
65	A systematic review of acute pancreatitis as an adverse event of type 2 diabetes drugs: from hard facts to a balanced position. Diabetes, Obesity and Metabolism, 2014, 16, 1041-1047.	4.4	20
66	Prevalence, incidence and associated comorbidities of treated hypothyroidism: an update from a European population. European Journal of Endocrinology, 2017, 176, 533-542.	3.7	20
67	Italian Association of Clinical Endocrinologists (AME) & Italian Association of Clinical Diabetologists (AMD) Position Statement. Endocrine, 2015, 49, 339-352.	2.3	19
68	Apparent Treatment Resistant Hypertension, Blood Pressure Control and the Progression of Chronic Kidney Disease in Patients with Type 2 Diabetes. Kidney and Blood Pressure Research, 2018, 43, 422-438.	2.0	19
69	Association of physicians' accuracy in recording with quality of care in cardiovascular medicine. European Journal of Cardiovascular Prevention and Rehabilitation, 2009, 16, 722-728.	2.8	17
70	Mortality, incidence of cardiovascular diseases, and educational level among the diabetic and non-diabetic populations in two large Italian cities. Diabetes Research and Clinical Practice, 2011, 92, 205-212.	2.8	17
71	Inappropriately high left ventricular mass in patients with type 2 diabetes mellitus and no overt cardiac disease. The DYDA study. Journal of Hypertension, 2011, 29, 1994-2003.	0.5	17
72	The under-use of statin in type 2 diabetic patients attending diabetic clinics in Italy. Nutrition, Metabolism and Cardiovascular Diseases, 2007, 17, 32-40.	2.6	16

#	Article	IF	CITATIONS
73	Incidence of hospitalization and mortality in patients with diabetic foot regardless of amputation: a population study. Acta Diabetologica, 2020, 57, 221-228.	2.5	16
74	Left ventricular dysfunction and outcome at two-year follow-up in patients with type 2 diabetes: The DYDA study. Diabetes Research and Clinical Practice, 2013, 101, 236-242.	2.8	15
<b>7</b> 5	The role of the care model in modifying prognosis in diabetes. Nutrition, Metabolism and Cardiovascular Diseases, 2013, 23, 11-16.	2.6	15
76	Normoalbuminuric kidney impairment in patients with T1DM: insights from annals initiative. Diabetology and Metabolic Syndrome, 2018, 10, 60.	2.7	15
77	Management of newly diagnosed patients with type 2 diabetes: what are the attitudes of physicians? A SUBITO!AMD survey on the early diabetes treatment in Italy. Acta Diabetologica, 2012, 49, 429-433.	2.5	14
78	Global cardiovascular risk management in different Italian regions: An analysis of the evaluation of final feasible effect of control training and ultra sensitisation (EFFECTUS) educational program. Nutrition, Metabolism and Cardiovascular Diseases, 2012, 22, 635-642.	2.6	14
79	Improving quality of care in people with Type 2 diabetes through the Associazione Medici Diabetologiâ€annals initiative: a longâ€ŧerm costâ€effectiveness analysis. Diabetic Medicine, 2014, 31, 615-623.	2.3	14
80	Global Cardiovascular Risk Assessment in Different Clinical Settings. High Blood Pressure and Cardiovascular Prevention, 2009, 16, 55-63.	2.2	13
81	Use of Electronic Support for Implementing Global Cardiovascular Risk Management. High Blood Pressure and Cardiovascular Prevention, 2010, 17, 37-47.	2.2	13
82	Five-Year Predictors of Insulin Initiation in People with Type 2 Diabetes under Real-Life Conditions. Journal of Diabetes Research, 2018, 2018, 1-10.	2.3	13
83	Diabetes-specific variables associated withÂquality of life changes in young diabetic people: The type 1 diabetes Registry of Turin (Italy). Nutrition, Metabolism and Cardiovascular Diseases, 2013, 23, 1031-1036.	2.6	12
84	Comparison of direct costs of type 2 diabetes care: Different care models with different outcomes. Nutrition, Metabolism and Cardiovascular Diseases, 2014, 24, 717-724.	2.6	12
85	Incidence and correlated factors of beta cell failure in a 4-year follow-up of patients with type 2 diabetes: a longitudinal analysis of the BETADECLINE study. Acta Diabetologica, 2016, 53, 761-767.	2.5	12
86	Impact of Diabetes Mellitus on the Clinical Management of Global Cardiovascular Risk: Analysis of the Results of the Evaluation of Final Feasible Effect of Control Training and Ultra Sensitization (EFFECTUS) Educational Program. Clinical Cardiology, 2011, 34, 560-566.	1.8	11
87	Cardiovascular Effects of Pioglitazone or Sulfonylureas According to Pretreatment Risk: Moving Toward Personalized Care. Journal of Clinical Endocrinology and Metabolism, 2019, 104, 3296-3302.	3.6	11
88	Long-term blood pressure variability, incidence of hypertension and changes in renal function in type 2 diabetes. Journal of Hypertension, 2020, 38, 2279-2286.	0.5	11
89	Changes in albuminuria and renal outcome in patients with type 2 diabetes and hypertension. Journal of Hypertension, 2018, 36, 1719-1728.	0.5	10
90	Beta cell stress in a 4â€year followâ€up of patients with type 2 diabetes: A longitudinal analysis of the <i>BetaDecline</i> Study. Diabetes/Metabolism Research and Reviews, 2018, 34, e3016.	4.0	10

#	Article	IF	CITATIONS
91	Portrait of women with type 1 or type 2 diabetes of childbearing age attending diabetes clinics in Italy: the AMD-Annals initiative. Acta Diabetologica, 2018, 55, 193-199.	2.5	9
92	Could clinical inertia in part explain the unexpected association of insulin therapy with poorer cardiovascular outcomes in observational studies on diabetes?. Diabetes Research and Clinical Practice, 2011, 92, e47-e48.	2.8	8
93	Comment on: Inzucchi et al. Management of Hyperglycemia in Type 2 Diabetes: A Patient-Centered Approach. Position Statement of the American Diabetes Association (ADA) and the European Association for the Study of Diabetes (EASD). Diabetes Care 2012;35:1364-1379. Diabetes Care, 2012, 35, e71-e71.	8.6	8
94	Secondary prevention of coronary artery disease in high-risk diabetic patients. Nutrition, Metabolism and Cardiovascular Diseases, 2003, 13, 238-243.	2.6	7
95	Self-Monitoring of Blood Glucose in Type 2 Diabetes: Steps toward consensus. Diabetes Care, 2007, 30, e105-e105.	8.6	7
96	An Analysis of the Management of Cardiovascular Risk Factors in Routine Clinical Practice in Italy. High Blood Pressure and Cardiovascular Prevention, 2011, 18, 19-30.	2.2	7
97	From swab testing to health outcomes within the T2DM population: Impact of diabetes background on COVID19 progression. Diabetes Research and Clinical Practice, 2021, 180, 109021.	2.8	7
98	Antihyperglycemic treatment in patients with type 2 diabetes in Italy: the impact of age and kidney function. Oncotarget, 2017, 8, 62039-62048.	1.8	7
99	Determinants of good metabolic control without weight gain in type 2 diabetes management: a machine learning analysis. BMJ Open Diabetes Research and Care, 2020, 8, e001362.	2.8	6
100	To what extent is the new position statement of the American Diabetes Association (ADA) and the European Association for the Study of Diabetes (EASD) †personalised'?. Diabetologia, 2012, 55, 2853-2855.	6.3	5
101	Incretin-based therapy and risk of cholangiocarcinoma: a nested case–control study in a population of subjects with type 2 diabetes. Acta Diabetologica, 2020, 57, 401-408.	2.5	5
102	Prescription of Sulphonylureas among Patients with Type 2 Diabetes Mellitus in Italy: Results from the Retrospective, Observational Multicentre Cross-Sectional SUSCIPE (Sulphonyl_UreaS_Correct_Internal_Prescription_Evaluation) Study. Diabetes Therapy, 2020, 11, 2105-2119.	2.5	5
103	Hepatic fibrosis of any origin in a large population of type 2 diabetes patients. Nutrition, Metabolism and Cardiovascular Diseases, 2021, 31, 2887-2894.	2.6	5
104	Short Course of Insulin Treatment versus Metformin in Newly Diagnosed Patients with Type 2 Diabetes. Journal of Clinical Medicine, 2018, 7, 235.	2.4	4
105	Effects of linagliptin on left ventricular DYsfunction in patients with type 2 DiAbetes and concentric left ventricular geometry: results of the DYDA 2 trial. European Journal of Preventive Cardiology, 2021, 28, 8-17.	1.8	4
106	Factors associated with a rapid normalization of HbA1c in newly diagnosed type 2 diabetes patients seen in a specialist setting. Acta Diabetologica, 2013, 50, 81-87.	2.5	3
107	Cardiovascular Biomarkers, Cardiac Dysfunction, and Outcomes in Patients With Type 2 Diabetes: A Prospective, Multicenter Study. Diabetes Care, 2013, 36, e137-e138.	8.6	3
108	Results of the Adequacy of glycemiC Control in pAtients with type 2 Diabetes mEllitus treated with Metformin monotherapY at the maximal-tolerated dose (ACCADEMY) study. Endocrine, 2016, 52, 507-515.	2.3	3

#	Article	IF	CITATIONS
109	Effects of Dipeptidyl Peptidase-4 Inhibitor Linagliptin on Left Ventricular Dysfunction in Patients with Type 2 Diabetes and Concentric Left Ventricular Geometry (the DYDA 2â,,¢ Trial). Rationale, Design, and Baseline Characteristics of the Study Population. Cardiovascular Drugs and Therapy, 2019, 33, 547-555.	2.6	3
110	Instructive lessons from the analysis of assistance in diabetes during the first phase of COVID-19 pandemic. Acta Diabetologica, 2022, 59, 861-864.	2.5	3
111	Impact of physicians' age on the clinical management of global cardiovascular risk: analysis of the results of the Evaluation of Final Feasible Effect of Control Training and Ultra Sensitisation Educational Programme. International Journal of Clinical Practice, 2011, 65, 649-657.	1.7	2
112	Association of kidney disease measures with risk of renal function worsening in patients with type 1 diabetes. BMC Nephrology, 2018, 19, 347.	1.8	2
113	Clinical characteristics and patterns of care of newly diagnosed type 2 diabetic patients. Nutrition, Metabolism and Cardiovascular Diseases, 2007, 17, e31-e33.	2.6	1

The implementation of international standardization of glycated hemoglobin. A "red-letter-day" for glycated hemoglobin in Italy: 1/1/11. Italian Recommendations of GLAD Working Group (A1c delegates) Tj ETQq0 @@rgBT /Overlock 10