## Carlo Bruno Giorda

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/4650393/publications.pdf

Version: 2024-02-01

114 papers 3,691 citations

34 h-index 54 g-index

115 all docs

115 docs citations

115 times ranked

5218 citing authors

| #  | Article  | IF  | CITATIONS |
|----|--|-----|-----------|
| 1  | Instructive lessons from the analysis of assistance in diabetes during the first phase of COVID-19 pandemic. Acta Diabetologica, 2022, 59, 861-864.  | 2.5 | 3         |
| 2  | Effects of linagliptin on left ventricular DYsfunction in patients with type 2 DiAbetes and concentric left ventricular geometry: results of the DYDA 2 trial. European Journal of Preventive Cardiology, 2021, 28, 8-17.  | 1.8 | 4         |
| 3  | Hepatic fibrosis of any origin in a large population of type 2 diabetes patients. Nutrition, Metabolism and Cardiovascular Diseases, 2021, 31, 2887-2894.  | 2.6 | 5         |
| 4  | From swab testing to health outcomes within the T2DM population: Impact of diabetes background on COVID19 progression. Diabetes Research and Clinical Practice, 2021, 180, 109021.   | 2.8 | 7         |
| 5  | Incidence of hospitalization and mortality in patients with diabetic foot regardless of amputation: a population study. Acta Diabetologica, 2020, 57, 221-228.   | 2.5 | 16        |
| 6  | Incretin-based therapy and risk of cholangiocarcinoma: a nested case–control study in a population of subjects with type 2 diabetes. Acta Diabetologica, 2020, 57, 401-408.  | 2.5 | 5         |
| 7  | Long-term blood pressure variability, incidence of hypertension and changes in renal function in type 2 diabetes. Journal of Hypertension, 2020, 38, 2279-2286.  | 0.5 | 11        |
| 8  | Prescription of Sulphonylureas among Patients with Type 2 Diabetes Mellitus in Italy: Results from the Retrospective, Observational Multicentre Cross-Sectional SUSCIPE (Sulphonyl_UreaS_Correct_Internal_Prescription_Evaluation) Study. Diabetes Therapy, 2020, 11, 2105-2119.   | 2.5 | 5         |
| 9  | Determinants of good metabolic control without weight gain in type 2 diabetes management: a machine learning analysis. BMJ Open Diabetes Research and Care, 2020, 8, e001362.  | 2.8 | 6         |
| 10 | Cardiovascular Effects of Pioglitazone or Sulfonylureas According to Pretreatment Risk: Moving Toward Personalized Care. Journal of Clinical Endocrinology and Metabolism, 2019, 104, 3296-3302.   | 3.6 | 11        |
| 11 | Impact of severe and symptomatic hypoglycemia on quality of life and fear of hypoglycemia in type 1 and type 2 diabetes. Results of the Hypos-1 observational study. Nutrition, Metabolism and Cardiovascular Diseases, 2019, 29, 736-743.   | 2.6 | 67        |
| 12 | Achievement of low density lipoprotein (LDL) cholesterol targets in primary and secondary prevention: Analysis of a large real practice database in Italy. Atherosclerosis, 2019, 285, 40-48.  | 0.8 | 39        |
| 13 | Overall Quality of Care Predicts the Variability of Key Risk Factors for Complications in Type 2<br>Diabetes: An Observational, Longitudinal Retrospective Study. Diabetes Care, 2019, 42, 514-519.  | 8.6 | 28        |
| 14 | Long-term blood pressure variability and development of chronic kidney disease in type 2 diabetes. Journal of Hypertension, 2019, 37, 805-813.   | 0.5 | 23        |
| 15 | Natural history and risk factors for diabetic kidney disease in patients with T2D: lessons from the AMD-annals. Journal of Nephrology, 2019, 32, 517-525.  | 2.0 | 30        |
| 16 | Effects of Dipeptidyl Peptidase-4 Inhibitor Linagliptin on Left Ventricular Dysfunction in Patients with Type 2 Diabetes and Concentric Left Ventricular Geometry (the DYDA 2â,, † Trial). Rationale, Design, and Baseline Characteristics of the Study Population. Cardiovascular Drugs and Therapy, 2019, 33, 547-555. | 2.6 | 3         |
| 17 | Changes in albuminuria and renal outcome in patients with type 2 diabetes and hypertension. Journal of Hypertension, 2018, 36, 1719-1728.  | 0.5 | 10        |
| 18 | Ten-year comparative analysis of incidence, prognosis, and associated factors for dialysis and renal transplantation in type 1 and type 2 diabetes versus non-diabetes. Acta Diabetologica, 2018, 55, 733-740.   | 2.5 | 38        |

| #  | Article   | IF   | CITATIONS |
|----|---|------|-----------|
| 19 | Beta cell stress in a 4â€year followâ€up of patients with type 2 diabetes: A longitudinal analysis of the <i>BetaDecline</i> Study. Diabetes/Metabolism Research and Reviews, 2018, 34, e3016.  | 4.0  | 10        |
| 20 | Apparent Treatment Resistant Hypertension, Blood Pressure Control and the Progression of Chronic Kidney Disease in Patients with Type 2 Diabetes. Kidney and Blood Pressure Research, 2018, 43, 422-438.  | 2.0  | 19        |
| 21 | Dietary intake and major food sources of polyphenols in people with type 2 diabetes: The TOSCA.IT Study. European Journal of Nutrition, 2018, 57, 679-688.  | 3.9  | 38        |
| 22 | Trend over time in hepatic fibrosis score in a cohort of type 2 diabetes patients. Diabetes Research and Clinical Practice, 2018, 135, 65-72.   | 2.8  | 21        |
| 23 | Portrait of women with type $1$ or type $2$ diabetes of childbearing age attending diabetes clinics in Italy: the AMD-Annals initiative. Acta Diabetologica, 2018, 55, 193-199.   | 2.5  | 9         |
| 24 | Association of kidney disease measures with risk of renal function worsening in patients with type 1 diabetes. BMC Nephrology, 2018, 19, 347.   | 1.8  | 2         |
| 25 | Five-Year Predictors of Insulin Initiation in People with Type 2 Diabetes under Real-Life Conditions.<br>Journal of Diabetes Research, 2018, 2018, 1-10.  | 2.3  | 13        |
| 26 | Diabetic kidney disease in the elderly: prevalence and clinical correlates. BMC Geriatrics, 2018, 18, 38.   | 2.7  | 47        |
| 27 | Short Course of Insulin Treatment versus Metformin in Newly Diagnosed Patients with Type 2<br>Diabetes. Journal of Clinical Medicine, 2018, 7, 235.   | 2.4  | 4         |
| 28 | Normoalbuminuric kidney impairment in patients with T1DM: insights from annals initiative. Diabetology and Metabolic Syndrome, 2018, 10, 60.  | 2.7  | 15        |
| 29 | Prevalence, incidence and associated comorbidities of treated hypothyroidism: an update from a European population. European Journal of Endocrinology, 2017, 176, 533-542.  | 3.7  | 20        |
| 30 | Variability in <scp>HbAlc</scp> , blood pressure, lipid parameters and serum uric acid, and risk of development of chronic kidney disease in type 2 diabetes. Diabetes, Obesity and Metabolism, 2017, 19, 1570-1578.  | 4.4  | 70        |
| 31 | Occurrence over time and regression of nonalcoholic fatty liver disease in type 2 diabetes.  Diabetes/Metabolism Research and Reviews, 2017, 33, e2878.   | 4.0  | 22        |
| 32 | Epidemiology of diabetic kidney disease in adult patients with type 1 diabetes in Italy: The AMDâ€Annals initiative. Diabetes/Metabolism Research and Reviews, 2017, 33, e2873.   | 4.0  | 26        |
| 33 | Healthcare resource use, direct and indirect costs of hypoglycemia in type 1 and type 2 diabetes, and nationwide projections. Results of the HYPOS-1 study. Nutrition, Metabolism and Cardiovascular Diseases, 2017, 27, 209-216.   | 2.6  | 34        |
| 34 | Resistant Hypertension, Timeâ€Updated Blood Pressure Values and Renal Outcome in Type 2 Diabetes Mellitus. Journal of the American Heart Association, 2017, 6, .  | 3.7  | 21        |
| 35 | Effects on the incidence of cardiovascular events of the addition of pioglitazone versus sulfonylureas in patients with type 2 diabetes inadequately controlled with metformin (TOSCA.IT): a randomised, multicentre trial. Lancet Diabetes and Endocrinology, the, 2017, 5, 887-897. | 11.4 | 231       |
| 36 | Predictors of chronic kidney disease in type 1 diabetes: a longitudinal study from the AMD Annals initiative. Scientific Reports, 2017, 7, 3313.  | 3.3  | 23        |

| #  | Article   | IF  | Citations |
|----|---|-----|-----------|
| 37 | Association of kidney disease measures with risk of renal function worsening in patients with hypertension and type 2 diabetes. Journal of Diabetes and Its Complications, 2017, 31, 419-426.   | 2.3 | 22        |
| 38 | Polyphenol intake and cardiovascular risk factors in a population withÂtype 2 diabetes: The TOSCA.IT study. Clinical Nutrition, 2017, 36, 1686-1692.  | 5.0 | 52        |
| 39 | Metabolic syndrome, serum uric acid and renal risk in patients with T2D. PLoS ONE, 2017, 12, e0176058.  | 2.5 | 25        |
| 40 | Antihyperglycemic treatment in patients with type 2 diabetes in Italy: the impact of age and kidney function. Oncotarget, 2017, 8, 62039-62048.   | 1.8 | 7         |
| 41 | The Burden of NAFLD and Its Characteristics in a Nationwide Population with Type 2 Diabetes. Journal of Diabetes Research, 2016, 2016, 1-9.   | 2.3 | 68        |
| 42 | Sexual Dysfunction in Type 2 Diabetes at Diagnosis: Progression over Time and Drug and Non-Drug Correlated Factors. PLoS ONE, 2016, 11, e0157915.   | 2.5 | 23        |
| 43 | Blood pressure status and the incidence of diabetic kidney disease in patients with hypertension and type 2 diabetes. Journal of Hypertension, 2016, 34, 2090-2098.   | 0.5 | 28        |
| 44 | Sex differences in food choices, adherence to dietary recommendations and plasma lipid profile in type 2 diabetes – The TOSCA.IT study. Nutrition, Metabolism and Cardiovascular Diseases, 2016, 26, 879-885.                               | 2.6 | 43        |
| 45 | Incidence and correlated factors of beta cell failure in a 4-year follow-up of patients with type 2 diabetes: a longitudinal analysis of the BETADECLINE study. Acta Diabetologica, 2016, 53, 761-767.                                      | 2.5 | 12        |
| 46 | Plasma Triglycerides and HDL-C Levels Predict the Development of Diabetic Kidney Disease in Subjects With Type 2 Diabetes: The AMD Annals Initiative. Diabetes Care, 2016, 39, 2278-2287.   | 8.6 | 93        |
| 47 | Predictors of chronic kidney disease in type 2 diabetes. Medicine (United States), 2016, 95, e4007.   | 1.0 | 48        |
| 48 | Results of the Adequacy of glycemiC Control in pAtients with type 2 Diabetes mEllitus treated with Metformin monotherapy at the maximal-tolerated dose (ACCADEMY) study. Endocrine, 2016, 52, 507-515.                                      | 2.3 | 3         |
| 49 | Gender-Disparities in Adults with Type 1 Diabetes: More Than a Quality of Care Issue. A Cross-Sectional Observational Study from the AMD Annals Initiative. PLoS ONE, 2016, 11, e0162960.   | 2.5 | 31        |
| 50 | Age- and Gender-Related Differences in LDL-Cholesterol Management in Outpatients with Type 2 Diabetes Mellitus. International Journal of Endocrinology, 2015, 2015, 1-8.  | 1.5 | 38        |
| 51 | Rationale, design, and baseline characteristics in Evaluation of LIXisenatide in Acute Coronary<br>Syndrome, a long-term cardiovascular end point trial of lixisenatide versus placebo. American Heart<br>Journal, 2015, 169, 631-638.e7.   | 2.7 | 88        |
| 52 | Hospitalisation for heart failure and mortality associated with dipeptidyl peptidase 4 (DPP-4) inhibitor use in an unselected population of subjects with type 2 diabetes: a nested case-control study. BMJ Open, 2015, 5, e007959-e007959. | 1.9 | 23        |
| 53 | Italian Association of Clinical Endocrinologists (AME) & Italian Association of Clinical Diabetologists (AMD) Position Statement. Endocrine, 2015, 49, 339-352.   | 2.3 | 19        |
| 54 | Trends over 8Âyears in quality of diabetes care: results of the AMD Annals continuous quality improvement initiative. Acta Diabetologica, 2015, 52, 557-571.  | 2.5 | 36        |

| #  | Article  | IF   | CITATIONS |
|----|--|------|-----------|
| 55 | Incidence and risk factors for severe and symptomatic hypoglycemia in type 1 diabetes. Results of the HYPOS-1 study. Acta Diabetologica, 2015, 52, 845-853.  | 2.5  | 79        |
| 56 | Achievement of therapeutic targets in patients with diabetes and chronic kidney disease: insights from the Associazione Medici Diabetologi Annals initiative. Nephrology Dialysis Transplantation, 2015, 30, 1526-1533.                      | 0.7  | 39        |
| 57 | Serum Uric Acid and Risk of CKD in Type 2 Diabetes. Clinical Journal of the American Society of Nephrology: CJASN, 2015, 10, 1921-1929.  | 4.5  | 136       |
| 58 | Incretin-based therapies and acute pancreatitis risk: a systematic review and meta-analysis of observational studies. Endocrine, 2015, 48, 461-471.  | 2.3  | 67        |
| 59 | Kidney dysfunction and related cardiovascular risk factors among patients with type 2 diabetes.<br>Nephrology Dialysis Transplantation, 2014, 29, 657-662.   | 0.7  | 49        |
| 60 | Sexual Dysfunction at the Onset of Type 2 Diabetes: The Interplay of Depression, Hormonal and Cardiovascular Factors. Journal of Sexual Medicine, 2014, 11, 2065-2073.   | 0.6  | 83        |
| 61 | Improving quality of care in people with Type 2 diabetes through the Associazione Medici<br>Diabetologiâ€annals initiative: a longâ€ŧerm costâ€effectiveness analysis. Diabetic Medicine, 2014, 31, 615-623.                                 | 2.3  | 14        |
| 62 | Pharmacokinetics, safety, and efficacy of DPP-4 inhibitors and GLP-1 receptor agonists in patients with type 2 diabetes mellitus and renal or hepatic impairment. A systematic review of the literature. Endocrine, 2014, 46, 406-419.       | 2.3  | 73        |
| 63 | Incretin therapies and risk of hospital admission for acute pancreatitis in an unselected population of European patients with type 2 diabetes: a case-control study. Lancet Diabetes and Endocrinology,the, 2014, 2, 111-115.               | 11.4 | 36        |
| 64 | Comparison of direct costs of type 2 diabetes care: Different care models with different outcomes. Nutrition, Metabolism and Cardiovascular Diseases, 2014, 24, 717-724.   | 2.6  | 12        |
| 65 | A systematic review of acute pancreatitis as an adverse event of type 2 diabetes drugs: from hard facts to a balanced position. Diabetes, Obesity and Metabolism, 2014, 16, 1041-1047.   | 4.4  | 20        |
| 66 | Factors Associated with Beta-Cell Dysfunction in Type 2 Diabetes: The BETADECLINE Study. PLoS ONE, 2014, 9, e109702.   | 2.5  | 37        |
| 67 | Sex Disparities in the Quality of Diabetes Care: Biological and Cultural Factors May Play a Different Role for Different Outcomes: A cross-sectional observational study from the AMD Annals initiative. Diabetes Care, 2013, 36, 3162-3168. | 8.6  | 102       |
| 68 | Left ventricular dysfunction and outcome at two-year follow-up in patients with type 2 diabetes: The DYDA study. Diabetes Research and Clinical Practice, 2013, 101, 236-242.  | 2.8  | 15        |
| 69 | Factors associated with a rapid normalization of HbA1c in newly diagnosed type 2 diabetes patients seen in a specialist setting. Acta Diabetologica, 2013, 50, 81-87.  | 2.5  | 3         |
| 70 | Diabetes-specific variables associated withÂquality of life changes in young diabetic people: The type 1 diabetes Registry of Turin (Italy). Nutrition, Metabolism and Cardiovascular Diseases, 2013, 23, 1031-1036.                         | 2.6  | 12        |
| 71 | The role of the care model in modifying prognosis in diabetes. Nutrition, Metabolism and Cardiovascular Diseases, 2013, 23, 11-16.   | 2.6  | 15        |
| 72 | Cardiovascular Biomarkers, Cardiac Dysfunction, and Outcomes in Patients With Type 2 Diabetes: A Prospective, Multicenter Study. Diabetes Care, 2013, 36, e137-e138.   | 8.6  | 3         |

| #  | Article  | IF                 | CITATIONS           |
|----|--|--------------------|---------------------|
| 73 | The SUBITO-DE study: sexual dysfunction in newly diagnosed type 2 diabetes male patients. Journal of Endocrinological Investigation, 2013, 36, 864-8.  | 3.3                | 27                  |
| 74 | Comment on: Inzucchi et al. Management of Hyperglycemia in Type 2 Diabetes: A Patient-Centered Approach. Position Statement of the American Diabetes Association (ADA) and the European Association for the Study of Diabetes (EASD). Diabetes Care 2012;35:1364-1379. Diabetes Care, 2012, 35, e71-e71. | 8.6                | 8                   |
| 75 | Analysis of midwall shortening reveals high prevalence of left ventricular myocardial dysfunction in patients with diabetes mellitus: the DYDA study. European Journal of Preventive Cardiology, 2012, 19, 935-943.  | 1.8                | 28                  |
| 76 | Management of newly diagnosed patients with type 2 diabetes: what are the attitudes of physicians? A SUBITO!AMD survey on the early diabetes treatment in Italy. Acta Diabetologica, 2012, 49, 429-433.  | 2.5                | 14                  |
| 77 | Global cardiovascular risk management in different Italian regions: An analysis of the evaluation of final feasible effect of control training and ultra sensitisation (EFFECTUS) educational program.  Nutrition, Metabolism and Cardiovascular Diseases, 2012, 22, 635-642.                            | 2.6                | 14                  |
| 78 | Addition of either pioglitazone or a sulfonylurea in type 2 diabetic patients inadequately controlled with metformin alone: Impact on cardiovascular events. A randomized controlled trial. Nutrition, Metabolism and Cardiovascular Diseases, 2012, 22, 997-1006.                                       | 2.6                | 42                  |
| 79 | To what extent is the new position statement of the American Diabetes Association (ADA) and the European Association for the Study of Diabetes (EASD) †personalised'?. Diabetologia, 2012, 55, 2853-2855.  | 6.3                | 5                   |
| 80 | The Impact of Adherence to Screening Guidelines and of Diabetes Clinics Referral on Morbidity and Mortality in Diabetes. PLoS ONE, 2012, 7, e33839.  | 2.5                | 45                  |
| 81 | The implementation of international standardization of glycated hemoglobin. A "red-letter-day" for glycated hemoglobin in Italy: $1/1/11$ . Italian Recommendations of GLAD Working Group (A1c delegates) Tj ETQq1   | ls <b>0</b> x78431 | . <b>⊕</b> rgBT /O√ |
| 82 | An Analysis of the Management of Cardiovascular Risk Factors in Routine Clinical Practice in Italy. High Blood Pressure and Cardiovascular Prevention, 2011, 18, 19-30.  | 2.2                | 7                   |
| 83 | Could clinical inertia in part explain the unexpected association of insulin therapy with poorer cardiovascular outcomes in observational studies on diabetes?. Diabetes Research and Clinical Practice, 2011, 92, e47-e48.  | 2.8                | 8                   |
| 84 | Mortality, incidence of cardiovascular diseases, and educational level among the diabetic and non-diabetic populations in two large Italian cities. Diabetes Research and Clinical Practice, 2011, 92, 205-212.  | 2.8                | 17                  |
| 85 | Inappropriately high left ventricular mass in patients with type 2 diabetes mellitus and no overt cardiac disease. The DYDA study. Journal of Hypertension, 2011, 29, 1994-2003.   | 0.5                | 17                  |
| 86 | Impact of physicians' age on the clinical management of global cardiovascular risk: analysis of the results of the Evaluation of Final Feasible Effect of Control Training and Ultra Sensitisation Educational Programme. International Journal of Clinical Practice, 2011, 65, 649-657.                 | 1.7                | 2                   |
| 87 | Impact of Diabetes Mellitus on the Clinical Management of Global Cardiovascular Risk: Analysis of the Results of the Evaluation of Final Feasible Effect of Control Training and Ultra Sensitization (EFFECTUS) Educational Program. Clinical Cardiology, 2011, 34, 560-566.                             | 1.8                | 11                  |
| 88 | Mild hyperhomocysteinemia, C677T polymorphism on methylenetetrahydrofolate reductase gene and the risk of macroangiopathy in type 2 diabetes: a prospective study. Acta Diabetologica, 2011, 48, 95-101.   | 2.5                | 21                  |
| 89 | Predictors of early-stage left ventricular dysfunction in type 2 diabetes: results of DYDA study. European Journal of Cardiovascular Prevention and Rehabilitation, 2011, 18, 415-423.   | 2.8                | 28                  |
| 90 | The impact of diabetes mellitus on healthcare costs in Italy. Expert Review of Pharmacoeconomics and Outcomes Research, 2011, 11, 709-719.   | 1.4                | 34                  |

| #   | Article  | IF                | Citations   |
|-----|--|-------------------|-------------|
| 91  | Ezetimibe + simvastatin versus doubling the dose of simvastatin in high cardiovascular risk diabetics: a multicenter, randomized trial (the LEAD study). Cardiovascular Diabetology, 2010, 9, 20.  | 6.8               | 24          |
| 92  | Fourâ€year impact of a continuous quality improvement effort implemented by a network of diabetes outpatient clinics: the AMDâ€Annals initiative. Diabetic Medicine, 2010, 27, 1041-1048.  | 2.3               | 55          |
| 93  | Recommendations for the implementation of international standardization of glycated hemoglobin in Italy. Clinical Chemistry and Laboratory Medicine, 2010, 48, 623-626.  | 2.3               | 20          |
| 94  | Obesity and changes in urine albumin/creatinine ratio in patients with type 2 diabetes: TheÂDEMAND Study. Nutrition, Metabolism and Cardiovascular Diseases, 2010, 20, 110-116.  | 2.6               | 27          |
| 95  | Use of Electronic Support for Implementing Global Cardiovascular Risk Management. High Blood Pressure and Cardiovascular Prevention, 2010, 17, 37-47.  | 2.2               | 13          |
| 96  | Determinants of Quality in Diabetes Care Process. Diabetes Care, 2009, 32, 1986-1992.  | 8.6               | 58          |
| 97  | Association of physicians' accuracy in recording with quality of care in cardiovascular medicine.<br>European Journal of Cardiovascular Prevention and Rehabilitation, 2009, 16, 722-728.  | 2.8               | 17          |
| 98  | Global Cardiovascular Risk Assessment in Different Clinical Settings. High Blood Pressure and Cardiovascular Prevention, 2009, 16, 55-63.  | 2.2               | 13          |
| 99  | Recurrence of Cardiovascular Events in Patients With Type 2 Diabetes. Diabetes Care, 2008, 31, 2154-2159.  | 8.6               | 71          |
| 100 | Baseline Quality-of-Care Data From a Quality-Improvement Program Implemented by a Network of Diabetes Outpatient Clinics. Diabetes Care, 2008, 31, 2166-2168.  | 8.6               | 61          |
| 101 | ldentifying patients with type 2 diabetes at high risk of microalbuminuria: results of the DEMAND (Developing Education on Microalbuminuria for Awareness of reNal and cardiovascular risk in) Tj ETQq1 1 0.7843   | 1 <b>⊕</b> ngBT/0 | Ovæølock 10 |
| 102 | Incidence and Risk Factors for Stroke in Type 2 Diabetic Patients. Stroke, 2007, 38, 1154-1160.  | 2.0               | 98          |
| 103 | Self-Monitoring of Blood Glucose in Type 2 Diabetes: Steps toward consensus. Diabetes Care, 2007, 30, e105-e105.   | 8.6               | 7           |
| 104 | Incidence of Coronary Heart Disease in Type 2 Diabetic Men and Women. Diabetes Care, 2007, 30, 1241-1247.  | 8.6               | 144         |
| 105 | The under-use of statin in type 2 diabetic patients attending diabetic clinics in Italy. Nutrition, Metabolism and Cardiovascular Diseases, 2007, 17, 32-40.   | 2.6               | 16          |
| 106 | Clinical characteristics and patterns of care of newly diagnosed type 2 diabetic patients. Nutrition, Metabolism and Cardiovascular Diseases, 2007, 17, e31-e33.   | 2.6               | 1           |
| 107 | Aspirin and Simvastatin Combination for Cardiovascular Events Prevention Trial in Diabetes (ACCEPT-D): design of a randomized study of the efficacy of low-dose aspirin in the prevention of cardiovascular events in subjects with diabetes mellitus treated with statins. Trials, 2007, 8, 21. | 1.6               | 140         |
| 108 | Persistent platelet activation in patients with type 2 diabetes treated with low doses of aspirin. Journal of Thrombosis and Haemostasis, 2007, 5, 2197-2203.  | 3.8               | 32          |

| #   | Article  | IF  | CITATION |
|-----|--|-----|----------|
| 109 | The impact of second-level specialized care on hospitalization in persons with diabetes: a multilevel population-based study. Diabetic Medicine, 2006, 23, 377-383.  | 2.3 | 20       |
| 110 | The prevalence of coronary heart disease in TypeÂ2 diabetic patients in Italy: the DAI study. Diabetic Medicine, 2004, 21, 738-745.  | 2.3 | 29       |
| 111 | Correlates of total homocysteine plasma concentration in type 2 diabetes. European Journal of Clinical Investigation, 2004, 34, 197-204.   | 3.4 | 25       |
| 112 | Secondary prevention of coronary artery disease in high-risk diabetic patients. Nutrition, Metabolism and Cardiovascular Diseases, 2003, 13, 238-243.  | 2.6 | 7        |
| 113 | $\hat{l}\pm 1$ -blocker doxazosin improves peripheral insulin sensitivity in diabetic hypertensive patients. Metabolism: Clinical and Experimental, 1995, 44, 673-676.   | 3.4 | 29       |
| 114 | Effects of doxazosin, a selective $\hat{l}\pm 1$ -inhibitor, on plasma insulin and blood glucose response to a glucose tolerance test in essential hypertension. Metabolism: Clinical and Experimental, 1993, 42, 1440-1442. | 3.4 | 23       |