

Jing Xu

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4652009/publications.pdf>

Version: 2024-02-01

27
papers

1,149
citations

471509

17
h-index

526287

27
g-index

27
all docs

27
docs citations

27
times ranked

1999
citing authors

#	ARTICLE	IF	CITATIONS
1	Tumor-derived adenosine promotes macrophage proliferation in human hepatocellular carcinoma. <i>Journal of Hepatology</i> , 2021, 74, 627-637.	3.7	64
2	Reprogramming immunosuppressive myeloid cells by activated T cells promotes the response to anti-PD-1 therapy in colorectal cancer. <i>Signal Transduction and Targeted Therapy</i> , 2021, 6, 4.	17.1	51
3	High S100A9+ cell density predicts a poor prognosis in hepatocellular carcinoma patients after curative resection. <i>Aging</i> , 2021, 13, 16367-16380.	3.1	16
4	HHLA2 Expression is Associated with Poor Survival in Patients with Hepatocellular Carcinoma. <i>Biologics: Targets and Therapy</i> , 2021, Volume 15, 329-341.	3.2	5
5	C-Reactive Protein Is an Indicator of the Immunosuppressive Microenvironment Fostered by Myeloid Cells in Hepatocellular Carcinoma. <i>Frontiers in Oncology</i> , 2021, 11, 774823.	2.8	5
6	Myeloid signature reveals immune contexture and predicts the prognosis of hepatocellular carcinoma. <i>Journal of Clinical Investigation</i> , 2020, 130, 4679-4693.	8.2	42
7	Macrophages induce CD47 upregulation via IL-6 and correlate with poor survival in hepatocellular carcinoma patients. <i>Oncolmmunology</i> , 2019, 8, e1652540.	4.6	55
8	CD103 ⁺ tumor-infiltrating lymphocytes predict favorable prognosis in patients with esophageal squamous cell carcinoma. <i>Journal of Cancer</i> , 2019, 10, 5234-5243.	2.5	16
9	Immune Cell Infiltration in the Microenvironment of Liver Oligometastasis from Colorectal Cancer: Intratumoural CD8/CD3 Ratio Is a Valuable Prognostic Index for Patients Undergoing Liver Metastasectomy. <i>Cancers</i> , 2019, 11, 1922.	3.7	11
10	Vessels That Encapsulate Tumor Clusters (VETC) Pattern Is a Predictor of Sorafenib Benefit in Patients with Hepatocellular Carcinoma. <i>Hepatology</i> , 2019, 70, 824-839.	7.3	62
11	Monocytes/Macrophages promote vascular CXCR4 expression via the ERK pathway in hepatocellular carcinoma. <i>Oncolmmunology</i> , 2018, 7, e1408745.	4.6	29
12	Expression patterns of programmed death ligand 1 correlate with different microenvironments and patient prognosis in hepatocellular carcinoma. <i>British Journal of Cancer</i> , 2018, 119, 80-88.	6.4	74
13	Spleen mediates a distinct hematopoietic progenitor response supporting tumor-promoting myelopoiesis. <i>Journal of Clinical Investigation</i> , 2018, 128, 3425-3438.	8.2	111
14	High NKG2A expression contributes to NK cell exhaustion and predicts a poor prognosis of patients with liver cancer. <i>Oncolmmunology</i> , 2017, 6, e1264562.	4.6	180
15	Vascular CXCR4 Expression Promotes Vessel Sprouting and Sensitivity to Sorafenib Treatment in Hepatocellular Carcinoma. <i>Clinical Cancer Research</i> , 2017, 23, 4482-4492.	7.0	38
16	Distinct patterns and prognostic values of tumor-infiltrating macrophages in hepatocellular carcinoma and gastric cancer. <i>Journal of Translational Medicine</i> , 2017, 15, 37.	4.4	41
17	Association of decreased expression of the macrophage scavenger receptor MARCO with tumor progression and poor prognosis in human hepatocellular carcinoma. <i>Journal of Gastroenterology and Hepatology (Australia)</i> , 2017, 32, 1107-1114.	2.8	32
18	Density of CD8+ lymphocytes in biopsy samples combined with the circulating lymphocyte ratio predicts pathologic complete response to chemoradiotherapy for rectal cancer. <i>Cancer Management and Research</i> , 2017, Volume 9, 701-708.	1.9	11

#	ARTICLE	IF	CITATIONS
19	The Bidirectional Regulation between MYL5 and HIF-1 α Promotes Cervical Carcinoma Metastasis. <i>Theranostics</i> , 2017, 7, 3768-3780.	10.0	17
20	GLUT1 and ASCT2 as Predictors for Prognosis of Hepatocellular Carcinoma. <i>PLoS ONE</i> , 2016, 11, e0168907.	2.5	79
21	CD200R, a co-inhibitory receptor on immune cells, predicts the prognosis of human hepatocellular carcinoma. <i>Immunology Letters</i> , 2016, 178, 105-113.	2.5	10
22	CD169 identifies an activated CD8+T cell subset in regional lymph nodes that predicts favorable prognosis in colorectal cancer patients. <i>Oncolmmunology</i> , 2016, 5, e1177690.	4.6	15
23	Tumor-infiltrating macrophages express interleukin-25 and predict a favorable prognosis in patients with gastric cancer after radical resection. <i>Oncotarget</i> , 2016, 7, 11083-11093.	1.8	13
24	Lactate Dehydrogenase Is an Important Prognostic Indicator for Hepatocellular Carcinoma after Partial Hepatectomy. <i>Translational Oncology</i> , 2015, 8, 497-503.	3.7	28
25	Elevated expression of Cripto-1 correlates with poor prognosis in hepatocellular carcinoma. <i>Oncotarget</i> , 2015, 6, 35116-35128.	1.8	29
26	The predictive value of centre tumour CD8+ T cells in patients with hepatocellular carcinoma: comparison with Immunoscore. <i>Oncotarget</i> , 2015, 6, 35602-35615.	1.8	60
27	CXCL17 Expression Predicts Poor Prognosis and Correlates with Adverse Immune Infiltration in Hepatocellular Carcinoma. <i>PLoS ONE</i> , 2014, 9, e110064.	2.5	55