

Dietmar Kuhl

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4680476/publications.pdf>

Version: 2024-02-01

104
papers

13,201
citations

41344

49
h-index

30087

103
g-index

105
all docs

105
docs citations

105
times ranked

13669
citing authors

#	ARTICLE	IF	CITATIONS
1	The adaptor protein PICK1 targets the sorting receptor SorLA. <i>Molecular Brain</i> , 2022, 15, 18.	2.6	3
2	Converging roles of PSENEN/PEN2 and CLN3 in the autophagy-lysosome system. <i>Autophagy</i> , 2022, 18, 2068-2085.	9.1	12
3	Neuronal activity regulates alternative exon usage. <i>Molecular Brain</i> , 2020, 13, 148.	2.6	7
4	Disturbed Prefrontal Cortex Activity in the Absence of Schizophrenia-Like Behavioral Dysfunction in Arc/Arg3.1 Deficient Mice. <i>Journal of Neuroscience</i> , 2019, 39, 8149-8163.	3.6	11
5	Cognitive impairment and autistic-like behaviour in SAPAP4-deficient mice. <i>Translational Psychiatry</i> , 2019, 9, 7.	4.8	13
6	Amyloidosis causes downregulation of <i>SorLA</i> , <i>SorCS1</i> and <i>SorCS3</i> expression in mice. <i>Biological Chemistry</i> , 2019, 400, 1181-1189.	2.5	13
7	Arc/Arg3.1 mediates a critical period for spatial learning and hippocampal networks. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2018, 115, 12531-12536.	7.1	38
8	Structural Properties of Synaptic Transmission and Temporal Dynamics at Excitatory Layer 5B Synapses in the Adult Rat Somatosensory Cortex. <i>Frontiers in Synaptic Neuroscience</i> , 2018, 10, 24.	2.5	31
9	SGK1 induces vascular smooth muscle cell calcification through NF- κ B signaling. <i>Journal of Clinical Investigation</i> , 2018, 128, 3024-3040.	8.2	114
10	SGK1-dependent ENaC processing and trafficking in mice with high dietary K intake and elevated aldosterone. <i>American Journal of Physiology - Renal Physiology</i> , 2017, 312, F65-F76.	2.7	33
11	Neuronal activity-regulated alternative mRNA splicing. <i>International Journal of Biochemistry and Cell Biology</i> , 2017, 91, 184-193.	2.8	23
12	Profiling the MAPK/ERK dependent and independent activity regulated transcriptional programs in the murine hippocampus in vivo. <i>Scientific Reports</i> , 2017, 7, 45101.	3.3	48
13	SGK1 up-regulates Orai1 expression and VSMC migration during neointima formation after arterial injury. <i>Thrombosis and Haemostasis</i> , 2017, 117, 1002-1005.	3.4	10
14	TMIC-20. INHIBITION OF SLC7A11 REDUCES EXCITATORY SYNAPTIC INPUT OF PERITUMORAL NEURONS IN GLIOMA PATIENTS. <i>Neuro-Oncology</i> , 2017, 19, vi247-vi247.	1.2	0
15	The Kinesin KIF21B Regulates Microtubule Dynamics and Is Essential for Neuronal Morphology, Synapse Function, and Learning and Memory. <i>Cell Reports</i> , 2016, 15, 968-977.	6.4	70
16	Effects of Arc/Arg3.1 gene deletion on rhythmic synchronization of hippocampal CA1 neurons during locomotor activity and sleep. <i>Neurobiology of Learning and Memory</i> , 2016, 131, 155-165.	1.9	9
17	Revisiting the neuronal localization and trafficking of <i>CLN3</i> in juvenile neuronal ceroid lipofuscinosis. <i>Journal of Neurochemistry</i> , 2016, 139, 456-470.	3.9	24
18	Arc/Arg3.1 governs inflammatory dendritic cell migration from the skin and thereby controls T cell activation. <i>Science Immunology</i> , 2016, 1, eaaf8665.	11.9	40

#	ARTICLE	IF	CITATIONS
19	BDNF-induced LTP is associated with rapid Arc/Arg3.1-dependent enhancement in adult hippocampal neurogenesis. <i>Scientific Reports</i> , 2016, 6, 21222.	3.3	74
20	Pivotal Role of Serum- and Glucocorticoid-Inducible Kinase 1 in Vascular Inflammation and Atherogenesis. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2015, 35, 547-557.	2.4	55
21	Sor<sc>CS</sc>1 variants and amyloid precursor protein (<sc>APP</sc>) are coâ€transported in neurons but only Sor<sc>CS</sc>1c modulates anterograde <sc>APP</sc> transport. <i>Journal of Neurochemistry</i> , 2015, 135, 60-75.	3.9	20
22	Lack of the serum- and glucocorticoid-inducible kinase SGK1 improves muscle force characteristics and attenuates fibrosis in dystrophic mdx mouse muscle. <i>Pflugers Archiv European Journal of Physiology</i> , 2015, 467, 1965-1974.	2.8	17
23	Serum- and Glucocorticoid-Inducible Kinase 1 Sensitive NF-#954;B Signaling in Dendritic Cells. <i>Cellular Physiology and Biochemistry</i> , 2014, 34, 943-954.	1.6	34
24	Activation of serum/glucocorticoidâ€induced kinase 1 (SGK1) is important to maintain skeletal muscle homeostasis and prevent atrophy. <i>EMBO Molecular Medicine</i> , 2013, 5, 80-91.	6.9	100
25	Sgk1-Dependent Stimulation of Cardiac Na⁺/H⁺Exchanger Nhe1 by Dexamethasone. <i>Cellular Physiology and Biochemistry</i> , 2013, 32, 25-38.	1.6	654
26	Genome-Wide Profiling of the Activity-Dependent Hippocampal Transcriptome. <i>PLoS ONE</i> , 2013, 8, e76903.	2.5	38
27	SGK1 Sensitivity of Platelet Migration. <i>Cellular Physiology and Biochemistry</i> , 2012, 30, 259-268.	1.6	37
28	The serum- and glucocorticoid-inducible kinase 1 (SGK1) influences platelet calcium signaling and function by regulation of Orai1 expression in megakaryocytes. <i>Blood</i> , 2012, 119, 251-261.	1.4	126
29	Sgk1 sensitivity of Na ⁺ /H ⁺ exchanger activity and cardiac remodeling following pressure overload. <i>Basic Research in Cardiology</i> , 2012, 107, 236.	5.9	47
30	Arc/Arg3.1 Regulates an Endosomal Pathway Essential for Activity-Dependent Î²-Amyloid Generation. <i>Cell</i> , 2011, 147, 615-628.	28.9	183
31	Activity-Induced Notch Signaling in Neurons Requires Arc/Arg3.1 and Is Essential for Synaptic Plasticity in Hippocampal Networks. <i>Neuron</i> , 2011, 69, 437-444.	8.1	184
32	SGK1-dependent stimulation of intestinal SGLT1 activity by vitamin D. <i>Pflugers Archiv European Journal of Physiology</i> , 2011, 462, 489-494.	2.8	11
33	Stimulation of Ca ²⁺ â€channel Orai1/STIM1 by serumâ€and glucocorticoidâ€inducible kinase 1 (SGK1). <i>FASEB Journal</i> , 2011, 25, 2012-2021.	0.5	82
34	Arc-dependent synapse-specific homeostatic plasticity. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2011, 108, 816-821.	7.1	165
35	Widespread transcription at neuronal activity-regulated enhancers. <i>Nature</i> , 2010, 465, 182-187.	27.8	2,120
36	SRF binding to SRE 6.9 in the Arc promoter is essential for LTD in cultured Purkinje cells. <i>Nature Neuroscience</i> , 2010, 13, 1082-1089.	14.8	72

#	ARTICLE	IF	CITATIONS
37	SGK1-dependent Intestinal Tumor Growth in APC-deficient Mice. <i>Cellular Physiology and Biochemistry</i> , 2010, 25, 271-278.	1.6	34
38	Serum- and glucocorticoid-regulated kinase 1 is upregulated following unilateral ureteral obstruction causing epithelialâ€mesenchymal transition. <i>Kidney International</i> , 2010, 78, 668-678.	5.2	58
39	A Specific Requirement of Arc/Arg3.1 for Visual Experience-Induced Homeostatic Synaptic Plasticity in Mouse Primary Visual Cortex. <i>Journal of Neuroscience</i> , 2010, 30, 7168-7178.	3.6	123
40	Nociceptive Stimulation Induces Expression of Arc/Arg3.1 in the Spinal Cord with a Preference for Neurons Containing Enkephalin. <i>Molecular Pain</i> , 2010, 6, 1744-8069-6-43.	2.1	21
41	The Serum and Glucocorticoid-Regulated Kinase 1 in Hypoxic Renal Injury. <i>Cellular Physiology and Biochemistry</i> , 2009, 24, 577-584.	1.6	24
42	Pioglitazone Induced Gastric Acid Secretion. <i>Cellular Physiology and Biochemistry</i> , 2009, 24, 193-200.	1.6	14
43	SGK1-sensitive renal tubular glucose reabsorption in diabetes. <i>American Journal of Physiology - Renal Physiology</i> , 2009, 296, F859-F866.	2.7	29
44	Expression and phosphorylation of the Na ⁺ -Cl ⁻ cotransporter NCC in vivo is regulated by dietary salt, potassium, and SGK1. <i>American Journal of Physiology - Renal Physiology</i> , 2009, 297, F704-F712.	2.7	225
45	Hyperaldosteronism, hypovolemia, and increased blood pressure in mice expressing defective APC. <i>American Journal of Physiology - Regulatory Integrative and Comparative Physiology</i> , 2009, 297, R571-R575.	1.8	23
46	Impaired Mast Cell Activation in Gene-Targeted Mice Lacking the Serum- and Glucocorticoid-Inducible Kinase SGK1. <i>Journal of Immunology</i> , 2009, 183, 4395-4402.	0.8	29
47	Relative resistance of SGK1 knockout mice against chemical carcinogenesis. <i>IUBMB Life</i> , 2009, 61, 768-776.	3.4	53
48	SGK1 dependence of insulin induced hypokalemia. <i>Pflügers Archiv European Journal of Physiology</i> , 2009, 457, 955-961.	2.8	16
49	Stimulation of electrogenic intestinal dipeptide transport by the glucocorticoid dexamethasone. <i>Pflügers Archiv European Journal of Physiology</i> , 2009, 459, 191-202.	2.8	14
50	Fluorescent Arc/Arg3.1 indicator mice: A versatile tool to study brain activity changes in vitro and in vivo. <i>Journal of Neuroscience Methods</i> , 2009, 184, 25-36.	2.5	43
51	Lack of the serum and glucocorticoid-inducible kinase SGK1 attenuates the volume retention after treatment with the PPAR β agonist pioglitazone. <i>Pflügers Archiv European Journal of Physiology</i> , 2008, 456, 425-436.	2.8	28
52	Two rat brain Staufen isoforms differentially bind RNA. <i>Journal of Neurochemistry</i> , 2008, 76, 155-165.	3.9	62
53	Different Motifs Regulate Trafficking of SorCS1 Isoforms. <i>Traffic</i> , 2008, 9, 980-994.	2.7	39
54	Elongation Factor 2 and Fragile X Mental Retardation Protein Control the Dynamic Translation of Arc/Arg3.1 Essential for mGluR-LTD. <i>Neuron</i> , 2008, 59, 70-83.	8.1	471

#	ARTICLE	IF	CITATIONS
55	SGK1-Dependent Upregulation of Connective Tissue Growth Factor by Angiotensin II. <i>Kidney and Blood Pressure Research</i> , 2008, 31, 80-86.	2.0	16
56	Role of Serum- and Glucocorticoid-Inducible Kinase SGK1 in Glucocorticoid Regulation of Renal Electrolyte Excretion and Blood Pressure. <i>Kidney and Blood Pressure Research</i> , 2008, 31, 280-289.	2.0	19
57	Role of maternal glucocorticoid inducible kinase SGK1 in fetal programming of blood pressure in response to prenatal diet. <i>American Journal of Physiology - Regulatory Integrative and Comparative Physiology</i> , 2008, 294, R2008-R2013.	1.8	28
58	Dexamethasone increases Na ⁺ /K ⁺ ATPase activity in insulin secreting cells through SGK1. <i>Biochemical and Biophysical Research Communications</i> , 2007, 352, 662-667.	2.1	25
59	Role of the serum and glucocorticoid inducible kinase SGK1 in glucocorticoid stimulation of gastric acid secretion. <i>Pflügers Archiv European Journal of Physiology</i> , 2007, 455, 493-503.	2.8	29
60	Aldosterone-induced Sgk1 relieves Dot1a-Af9â€œmediated transcriptional repression of epithelial Na ⁺ channel β . <i>Journal of Clinical Investigation</i> , 2007, 117, 773-783.	8.2	150
61	Arc/Arg3.1 Is Essential for the Consolidation of Synaptic Plasticity and Memories. <i>Neuron</i> , 2006, 52, 437-444.	8.1	743
62	Arc/Arg3.1 Interacts with the Endocytic Machinery to Regulate AMPA Receptor Trafficking. <i>Neuron</i> , 2006, 52, 445-459.	8.1	691
63	Arc/Arg3.1 Mediates Homeostatic Synaptic Scaling of AMPA Receptors. <i>Neuron</i> , 2006, 52, 475-484.	8.1	684
64	The N-terminus of the serum- and glucocorticoid-inducible kinase Sgk1 specifies mitochondrial localization and rapid turnover. <i>Biochemical Journal</i> , 2006, 399, 69-76.	3.7	28
65	Renal Ca ²⁺ handling in sgk1 knockout mice. <i>Pflügers Archiv European Journal of Physiology</i> , 2006, 452, 444-452.	2.8	17
66	SGK1 is not required for regulation of colonic ENaC activity. <i>Pflügers Archiv European Journal of Physiology</i> , 2006, 453, 97-105.	2.8	17
67	SGK1-dependent cardiac CTGF formation and fibrosis following DOCA treatment. <i>Journal of Molecular Medicine</i> , 2006, 84, 396-404.	3.9	111
68	Blunted DOCA/high salt induced albuminuria and renal tubulointerstitial damage in gene-targeted mice lacking SGK1. <i>Journal of Molecular Medicine</i> , 2006, 84, 737-746.	3.9	49
69	Resistance of mice lacking the serum- and glucocorticoid-inducible kinase SGK1 against salt-sensitive hypertension induced by a high-fat diet. <i>American Journal of Physiology - Renal Physiology</i> , 2006, 291, F1264-F1273.	2.7	62
70	Intestinal function of gene-targeted mice lacking serum- and glucocorticoid-inducible kinase 1. <i>American Journal of Physiology - Renal Physiology</i> , 2006, 290, G1114-G1123.	3.4	53
71	Blunted hypertensive effect of combined fructose and high-salt diet in gene-targeted mice lacking functional serum- and glucocorticoid-inducible kinase SGK1. <i>American Journal of Physiology - Regulatory Integrative and Comparative Physiology</i> , 2006, 290, R935-R944.	1.8	64
72	Renal function of gene-targeted mice lacking both SGK1 and SGK3. <i>American Journal of Physiology - Regulatory Integrative and Comparative Physiology</i> , 2006, 290, R945-R950.	1.8	44

#	ARTICLE	IF	CITATIONS
73	DOCA-induced Phosphorylation of Glycogen Synthase Kinase 3 β . Cellular Physiology and Biochemistry, 2006, 17, 137-144.	1.6	37
74	Serum- and Glucocorticoid-Inducible Kinase 1 Mediates Salt Sensitivity of Glucose Tolerance. Diabetes, 2006, 55, 2059-2066.	0.6	41
75	Deranged Kv channel regulation in fibroblasts from mice lacking the serum and glucocorticoid inducible kinase SGK1. Journal of Cellular Physiology, 2005, 204, 87-98.	4.1	11
76	Role of Sgk1 in salt and potassium homeostasis. American Journal of Physiology - Regulatory Integrative and Comparative Physiology, 2005, 288, R4-R10.	1.8	64
77	SGK1 as a determinant of kidney function and salt intake in response to mineralocorticoid excess. American Journal of Physiology - Regulatory Integrative and Comparative Physiology, 2005, 289, R395-R401.	1.8	66
78	Serum- and Glucocorticoid-Inducible Kinase 1 (SGK1) Mediates Glucocorticoid-Induced Inhibition of Insulin Secretion. Diabetes, 2005, 54, 1090-1099.	0.6	155
79	Regulation of the excitatory amino acid transporter EAAT5 by the serum and glucocorticoid dependent kinases SGK1 and SGK3. Biochemical and Biophysical Research Communications, 2005, 329, 738-742.	2.1	34
80	Impaired Regulation of Renal K ⁺ Elimination in the sgk1-Knockout Mouse. Journal of the American Society of Nephrology: JASN, 2004, 15, 885-891.	6.1	115
81	The three sorCS genes are differentially expressed and regulated by synaptic activity. Journal of Neurochemistry, 2004, 88, 1470-1476.	3.9	66
82	MGluRs regulate the expression of neuronal calcium sensor proteins NCS-1 and VILIP-1 and the immediate early gene arg3.1/arc in the hippocampus in vivo. Biochemical and Biophysical Research Communications, 2004, 322, 1073-1079.	2.1	29
83	Exploitation of KESTREL to identify NDRG family members as physiological substrates for SGK1 and GSK3. Biochemical Journal, 2004, 384, 477-488.	3.7	299
84	Cerebral localization and regulation of the cell volume-sensitive serum- and glucocorticoid-dependent kinase SGK1. Pflugers Archiv European Journal of Physiology, 2002, 443, 617-624.	2.8	75
85	Induction of Glycerol Phosphate Dehydrogenase Gene Expression During Seizure and Analgesia. Journal of Neurochemistry, 2002, 75, 1419-1428.	3.9	6
86	Impaired renal Na ⁺ retention in the sgk1-knockout mouse. Journal of Clinical Investigation, 2002, 110, 1263-1268.	8.2	271
87	Impaired renal Na ⁺ retention in the sgk1-knockout mouse. Journal of Clinical Investigation, 2002, 110, 1263-1268.	8.2	196
88	Impaired renal Na ⁺ retention in the sgk1-knockout mouse. Journal of Clinical Investigation, 2002, 110, 1263-1268.	8.2	60
89	Arg3.1/Arc mRNA Induction by Ca ²⁺ and cAMP Requires Protein Kinase A and Mitogen-Activated Protein Kinase/Extracellular Regulated Kinase Activation. Journal of Neuroscience, 2001, 21, 5484-5493.	3.6	239
90	Odors regulate Arc expression in neuronal ensembles engaged in odor processing. NeuroReport, 2000, 11, 1809-1813.	1.2	37

#	ARTICLE	IF	CITATIONS
91	Adhesion induced expression of the serine/threonine kinase Fnk in human macrophages. <i>Oncogene</i> , 2000, 19, 4832-4839.	5.9	62
92	The polo-like protein kinases Fnk and Snk associate with a Ca ²⁺ - and integrin-binding protein and are regulated dynamically with synaptic plasticity. <i>EMBO Journal</i> , 1999, 18, 5528-5539.	7.8	200
93	Novelty-induced increased expression of immediate-early genes c-fos and arg 3.1 in the mouse brain. , 1999, 38, 234-246.		126
94	Pim kinase expression is induced by LTP stimulation and required for the consolidation of enduring LTP. <i>EMBO Journal</i> , 1999, 18, 3359-3369.	7.8	72
95	Dendritic localization of mRNAs. <i>Current Opinion in Neurobiology</i> , 1998, 8, 600-606.	4.2	94
96	A subtractive hybridisation method for the enrichment of moderately induced sequences. <i>Nucleic Acids Research</i> , 1998, 26, 1359-1361.	14.5	8
97	A different form of long-lasting potentiation revealed in tissue plasminogen activator mutant mice. <i>Journal of Neuroscience</i> , 1996, 16, 2057-2063.	3.6	204
98	A tri-hybrid system for the analysis and detection of RNA-protein interactions. <i>Nucleic Acids Research</i> , 1996, 24, 4838-4840.	14.5	79
99	Somatodendritic expression of an immediate early gene is regulated by synaptic activity.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1995, 92, 5734-5738.	7.1	659
100	Tissue-plasminogen activator is induced as an immediate-early gene during seizure, kindling and long-term potentiation. <i>Nature</i> , 1993, 361, 453-457.	27.8	771
101	Long-term sensitization training in <i>Aplysia</i> leads to an increase in the expression of BiP, the major protein chaperon of the ER.. <i>Journal of Cell Biology</i> , 1992, 119, 1069-1076.	5.2	83
102	Long-Term sensitization training in <i>Aplysia</i> leads to an increase in calreticulin, a major presynaptic calcium-binding protein. <i>Neuron</i> , 1992, 9, 1013-1024.	8.1	76
103	Different pathways mediate virus inducibility of the human IFN- β 1 and IFN- β 2 genes. <i>Cell</i> , 1990, 60, 767-779.	28.9	177
104	Reversible silencing of enhancers by sequences derived from the human IFN- β promoter. <i>Cell</i> , 1987, 50, 1057-1069.	28.9	133