## Alexander M Van Der Linden

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/4727446/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Loss of circRNAs from the <i>crhâ€l </i> gene extends the mean lifespan in <i>Caenorhabditis elegans</i> . Aging Cell, 2022, 21, e13560.	6.7	6
2	Dietary vitamin B12 regulates chemosensory receptor gene expression via the MEF2 transcription factor in <i>Caenorhabditis elegans</i> . G3: Genes, Genomes, Genetics, 2022, 12, .	1.8	4
3	A salt-induced kinase is required for the metabolic regulation of sleep. PLoS Biology, 2020, 18, e3000220.	5.6	37
4	Regulation of sleep by KIN-29 is not developmental. MicroPublication Biology, 2020, 2020, .	0.1	0
5	The salt-inducible kinase KIN-29 regulates lifespan via the class II histone-deacetylase HDA-4. MicroPublication Biology, 2020, 2020, .	0.1	Ο
6	Global accumulation of circRNAs during aging in Caenorhabditis elegans. BMC Genomics, 2018, 19, 8.	2.8	139
7	Increased food intake after starvation enhances sleep in Drosophila melanogaster. Journal of Genetics and Genomics, 2017, 44, 319-326.	3.9	18
8	Cell-Autonomous and Non-Cell-Autonomous Regulation of a Feeding State-Dependent Chemoreceptor Gene via MEF-2 and bHLH Transcription Factors. PLoS Genetics, 2016, 12, e1006237.	3.5	21
9	Long-term imaging of circadian locomotor rhythms of a freely crawling C. elegans population. Journal of Neuroscience Methods, 2015, 249, 66-74.	2.5	11
10	Plasticity of chemoreceptor gene expression: Sensory and circuit inputs modulate state-dependent chemoreceptors. Worm, 2015, 4, e1023497.	1.0	4
11	Feeding State, Insulin and NPR-1 Modulate Chemoreceptor Gene Expression via Integration of Sensory and Circuit Inputs. PLoS Genetics, 2014, 10, e1004707.	3.5	42
12	Differential hippocampal gene expression is associated with climateâ€related natural variation in memory and the hippocampus in foodâ€caching chickadees. Molecular Ecology, 2013, 22, 397-408.	3.9	29
13	Genome-Wide Analysis of Light- and Temperature-Entrained Circadian Transcripts in Caenorhabditis elegans. PLoS Biology, 2010, 8, e1000503.	5.6	60
14	<i>Cis</i> â€regulatory mechanisms of gene expression in an olfactory neuron type in <i>Caenorhabditis elegans</i> . Developmental Dynamics, 2009, 238, 3080-3092.	1.8	18
15	The ECL-4 PKG Acts With KIN-29 Salt-Inducible Kinase and Protein Kinase A to Regulate Chemoreceptor Gene Expression and Sensory Behaviors in <i>Caenorhabditis elegans</i> . Genetics, 2008, 180, 1475-1491.	2.9	47
16	Left-right olfactory asymmetry results from antagonistic functions of voltage-activated calcium channels and the Raw repeat protein OLRN-1 in C. elegans. Neural Development, 2007, 2, 24.	2.4	61
17	KIN-29 SIK regulates chemoreceptor gene expression via an MEF2 transcription factor and a class II HDAC. EMBO Journal, 2007, 26, 358-370.	7.8	84
18	Chemical Genetics Reveals an RGS/G-Protein Role in the Action of a Compound. PLoS Genetics, 2006, 2, e57.	3.5	32

#	Article	IF	CITATIONS
19	Shotgun Cloning of Transposon Insertions in the Genome ofCaenorhabditis elegans. Comparative and Functional Genomics, 2004, 5, 225-229.	2.0	9
20	Hyperactivation of the G12-Mediated Signaling Pathway in Caenorhabditis elegans Induces a Developmental Growth Arrest via Protein Kinase C. Current Biology, 2003, 13, 516-521.	3.9	21
21	Proteins Interacting withCaenorhabditis elegans CαSubunits. Comparative and Functional Genomics, 2003, 4, 479-491.	2.0	37
22	Genome-Wide RNAi of C. elegans Using the Hypersensitive rrf-3 Strain Reveals Novel Gene Functions. PLoS Biology, 2003, 1, e12.	5.6	545
23	The G-Protein β-Subunit GPB-2 in Caenorhabditis elegans Regulates the Goα-Gqα Signaling Network Through Interactions With the Regulator of G-Protein Signaling Proteins EGL-10 and EAT-16. Genetics, 2001, 158, 221-235.	2.9	56
24	G protein hyperactivation of the Caenorhabditis elegans adenylyl cyclase SGS-1 induces neuronal degeneration. EMBO Journal, 1998, 17, 5059-5065.	7.8	43