
## Margarida Ruas

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/4728389/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	<i>Tpcn2</i> knockout mice have improved insulin sensitivity and are protected against high-fat diet-induced weight gain. Physiological Genomics, 2018, 50, 605-614.	2.3	3
2	Synthesis of the Ca2+-mobilizing messengers NAADP and cADPR by intracellular CD38 enzyme in the mouse heart: Role in β-adrenoceptor signaling. Journal of Biological Chemistry, 2017, 292, 13243-13257.	3.4	44
3	PLCζ is the physiological trigger of the Ca2+ oscillations that induce embryogenesis in mammals but offspring can be conceived in its absence. Development (Cambridge), 2017, 144, 2914-2924.	2.5	95
4	Carvedilol Inhibits cADPR- and IP3-Induced Ca2+ Release. Messenger (Los Angeles, Calif: Print), 2016, 5, 92-99.	0.3	3
5	Ebolavirus Glycoprotein Directs Fusion through NPC1 <sup>+</sup> Endolysosomes. Journal of Virology, 2016, 90, 605-610.	3.4	67
6	Expression of Ca <sup>2+</sup> â€permeable twoâ€pore channels rescues <scp>NAADP</scp> signalling in <scp>TPC</scp> â€deficient cells. EMBO Journal, 2015, 34, 1743-1758.	7.8	144
7	Two-Pore Channels: Lessons from Mutant Mouse Models. Messenger (Los Angeles, Calif: Print), 2015, 4, 4-22.	0.3	22
8	Two-Pore Channel 2 activity is required for slow muscle cell-generated Ca2+ signaling during myogenesis in intact zebrafish. International Journal of Developmental Biology, 2015, 59, 313-325.	0.6	30
9	TPC: the NAADP discovery channel?. Biochemical Society Transactions, 2015, 43, 384-389.	3.4	41
10	Two-pore Channels (TPC2s) and Nicotinic Acid Adenine Dinucleotide Phosphate (NAADP) at Lysosomal-Sarcoplasmic Reticular Junctions Contribute to Acute and Chronic β-Adrenoceptor Signaling in the Heart. Journal of Biological Chemistry, 2015, 290, 30087-30098.	3.4	63
11	Nicotinic Acid Adenine Dinucleotide Phosphate (NAADP) and Endolysosomal Two-pore Channels Modulate Membrane Excitability and Stimulus-Secretion Coupling in Mouse Pancreatic Î <sup>2</sup> Cells. Journal of Biological Chemistry, 2015, 290, 21376-21392.	3.4	48
12	Reply to "TPC1 Knockout Knocks Out TPC1― Molecular and Cellular Biology, 2015, 35, 1884-1884.	2.3	1
13	Synthesis of [ <sup>32</sup> P]NAADP for the Radioreceptor Binding Assay. Cold Spring Harbor Protocols, 2014, 2014, pdb.prot076919.	0.3	1
14	Preparation and Use of Sea Urchin Egg Homogenates for Studying NAADP-Mediated Ca2+ Release. Cold Spring Harbor Protocols, 2014, 2014, pdb.prot076901-pdb.prot076901.	0.3	6
15	VEGF-induced neoangiogenesis is mediated by NAADP and two-pore channel-2–dependent Ca <sup>2+</sup> signaling. Proceedings of the National Academy of Sciences of the United States of America, 2014, 111, E4706-15.	7.1	138
16	Synthesis of Caged NAADP. Cold Spring Harbor Protocols, 2014, 2014, pdb.prot076943-pdb.prot076943.	0.3	0
17	TPC1 Has Two Variant Isoforms, and Their Removal Has Different Effects on Endo-Lysosomal Functions Compared to Loss of TPC2. Molecular and Cellular Biology, 2014, 34, 3981-3992.	2.3	76
18	Synthesis of NAADP-AM as a Membrane-Permeant NAADP Analog. Cold Spring Harbor Protocols, 2014, 2014, pdb.prot076927.	0.3	3

Margarida Ruas

#	Article	IF	CITATIONS
19	Measurement of Luminal pH of Acidic Stores as a Readout for NAADP Action. Cold Spring Harbor Protocols, 2014, 2014, pdb.prot076935.	0.3	1
20	Photoaffinity Labeling of High Affinity Nicotinic Acid Adenine Dinucleotide Phosphate (NAADP)-Binding Proteins in Sea Urchin Egg. Journal of Biological Chemistry, 2012, 287, 2308-2315.	3.4	110
21	Two-pore Channels Form Homo- and Heterodimers. Journal of Biological Chemistry, 2011, 286, 37058-37062.	3.4	51
22	Loss of activity mutations in phospholipase C zeta (PLCÂ) abolishes calcium oscillatory ability of human recombinant protein in mouse oocytes. Human Reproduction, 2011, 26, 3372-3387.	0.9	75
23	NAADP as an intracellular messenger regulating lysosomal calcium-release channels. Biochemical Society Transactions, 2010, 38, 1424-1431.	3.4	91
24	Purified TPC Isoforms Form NAADP Receptors with Distinct Roles for Ca2+ Signaling and Endolysosomal Trafficking. Current Biology, 2010, 20, 703-709.	3.9	234
25	TPC2 Is a Novel NAADP-sensitive Ca2+ Release Channel, Operating as a Dual Sensor of Luminal pH and Ca2+. Journal of Biological Chemistry, 2010, 285, 35039-35046.	3.4	197
26	TPC2 Proteins Mediate Nicotinic Acid Adenine Dinucleotide Phosphate (NAADP)- and Agonist-evoked Contractions of Smooth Muscle. Journal of Biological Chemistry, 2010, 285, 24925-24932.	3.4	71
27	NAADP mobilizes calcium from acidic organelles through two-pore channels. Nature, 2009, 459, 596-600.	27.8	687
28	Ca2+ Signaling Occurs via Second Messenger Release from Intraorganelle Synthesis Sites. Current Biology, 2008, 18, 1612-1618.	3.9	61
29	A <i>CDKN2A</i> Mutation in Familial Melanoma that Abrogates Binding of p16INK4a to CDK4 but not CDK6. Cancer Research, 2007, 67, 9134-9141.	0.9	47
30	CDK4 and CDK6 Delay Senescence by Kinase-Dependent and p16 INK4a -Independent Mechanisms. Molecular and Cellular Biology, 2007, 27, 4273-4282.	2.3	52
31	NAADP receptors. Cell Calcium, 2005, 38, 273-280.	2.4	51
32	Methods in Cyclic ADP-Ribose and NAADP Research. , 2005, , 265-334.		4
33	EMSY Links the BRCA2 Pathway to Sporadic Breast and Ovarian Cancer. Cell, 2003, 115, 523-535.	28.9	389
34	INK4a-deficient human diploid fibroblasts are resistant to RAS-induced senescence. EMBO Journal, 2002, 21, 2936-2945.	7.8	173
35	Functional evaluation of tumour-specific variants of p16INK4a/CDKN2A: correlation with protein structure information. Oncogene, 1999, 18, 5423-5434.	5.9	69
36	The p16INK4a/CDKN2A tumor suppressor and its relatives. Biochimica Et Biophysica Acta: Reviews on Cancer, 1998, 1378, F115-F177.	7.4	488

#	Article	IF	CITATIONS
37	Isolation and prevalidation of an Escherichia coli tester strain for the use in mechanistic and metabolic studies of genotoxins. Mutation Research - Environmental Mutagenesis and Related Subjects Including Methodology, 1994, 312, 99-109.	0.4	10