

# Hua Qu

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4739238/publications.pdf>

Version: 2024-02-01

17  
papers

537  
citations

759233

12  
h-index

888059

17  
g-index

17  
all docs

17  
docs citations

17  
times ranked

874  
citing authors

#	ARTICLE	IF	CITATIONS
1	Plasma Secretagoin is Increased in Individuals with Glucose Dysregulation. <i>Experimental and Clinical Endocrinology and Diabetes</i> , 2021, 129, 661-665.	1.2	3
2	The potential effects of clinical antidiabetic agents on <sc>SARSâ€CoV</sc>â€2. <i>Journal of Diabetes</i> , 2021, 13, 243-252.	1.8	15
3	Dedicator of Cytokines 5 Regulates Keratinocyte Function and Promotes Diabetic Wound Healing. <i>Diabetes</i> , 2021, 70, 1170-1184.	0.6	11
4	Deficiency of Mitochondrial Glycerol 3-Phosphate Dehydrogenase Exacerbates Podocyte Injury and the Progression of Diabetic Kidney Disease. <i>Diabetes</i> , 2021, 70, 1372-1387.	0.6	18
5	PSA controls hepatic lipid metabolism by regulating the NRF2 signaling pathway. <i>Journal of Molecular Cell Biology</i> , 2021, 13, 527-539.	3.3	8
6	Pituitary P62 deficiency leads to female infertility by impairing luteinizing hormone production. <i>Experimental and Molecular Medicine</i> , 2021, 53, 1238-1249.	7.7	9
7	Decreased Plasma Maresin 1 Concentration Is Associated with Diabetic Foot Ulcer. <i>Mediators of Inflammation</i> , 2020, 2020, 1-7.	3.0	16
8	A High Level of Circulating Valine Is a Biomarker for Type 2 Diabetes and Associated with the Hypoglycemic Effect of Sitagliptin. <i>Mediators of Inflammation</i> , 2019, 2019, 1-7.	3.0	16
9	Alteration of gut microbiota induced by DPP-4i treatment improves glucose homeostasis. <i>EBioMedicine</i> , 2019, 44, 665-674.	6.1	66
10	Deficiency of Mitochondrial Glycerol 3â€Phosphate Dehydrogenase Contributes to Hepatic Steatosis. <i>Hepatology</i> , 2019, 70, 84-97.	7.3	30
11	Mitochondrial glycerol 3â€phosphate dehydrogenase promotes skeletal muscle regeneration. <i>EMBO Molecular Medicine</i> , 2018, 10, .	6.9	24
12	Plasma Asprosin Levels Are Associated with Glucose Metabolism, Lipid, and Sex Hormone Profiles in Females with Metabolic-Related Diseases. <i>Mediators of Inflammation</i> , 2018, 2018, 1-12.	3.0	71
13	DPP-4 Inhibitors as Treatments for Type 1 Diabetes Mellitus: A Systematic Review and Meta-Analysis. <i>Journal of Diabetes Research</i> , 2018, 2018, 1-10.	2.3	30
14	Plasma Asprosin Concentrations Are Increased in Individuals with Glucose Dysregulation and Correlated with Insulin Resistance and First-Phase Insulin Secretion. <i>Mediators of Inflammation</i> , 2018, 2018, 1-7.	3.0	98
15	Global and Regional Effects of Bladder Cancer Risk Associated with Pioglitazone Therapy in Patients with Diabetes. <i>Scientific Reports</i> , 2017, 7, 15804.	3.3	13
16	Plasma CTRP-3 concentrations in Chinese patients with obesity and type II diabetes negatively correlate with insulin resistance. <i>Journal of Clinical Lipidology</i> , 2015, 9, 289-294.	1.5	43
17	Plasma Progranulin Concentrations Are Increased in Patients with Type 2 Diabetes and Obesity and Correlated with Insulin Resistance. <i>Mediators of Inflammation</i> , 2013, 2013, 1-6.	3.0	66