

# Jonathan D G Jones

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4766845/publications.pdf>

Version: 2024-02-01

352  
papers

87,274  
citations

613

124  
h-index

361

282  
g-index

395  
all docs

395  
docs citations

395  
times ranked

43137  
citing authors

#	ARTICLE	IF	CITATIONS
1	The Arabidopsis <i>WRR4A</i> and <i>WRR4B</i> paralogous NLR proteins both confer recognition of multiple <i>Albugo candida</i> effectors. <i>New Phytologist</i> , 2023, 237, 532-547.	3.5	7
2	An Improved Assembly of the <i>Albugo candida</i> Ac2V Genome Reveals the Expansion of the "CCG" Class of Effectors. <i>Molecular Plant-Microbe Interactions</i> , 2022, 35, 39-48.	1.4	6
3	Plant immune networks. <i>Trends in Plant Science</i> , 2022, 27, 255-273.	4.3	140
4	Thirty years of resistance: Zig-zag through the plant immune system. <i>Plant Cell</i> , 2022, 34, 1447-1478.	3.1	318
5	<i>Aegilops sharonensis</i> genome-assisted identification of stem rust resistance gene Sr62. <i>Nature Communications</i> , 2022, 13, 1607.	5.8	48
6	The Ry <sub>sto</sub> immune receptor recognises a broadly conserved feature of potyviral coat proteins. <i>New Phytologist</i> , 2022, 235, 1179-1195.	3.5	10
7	The host exocyst complex is targeted by a conserved bacterial type-III effector that promotes virulence. <i>Plant Cell</i> , 2022, 34, 3400-3424.	3.1	17
8	Identification of RipAZ1 as an avirulence determinant of <i>Ralstonia solanacearum</i> in <i>Solanum americanum</i> . <i>Molecular Plant Pathology</i> , 2021, 22, 317-333.	2.0	15
9	A complex resistance locus in <i>Solanum americanum</i> recognizes a conserved <i>Phytophthora</i> effector. <i>Nature Plants</i> , 2021, 7, 198-208.	4.7	62
10	Transient reprogramming of crop plants for agronomic performance. <i>Nature Plants</i> , 2021, 7, 159-171.	4.7	72
11	Mutual potentiation of plant immunity by cell-surface and intracellular receptors. <i>Nature</i> , 2021, 592, 110-115.	13.7	536
12	New Honorary Member of the BSPP. <i>Plant Pathology</i> , 2021, 70, 763-763.	1.2	0
13	Pathogen effector recognition-dependent association of NRG1 with EDS1 and SAG101 in TNL receptor immunity. <i>Nature Communications</i> , 2021, 12, 3335.	5.8	112
14	Channeling plant immunity. <i>Cell</i> , 2021, 184, 3358-3360.	13.5	14
15	Chromatin accessibility landscapes activated by cell-surface and intracellular immune receptors. <i>Journal of Experimental Botany</i> , 2021, 72, 7927-7941.	2.4	14
16	Evolutionary tradeoffs at the Arabidopsis <i>WRR4A</i> resistance locus underpin alternate <i>Albugo candida</i> race recognition specificities. <i>Plant Journal</i> , 2021, 107, 1490-1502.	2.8	5
17	Evolutionarily distinct resistance proteins detect a pathogen effector through its association with different host targets. <i>New Phytologist</i> , 2021, 232, 1368-1381.	3.5	6
18	Autoactive Arabidopsis RPS4 alleles require partner protein RRS1-R. <i>Plant Physiology</i> , 2021, 185, 761-764.	2.3	7

#	ARTICLE	IF	CITATIONS
19	Perception of structurally distinct effectors by the integrated WRKY domain of a plant immune receptor. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2021, 118, .	3.3	32
20	Extreme resistance to <i>Potato virus Y</i> in potato carrying the <i>Ry</i> gene is mediated by a TIR-NLR immune receptor. <i>Plant Biotechnology Journal</i> , 2020, 18, 655-667.	4.1	57
21	High-resolution expression profiling of selected gene sets during plant immune activation. <i>Plant Biotechnology Journal</i> , 2020, 18, 1610-1619.	4.1	21
22	RNA Splicing: A Novel Pathogen Effector Target. <i>Molecular Plant</i> , 2020, 13, 1348.	3.9	0
23	Identification of <i>Avramr1</i> from <i>Phytophthora infestans</i> using long read and cDNA pathogen-enrichment sequencing (PenSeq). <i>Molecular Plant Pathology</i> , 2020, 21, 1502-1512.	2.0	22
24	Induced proximity of a TIR signaling domain on a plant-mammalian NLR chimera activates defense in plants. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2020, 117, 18832-18839.	3.3	82
25	Two unequally redundant "helper" immune receptor families mediate <i>Arabidopsis thaliana</i> intracellular "sensor" immune receptor functions. <i>PLoS Biology</i> , 2020, 18, e3000783.	2.6	125
26	The NLR-Annotator Tool Enables Annotation of the Intracellular Immune Receptor Repertoire. <i>Plant Physiology</i> , 2020, 183, 468-482.	2.3	147
27	Breeding a fungal gene into wheat. <i>Science</i> , 2020, 368, 822-823.	6.0	8
28	Plant NLRs get by with a little help from their friends. <i>Current Opinion in Plant Biology</i> , 2020, 56, 99-108.	3.5	70
29	Estradiol-inducible <i>AvrRps4</i> expression reveals distinct properties of TIR-NLR-mediated effector-triggered immunity. <i>Journal of Experimental Botany</i> , 2020, 71, 2186-2197.	2.4	37
30	Phosphorylation-Regulated Activation of the <i>Arabidopsis</i> RRS1-R/RPS4 Immune Receptor Complex Reveals Two Distinct Effector Recognition Mechanisms. <i>Cell Host and Microbe</i> , 2020, 27, 769-781.e6.	5.1	50
31	Title is missing!. , 2020, 18, e3000783.		0
32	Title is missing!. , 2020, 18, e3000783.		0
33	Title is missing!. , 2020, 18, e3000783.		0
34	Title is missing!. , 2020, 18, e3000783.		0
35	Title is missing!. , 2020, 18, e3000783.		0
36	Title is missing!. , 2020, 18, e3000783.		0

#	ARTICLE	IF	CITATIONS
37	Using CRISPR/Cas9 genome editing in tomato to create a gibberellin-responsive dominant dwarf DELLA allele. <i>Plant Biotechnology Journal</i> , 2019, 17, 132-140.	4.1	64
38	A Species-Wide Inventory of NLR Genes and Alleles in <i>Arabidopsis thaliana</i> . <i>Cell</i> , 2019, 178, 1260-1272.e14.	13.5	265
39	A SWEET solution to rice blight. <i>Nature Biotechnology</i> , 2019, 37, 1280-1282.	9.4	20
40	The curious case of the bacterial engineer. <i>Nature Plants</i> , 2019, 5, 906-907.	4.7	0
41	Transgressive segregation reveals mechanisms of <i>Arabidopsis</i> immunity to <i>Brassica</i> -infecting races of white rust ( <i>Albugo candida</i> ). <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2019, 116, 2767-2773.	3.3	57
42	Floriculture: Ellis and Dodds's™ Illumination of Gene-for-Gene Biology. <i>Plant Cell</i> , 2019, 31, 1204-1205.	3.1	3
43	A pentangular plant inflammasome. <i>Science</i> , 2019, 364, 31-32.	6.0	28
44	Alien domains shaped the modular structure of plant NLR proteins. <i>Genome Biology and Evolution</i> , 2019, 11, 3466-3477.	1.1	21
45	Diverse NLR immune receptors activate defence via the RPW8-NLR NRG1. <i>New Phytologist</i> , 2019, 222, 966-980.	3.5	219
46	Optimization of T-DNA architecture for Cas9-mediated mutagenesis in <i>Arabidopsis</i> . <i>PLoS ONE</i> , 2019, 14, e0204778.	1.1	96
47	<i>Albugo candida</i> race diversity, ploidy and host-associated microbes revealed using DNA sequence capture on diseased plants in the field. <i>New Phytologist</i> , 2019, 221, 1529-1543.	3.5	41
48	Autoimmunity and effector recognition in <i>Arabidopsis thaliana</i> can be uncoupled by mutations in the RRS1 immune receptor. <i>New Phytologist</i> , 2019, 222, 954-965.	3.5	10
49	Pathogen enrichment sequencing (PenSeq) enables population genomic studies in oomycetes. <i>New Phytologist</i> , 2019, 221, 1634-1648.	3.5	43
50	Resistance gene cloning from a wild crop relative by sequence capture and association genetics. <i>Nature Biotechnology</i> , 2019, 37, 139-143.	9.4	280
51	Pm21 from <i>Haynaldia villosa</i> Encodes a CC-NBS-LRR Protein Conferring Powdery Mildew Resistance in Wheat. <i>Molecular Plant</i> , 2018, 11, 874-878.	3.9	181
52	A downy mildew effector evades recognition by polymorphism of expression and subcellular localization. <i>Nature Communications</i> , 2018, 9, 5192.	5.8	40
53	Distinct modes of derepression of an <i>Arabidopsis</i> immune receptor complex by two different bacterial effectors. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2018, 115, 10218-10227.	3.3	83
54	<i>Arabidopsis</i> downy mildew effector HaRxL106 suppresses plant immunity by binding to RADICAL-INDUCED CELL DEATH1. <i>New Phytologist</i> , 2018, 220, 232-248.	3.5	51

#	ARTICLE	IF	CITATIONS
55	A workflow for simplified analysis of ATAC-cap-seq data in R. <i>GigaScience</i> , 2018, 7, .	3.3	6
56	The transcriptional landscape of polyploid wheat. <i>Science</i> , 2018, 361, .	6.0	768
57	Shifting the limits in wheat research and breeding using a fully annotated reference genome. <i>Science</i> , 2018, 361, .	6.0	2,424
58	Deadlier than the malate. <i>Cell Research</i> , 2018, 28, 609-610.	5.7	1
59	<i>Arabidopsis</i> late blight: infection of a nonhost plant by <i>Albugo laibachii</i> enables full colonization by <i>Phytophthora infestans</i> . <i>Cellular Microbiology</i> , 2017, 19, e12628.	1.1	44
60	Mis-placed Congeniality: When Pathogens Ask Their Plant Hosts for Another Drink. <i>Developmental Cell</i> , 2017, 40, 116-117.	3.1	2
61	Two-faced TIRs trip the immune switch. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2017, 114, 2445-2446.	3.3	1
62	Genomic Rearrangements in <i>Arabidopsis</i> Considered as Quantitative Traits. <i>Genetics</i> , 2017, 205, 1425-1441.	1.2	21
63	Albugo-imposed changes to tryptophan-derived antimicrobial metabolite biosynthesis may contribute to suppression of non-host resistance to <i>Phytophthora infestans</i> in <i>Arabidopsis thaliana</i> . <i>BMC Biology</i> , 2017, 15, 20.	1.7	48
64	MutRenSeq: A Method for Rapid Cloning of Plant Disease Resistance Genes. <i>Methods in Molecular Biology</i> , 2017, 1659, 215-229.	0.4	22
65	Discovery and characterization of two new stem rust resistance genes in <i>Aegilops sharonensis</i> . <i>Theoretical and Applied Genetics</i> , 2017, 130, 1207-1222.	1.8	45
66	The highly buffered <i>Arabidopsis</i> immune signaling network conceals the functions of its components. <i>PLoS Genetics</i> , 2017, 13, e1006639.	1.5	138
67	Comparative analysis of targeted long read sequencing approaches for characterization of a plant's immune receptor repertoire. <i>BMC Genomics</i> , 2017, 18, 564.	1.2	51
68	Protein-protein interactions in the RPS4/RRS1 immune receptor complex. <i>PLoS Pathogens</i> , 2017, 13, e1006376.	2.1	103
69	Targeted capture and sequencing of gene-sized DNA molecules. <i>BioTechniques</i> , 2016, 61, 315-322.	0.8	48
70	Pathogen perception by NLRs in plants and animals: Parallel worlds. <i>BioEssays</i> , 2016, 38, 769-781.	1.2	81
71	Intracellular innate immune surveillance devices in plants and animals. <i>Science</i> , 2016, 354, .	6.0	834
72	Comparative analysis of plant immune receptor architectures uncovers host proteins likely targeted by pathogens. <i>BMC Biology</i> , 2016, 14, 8.	1.7	293

#	ARTICLE	IF	CITATIONS
73	Accelerated cloning of a potato late blight resistance gene using RenSeq and SMRT sequencing. <i>Nature Biotechnology</i> , 2016, 34, 656-660.	9.4	248
74	Rapid cloning of disease-resistance genes in plants using mutagenesis and sequence capture. <i>Nature Biotechnology</i> , 2016, 34, 652-655.	9.4	383
75	A pigeonpea gene confers resistance to Asian soybean rust in soybean. <i>Nature Biotechnology</i> , 2016, 34, 661-665.	9.4	87
76	Characterization of a <i>JAZ7</i> activation-tagged <i>Arabidopsis</i> mutant with increased susceptibility to the fungal pathogen <i>Fusarium oxysporum</i> . <i>Journal of Experimental Botany</i> , 2016, 67, 2367-2386.	2.4	68
77	Standards for plant synthetic biology: a common syntax for exchange of DNA parts. <i>New Phytologist</i> , 2015, 208, 13-19.	3.5	263
78	Probing formation of cargo/importin transport complexes in plant cells using a pathogen effector. <i>Plant Journal</i> , 2015, 81, 40-52.	2.8	48
79	Autoimmunity conferred by <i>chs3-2D</i> relies on <i>CSA1</i> , its adjacent TNL-encoding neighbour. <i>Scientific Reports</i> , 2015, 5, 8792.	1.6	47
80	Comparative genomic analysis of multiple strains of two unusual plant pathogens: <i>Pseudomonas corrugata</i> and <i>Pseudomonas mediterranea</i> . <i>Frontiers in Microbiology</i> , 2015, 6, 811.	1.5	50
81	A Plant Immune Receptor Detects Pathogen Effectors that Target WRKY Transcription Factors. <i>Cell</i> , 2015, 161, 1089-1100.	13.5	454
82	Fine mapping of the <i>Rpi-rzc1</i> gene conferring broad-spectrum resistance to potato late blight. <i>European Journal of Plant Pathology</i> , 2015, 143, 193-198.	0.8	14
83	NLR-parser: rapid annotation of plant NLR complements. <i>Bioinformatics</i> , 2015, 31, 1665-1667.	1.8	103
84	Two linked pairs of <i>Arabidopsis</i> TNL resistance genes independently confer recognition of bacterial effector <i>AvrRps4</i> . <i>Nature Communications</i> , 2015, 6, 6338.	5.8	147
85	Domestication: Sweet! A naturally transgenic crop. <i>Nature Plants</i> , 2015, 1, 15077.	4.7	0
86	The Top 10 oomycete pathogens in molecular plant pathology. <i>Molecular Plant Pathology</i> , 2015, 16, 413-434.	2.0	695
87	Plant immune receptors mimic pathogen virulence targets. <i>Oncotarget</i> , 2015, 6, 16824-16825.	0.8	4
88	<i>Hyaloperonospora arabidopsidis</i> (Downy Mildew) infection Assay in <i>Arabidopsis</i> . <i>Bio-protocol</i> , 2015, 5, .	0.2	11
89	Evidence for suppression of immunity as a driver for genomic introgressions and host range expansion in races of <i>Albugo candida</i> , a generalist parasite. <i>ELife</i> , 2015, 4, .	2.8	71
90	A novel approach for multi-domain and multi-gene family identification provides insights into evolutionary dynamics of disease resistance genes in core eudicot plants. <i>BMC Genomics</i> , 2014, 15, 966.	1.2	29

#	ARTICLE	IF	CITATIONS
91	The Nuclear Immune Receptor RPS4 Is Required for RRS1SLH1-Dependent Constitutive Defense Activation in <i>Arabidopsis thaliana</i> . <i>PLoS Genetics</i> , 2014, 10, e1004655.	1.5	121
92	Expression Profiling during Arabidopsis/Downy Mildew Interaction Reveals a Highly-Expressed Effector That Attenuates Responses to Salicylic Acid. <i>PLoS Pathogens</i> , 2014, 10, e1004443.	2.1	117
93	The Plasmodesmal Protein PDL1 Localises to Haustoria-Associated Membranes during Downy Mildew Infection and Regulates Callose Deposition. <i>PLoS Pathogens</i> , 2014, 10, e1004496.	2.1	130
94	Genomic DNA Library Preparation for Resistance Gene Enrichment and Sequencing (RenSeq) in Plants. <i>Methods in Molecular Biology</i> , 2014, 1127, 291-303.	0.4	24
95	Elevating crop disease resistance with cloned genes. <i>Philosophical Transactions of the Royal Society B: Biological Sciences</i> , 2014, 369, 20130087.	1.8	117
96	Structural Basis for Assembly and Function of a Heterodimeric Plant Immune Receptor. <i>Science</i> , 2014, 344, 299-303.	6.0	300
97	A locus conferring effective late blight resistance in potato cultivar Solanum Mira maps to chromosome XI. <i>Theoretical and Applied Genetics</i> , 2014, 127, 647-657.	1.8	28
98	Identification of unique SUN-interacting nuclear envelope proteins with diverse functions in plants. <i>Journal of Cell Biology</i> , 2014, 205, 677-692.	2.3	78
99	Convergent Targeting of a Common Host Protein-Network by Pathogen Effectors from Three Kingdoms of Life. <i>Cell Host and Microbe</i> , 2014, 16, 364-375.	5.1	367
100	EXPRSS: an Illumina based high-throughput expression-profiling method to reveal transcriptional dynamics. <i>BMC Genomics</i> , 2014, 15, 341.	1.2	36
101	Defining the full tomato NB-LRR resistance gene repertoire using genomic and cDNA RenSeq. <i>BMC Plant Biology</i> , 2014, 14, 120.	1.6	161
102	Direct Regulation of the NADPH Oxidase RBOHD by the PRR-Associated Kinase BIK1 during Plant Immunity. <i>Molecular Cell</i> , 2014, 54, 43-55.	4.5	744
103	A Golden Gate Modular Cloning Toolbox for Plants. <i>ACS Synthetic Biology</i> , 2014, 3, 839-843.	1.9	666
104	Targeted mutagenesis in the model plant <i>Nicotiana benthamiana</i> using Cas9 RNA-guided endonuclease. <i>Nature Biotechnology</i> , 2013, 31, 691-693.	9.4	951
105	Resistance gene enrichment sequencing (R-Seq) enables reannotation of the NB-LRR gene family from sequenced plant genomes and rapid mapping of resistance loci in segregating populations. <i>Plant Journal</i> , 2013, 76, 530-544.	2.8	367
106	Anthocyanins Double the Shelf Life of Tomatoes by Delaying Overripening and Reducing Susceptibility to Gray Mold. <i>Current Biology</i> , 2013, 23, 1094-1100.	1.8	292
107	The Variable Domain of a Plant Calcium-dependent Protein Kinase (CDPK) Confers Subcellular Localization and Substrate Recognition for NADPH Oxidase. <i>Journal of Biological Chemistry</i> , 2013, 288, 14332-14340.	1.6	129
108	Deployment of the <i>Burkholderia glumae</i> type III secretion system as an efficient tool for translocating pathogen effectors to monocot cells. <i>Plant Journal</i> , 2013, 74, 701-712.	2.8	45

#	ARTICLE	IF	CITATIONS
109	In Planta Effector Competition Assays Detect <i>Hyaloperonospora arabidopsidis</i> Effectors That Contribute to Virulence and Localize to Different Plant Subcellular Compartments. <i>Molecular Plant-Microbe Interactions</i> , 2013, 26, 745-757.	1.4	16
110	Regulation of Transcription of Nucleotide-Binding Leucine-Rich Repeat-Encoding Genes SNC1 and RPP4 via H3K4 Trimethylation. <i>Plant Physiology</i> , 2013, 162, 1694-1705.	2.3	93
111	A Downy Mildew Effector Attenuates Salicylic Acid-Triggered Immunity in Arabidopsis by Interacting with the Host Mediator Complex. <i>PLoS Biology</i> , 2013, 11, e1001732.	2.6	167
112	Crystallization and preliminary X-ray diffraction analyses of the TIR domains of three TIR-NB-LRR proteins that are involved in disease resistance in <i>Arabidopsis thaliana</i> . <i>Acta Crystallographica Section F: Structural Biology Communications</i> , 2013, 69, 1275-1280.	0.7	5
113	Identifying and Classifying Trait Linked Polymorphisms in Non-Reference Species by Walking Coloured de Bruijn Graphs. <i>PLoS ONE</i> , 2013, 8, e60058.	1.1	26
114	Mechanisms of Nuclear Suppression of Host Immunity by Effectors from the Arabidopsis Downy Mildew Pathogen <i>Hyaloperonospora arabidopsidis</i> (Hpa). <i>Cold Spring Harbor Symposia on Quantitative Biology</i> , 2012, 77, 285-293.	2.0	17
115	Draft Genome Sequence of <i>Pseudomonas syringae</i> Pathovar <i>Syringae</i> Strain FF5, Causal Agent of Stem Tip Dieback Disease on Ornamental Pear. <i>Journal of Bacteriology</i> , 2012, 194, 3733-3734.	1.0	10
116	Characterization of the membrane-associated HaRxL17Hpaeffector candidate. <i>Plant Signaling and Behavior</i> , 2012, 7, 145-149.	1.2	7
117	Coverage-based consensus calling (CbCC) of short sequence reads and comparison of CbCC results to identify SNPs in chickpea ( <i>Cicer arietinum</i> ; Fabaceae), a crop species without a reference genome. <i>American Journal of Botany</i> , 2012, 99, 186-192.	0.8	34
118	The <i>awr</i> Gene Family Encodes a Novel Class of <i>Ralstonia solanacearum</i> Type III Effectors Displaying Virulence and Avirulence Activities. <i>Molecular Plant-Microbe Interactions</i> , 2012, 25, 941-953.	1.4	66
119	Distinct regions of the <i>Pseudomonas syringae</i> coiled-coil effector AvrRps4 are required for activation of immunity. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2012, 109, 16371-16376.	3.3	81
120	Obligate biotroph parasitism: can we link genomes to lifestyles?. <i>Trends in Plant Science</i> , 2012, 17, 448-457.	4.3	102
121	Identification and localisation of the NB-LRR gene family within the potato genome. <i>BMC Genomics</i> , 2012, 13, 75.	1.2	290
122	HopAS1 recognition significantly contributes to Arabidopsis nonhost resistance to <i>Pseudomonas syringae</i> pathogens. <i>New Phytologist</i> , 2012, 193, 58-66.	3.5	32
123	Subcellular localization of the Hpa RxLR effector repertoire identifies a tonoplast-associated protein HaRxL17 that confers enhanced plant susceptibility. <i>Plant Journal</i> , 2012, 69, 252-265.	2.8	198
124	Subcellular targeting of an evolutionarily conserved plant defensin <i>MtD4</i> determines the outcome of plant-pathogen interaction in transgenic <i>Arabidopsis</i> . <i>Molecular Plant Pathology</i> , 2012, 13, 1032-1046.	2.0	29
125	Why genetically modified crops?. <i>Philosophical Transactions Series A, Mathematical, Physical, and Engineering Sciences</i> , 2011, 369, 1807-1816.	1.6	17
126	Molecular Cloning of ATR5Emoy2 from <i>Hyaloperonospora arabidopsidis</i> , an Avirulence Determinant That Triggers RPP5-Mediated Defense in Arabidopsis. <i>Molecular Plant-Microbe Interactions</i> , 2011, 24, 827-838.	1.4	102



#	ARTICLE	IF	CITATIONS
127	The microRNA miR393 re-directs secondary metabolite biosynthesis away from camalexin and towards glucosinolates. <i>Plant Journal</i> , 2011, 67, 218-231.	2.8	196
128	Hormone Crosstalk in Plant Disease and Defense: More Than Just JASMONATE-SALICYLATE Antagonism. <i>Annual Review of Phytopathology</i> , 2011, 49, 317-343.	3.5	1,564
129	Crystallization and preliminary X-ray analysis of the RXLR-type effector RXLR3 from the oomycete pathogen <i>Hyaloperonospora arabidopsidis</i> . <i>Acta Crystallographica Section F: Structural Biology Communications</i> , 2011, 67, 1417-1420.	0.7	2
130	Gene Gain and Loss during Evolution of Obligate Parasitism in the White Rust Pathogen of <i>Arabidopsis thaliana</i> . <i>PLoS Biology</i> , 2011, 9, e1001094.	2.6	271
131	Multiple Candidate Effectors from the Oomycete Pathogen <i>Hyaloperonospora arabidopsidis</i> Suppress Host Plant Immunity. <i>PLoS Pathogens</i> , 2011, 7, e1002348.	2.1	212
132	Genome-wide sequencing data reveals virulence factors implicated in banana <i>Xanthomonas</i> wilt. <i>FEMS Microbiology Letters</i> , 2010, 310, 182-192.	0.7	57
133	Genome-wide association study of 107 phenotypes in <i>Arabidopsis thaliana</i> inbred lines. <i>Nature</i> , 2010, 465, 627-631.	13.7	1,651
134	Interfamily transfer of a plant pattern-recognition receptor confers broad-spectrum bacterial resistance. <i>Nature Biotechnology</i> , 2010, 28, 365-369.	9.4	464
135	Signatures of Adaptation to Obligate Biotrophy in the <i>Hyaloperonospora arabidopsidis</i> Genome. <i>Science</i> , 2010, 330, 1549-1551.	6.0	492
136	Genome-wide survey of <i>Arabidopsis</i> natural variation in downy mildew resistance using combined association and linkage mapping. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2010, 107, 10302-10307.	3.3	120
137	Specific ER quality control components required for biogenesis of the plant innate immune receptor EFR. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2009, 106, 15973-15978.	3.3	241
138	<i>Rpi-vnt1.1</i> , a <i>Tm-2<sup>2</sup></i> Homolog from <i>Solanum venturii</i> , Confers Resistance to Potato Late Blight. <i>Molecular Plant-Microbe Interactions</i> , 2009, 22, 589-600.	1.4	194
139	Hormone (Dis)harmony Moulds Plant Health and Disease. <i>Science</i> , 2009, 324, 750-752.	6.0	416
140	Role of plant hormones in plant defence responses. <i>Plant Molecular Biology</i> , 2009, 69, 473-488.	2.0	2,187
141	Two distinct potato late blight resistance genes from <i>Solanum tuberosum</i> are located on chromosome 10. <i>Euphytica</i> , 2009, 165, 269-278.	0.6	37
142	A new resistance gene to powdery mildew identified in <i>Solanum neorossii</i> has been localized on the short arm of potato chromosome 6. <i>Euphytica</i> , 2009, 166, 331-339.	0.6	5
143	The <i>Pseudomonas syringae</i> effector protein, AvrRPS4, requires <i>in planta</i> processing and the KRVY domain to function. <i>Plant Journal</i> , 2009, 57, 1079-1091.	2.8	60
144	Control of the pattern-recognition receptor EFR by an ER protein complex in plant immunity. <i>EMBO Journal</i> , 2009, 28, 3428-3438.	3.5	267

#	ARTICLE	IF	CITATIONS
145	Genome sequence and analysis of the Irish potato famine pathogen <i>Phytophthora infestans</i> . <i>Nature</i> , 2009, 461, 393-398.	13.7	1,405
146	Application of 'next-generation' sequencing technologies to microbial genetics. <i>Nature Reviews Microbiology</i> , 2009, 7, 96-97.	13.6	269
147	In the News. <i>Nature Reviews Microbiology</i> , 2009, 7, 260-261.	13.6	158
148	<i>De novo</i> assembly of the <i>Pseudomonas syringae</i> pv. <i>syringae</i> B728a genome using Illumina/Solexa short sequence reads. <i>FEMS Microbiology Letters</i> , 2009, 291, 103-111.	0.7	77
149	A Biotic or Abiotic Stress?. , 2009, , 103-122.		1
150	The Major Specificity-Determining Amino Acids of the Tomato Cf-9 Disease Resistance Protein Are at Hypervariable Solvent-Exposed Positions in the Central Leucine-Rich Repeats. <i>Molecular Plant-Microbe Interactions</i> , 2009, 22, 1203-1213.	1.4	46
151	Autophagic Components Contribute to Hypersensitive Cell Death in Arabidopsis. <i>Cell</i> , 2009, 137, 773-783.	13.5	348
152	Mapping and Cloning of Late Blight Resistance Genes from <i>Solanum venturii</i> Using an Interspecific Candidate Gene Approach. <i>Molecular Plant-Microbe Interactions</i> , 2009, 22, 601-615.	1.4	148
153	A Draft Genome Sequence of <i>Pseudomonas syringae</i> pv. <i>tomato</i> T1 Reveals a Type III Effector Repertoire Significantly Divergent from That of <i>Pseudomonas syringae</i> pv. <i>tomato</i> DC3000. <i>Molecular Plant-Microbe Interactions</i> , 2009, 22, 52-62.	1.4	134
154	The TIR Domain of TIR-NB-LRR Resistance Proteins Is a Signaling Domain Involved in Cell Death Induction. <i>Molecular Plant-Microbe Interactions</i> , 2009, 22, 157-165.	1.4	185
155	Regions of the Cf-9B Disease Resistance Protein Able to Cause Spontaneous Necrosis in <i>Nicotiana benthamiana</i> Lie Within the Region Controlling Pathogen Recognition in Tomato. <i>Molecular Plant-Microbe Interactions</i> , 2009, 22, 1214-1226.	1.4	17
156	A new species of <i>Albugo</i> parasitic to <i>Arabidopsis thaliana</i> reveals new evolutionary patterns in white blister rusts ( <i>Albuginaceae</i> ). <i>Persoonia: Molecular Phylogeny and Evolution of Fungi</i> , 2009, 22, 123-128.	1.6	80
157	TECHNICAL ADVANCE: Induction of phenotypic variation by activation of genes harbouring a maize <i>Spm</i> element in their promoter regions using a TnpA-VP16 fusion protein. <i>Plant Journal</i> , 2008, 53, 587-594.	2.8	1
158	The auxin influx carrier LAX3 promotes lateral root emergence. <i>Nature Cell Biology</i> , 2008, 10, 946-954.	4.6	715
159	Characterization of Arabidopsis <i>mur3</i> mutations that result in constitutive activation of defence in petioles, but not leaves. <i>Plant Journal</i> , 2008, 56, 691-703.	2.8	40
160	Plant Pathogen Effectors: Getting Mixed Messages. <i>Current Biology</i> , 2008, 18, R128-R130.	1.8	15
161	DELLAs Control Plant Immune Responses by Modulating the Balance of Jasmonic Acid and Salicylic Acid Signaling. <i>Current Biology</i> , 2008, 18, 650-655.	1.8	614
162	The Downy Mildew Effector Proteins ATR1 and ATR13 Promote Disease Susceptibility in <i>Arabidopsis thaliana</i> . <i>Plant Cell</i> , 2008, 19, 4077-4090.	3.1	247

#	ARTICLE	IF	CITATIONS
163	A Genome-Wide Functional Investigation into the Roles of Receptor-Like Proteins in Arabidopsis. <i>Plant Physiology</i> , 2008, 147, 503-517.	2.3	266
164	The F-Box Protein ACRE189/ACIF1 Regulates Cell Death and Defense Responses Activated during Pathogen Recognition in Tobacco and Tomato. <i>Plant Cell</i> , 2008, 20, 697-719.	3.1	154
165	Roles of Plant Hormones in Plant Resistance and Susceptibility to Pathogens. , 2008, , 1-10.		8
166	Genome-wide patterns of single-feature polymorphism in <i>Arabidopsis thaliana</i> . <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2007, 104, 12057-12062.	3.3	157
167	Marker development for the genetic study of natural variation in <i>Arabidopsis thaliana</i> . <i>Bioinformatics</i> , 2007, 23, 3108-3109.	1.8	4
168	A flagellin-induced complex of the receptor FLS2 and BAK1 initiates plant defence. <i>Nature</i> , 2007, 448, 497-500.	13.7	1,619
169	Nuclear Accumulation of the Arabidopsis Immune Receptor RPS4 Is Necessary for Triggering EDS1-Dependent Defense. <i>Current Biology</i> , 2007, 17, 2023-2029.	1.8	281
170	Pathological hormone imbalances. <i>Current Opinion in Plant Biology</i> , 2007, 10, 372-379.	3.5	513
171	Inducible cell death in plant immunity. <i>Seminars in Cancer Biology</i> , 2007, 17, 166-187.	4.3	98
172	Molecular analysis of Agrobacterium T-DNA integration in tomato reveals a role for left border sequence homology in most integration events. <i>Molecular Genetics and Genomics</i> , 2007, 278, 411-420.	1.0	14
173	Multiple Avirulence Paralogues in Cereal Powdery Mildew Fungi May Contribute to Parasite Fitness and Defeat of Plant Resistance. <i>Plant Cell</i> , 2006, 18, 2402-2414.	3.1	245
174	CITRX thioredoxin is a putative adaptor protein connecting Cf-9 and the ACIK1 protein kinase during the Cf-9/Avr9- induced defence response. <i>FEBS Letters</i> , 2006, 580, 4236-4241.	1.3	48
175	Perception of the Bacterial PAMP EF-Tu by the Receptor EFR Restricts Agrobacterium-Mediated Transformation. <i>Cell</i> , 2006, 125, 749-760.	13.5	1,658
176	Dominant-negative interference with defence signalling by truncation mutations of the tomato Cf-9 disease resistance gene. <i>Plant Journal</i> , 2006, 46, 385-399.	2.8	6
177	The <i>Arabidopsis thaliana</i> TIR-NB-LRR R-protein, RPP1A; protein localization and constitutive activation of defence by truncated alleles in tobacco and Arabidopsis. <i>Plant Journal</i> , 2006, 47, 829-840.	2.8	103
178	The plant immune system. <i>Nature</i> , 2006, 444, 323-329.	13.7	10,939
179	A tomato mutant that shows stunting, wilting, progressive necrosis and constitutive expression of defence genes contains a recombinant Hcr9 gene encoding an autoactive protein. <i>Plant Journal</i> , 2006, 46, 369-384.	2.8	8
180	A Plant miRNA Contributes to Antibacterial Resistance by Repressing Auxin Signaling. <i>Science</i> , 2006, 312, 436-439.	6.0	1,762

#	ARTICLE	IF	CITATIONS
181	Reactive Oxygen Species Signaling in Response to Pathogens. <i>Plant Physiology</i> , 2006, 141, 373-378.	2.3	1,449
182	The E3 Ubiquitin Ligase Activity of Arabidopsis PLANT U-BOX17 and Its Functional Tobacco Homolog ACRE276 Are Required for Cell Death and Defense. <i>Plant Cell</i> , 2006, 18, 1084-1098.	3.1	215
183	Rewiring Mitogen-Activated Protein Kinase Cascade by Positive Feedback Confers Potato Blight Resistance. <i>Plant Physiology</i> , 2006, 140, 681-692.	2.3	79
184	A Bacterial Virulence Protein Suppresses Host Innate Immunity to Cause Plant Disease. <i>Science</i> , 2006, 313, 220-223.	6.0	438
185	The U-Box Protein CMPG1 Is Required for Efficient Activation of Defense Mechanisms Triggered by Multiple Resistance Genes in Tobacco and Tomato. <i>Plant Cell</i> , 2006, 18, 1067-1083.	3.1	195
186	Patterns of Dwarf expression and brassinosteroid accumulation in tomato reveal the importance of brassinosteroid synthesis during fruit development. <i>Plant Journal</i> , 2005, 42, 262-269.	2.8	120
187	Pathogen-induced, NADPH oxidase-derived reactive oxygen intermediates suppress spread of cell death in <i>Arabidopsis thaliana</i> . <i>Nature Genetics</i> , 2005, 37, 1130-1134.	9.4	513
188	<i>Solanum mochiquense</i> chromosome IX carries a novel late blight resistance gene Rpi-moc1. <i>Theoretical and Applied Genetics</i> , 2005, 110, 252-258.	1.8	107
189	Structure-Function Analysis of Cf-9, a Receptor-Like Protein with Extracytoplasmic Leucine-Rich Repeats. <i>Plant Cell</i> , 2005, 17, 1000-1015.	3.1	112
190	Rapid Phosphorylation of a Syntaxin during the Avr9/Cf-9-Race-Specific Signaling Pathway. <i>Plant Physiology</i> , 2005, 138, 2406-2416.	2.3	41
191	Involvement of PPS3 Phosphorylated by Elicitor-Responsive Mitogen-Activated Protein Kinases in the Regulation of Plant Cell Death. <i>Plant Physiology</i> , 2005, 139, 1914-1926.	2.3	59
192	Functional Analysis of Avr9/Cf-9 Rapidly Elicited Genes Identifies a Protein Kinase, ACIK1, That Is Essential for Full Cf-9-Dependent Disease Resistance in Tomato. <i>Plant Cell</i> , 2005, 17, 295-310.	3.1	164
193	Phylogenomic Analysis of the Receptor-Like Proteins of Rice and Arabidopsis. <i>Plant Physiology</i> , 2005, 138, 611-623.	2.3	211
194	Ethylene-mediated cross-talk between calcium-dependent protein kinase and MAPK signaling controls stress responses in plants. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2005, 102, 10736-10741.	3.3	292
195	<i>Cladosporium Avr2</i> Inhibits Tomato Rcr3 Protease Required for Cf-2-Dependent Disease Resistance. <i>Science</i> , 2005, 308, 1783-1786.	6.0	415
196	Arabidopsis Downy Mildew Resistance Gene RPP27 Encodes a Receptor-Like Protein Similar to CLAVATA2 and Tomato Cf-9. <i>Plant Physiology</i> , 2004, 135, 1100-1112.	2.3	52
197	Cold Tolerance, SFR2, and the Legacy of Gary Warren. <i>Plant Cell</i> , 2004, 16, 1955-1957.	3.1	1
198	Genetic Variation at the Tomato Cf-4/Cf-9 Locus Induced by EMS Mutagenesis and Intralocus Recombination. <i>Genetics</i> , 2004, 167, 459-470.	1.2	32

#	ARTICLE	IF	CITATIONS
199	The Transcriptional Innate Immune Response to flg22. Interplay and Overlap with Avr Gene-Dependent Defense Responses and Bacterial Pathogenesis. <i>Plant Physiology</i> , 2004, 135, 1113-1128.	2.3	562
200	Virus-induced gene silencing in <i>Solanum</i> species. <i>Plant Journal</i> , 2004, 39, 264-272.	2.8	200
201	Expression of RPS4 in tobacco induces an AvrRps4-independent HR that requires EDS1, SGT1 and HSP90. <i>Plant Journal</i> , 2004, 40, 213-224.	2.8	135
202	Gene shuffling-generated and natural variants of the tomato resistance gene Cf-9 exhibit different auto-necrosis-inducing activities in <i>Nicotiana</i> species. <i>Plant Journal</i> , 2004, 40, 942-956.	2.8	38
203	CITRX thioredoxin interacts with the tomato Cf-9 resistance protein and negatively regulates defence. <i>EMBO Journal</i> , 2004, 23, 2156-2165.	3.5	122
204	Bacterial disease resistance in <i>Arabidopsis</i> through flagellin perception. <i>Nature</i> , 2004, 428, 764-767.	13.7	1,487
205	The plant proteolytic machinery and its role in defence. <i>Current Opinion in Plant Biology</i> , 2004, 7, 400-407.	3.5	231
206	NADPH oxidase AtrbohD and AtrbohF genes function in ROS-dependent ABA signaling in <i>Arabidopsis</i> . <i>EMBO Journal</i> , 2003, 22, 2623-2633.	3.5	1,474
207	Multi-functional T-DNA/Ds tomato lines designed for gene cloning and molecular and physical dissection of the tomato genome. <i>Plant Molecular Biology</i> , 2003, 51, 83-98.	2.0	23
208	Rapid migration in gel filtration of the Cf-4 and Cf-9 resistance proteins is an intrinsic property of Cf proteins and not because of their association with high-molecular-weight proteins. <i>Plant Journal</i> , 2003, 35, 305-315.	2.8	33
209	Reactive oxygen species produced by NADPH oxidase regulate plant cell growth. <i>Nature</i> , 2003, 422, 442-446.	13.7	1,999
210	<i>Nicotiana benthamiana</i> gp91phox Homologs NbrbohA and NbrbohB Participate in H <sub>2</sub> O <sub>2</sub> Accumulation and Resistance to <i>Phytophthora infestans</i> . <i>Plant Cell</i> , 2003, 15, 706-718.	3.1	573
211	p-Coumaroylnoradrenaline, a Novel Plant Metabolite Implicated in Tomato Defense against Pathogens. <i>Journal of Biological Chemistry</i> , 2003, 278, 43373-43383.	1.6	88
212	ATIDB: <i>Arabidopsis thaliana</i> insertion database. <i>Nucleic Acids Research</i> , 2003, 31, 1245-1251.	6.5	36
213	CDPK-mediated signalling pathways: specificity and cross-talk. <i>Journal of Experimental Botany</i> , 2003, 55, 181-188.	2.4	342
214	Putting Plant Disease Resistance Genes to Work. , 2003, , 10-17.		1
215	Ubiquitin ligase-associated protein SGT1 is required for host and nonhost disease resistance in plants. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2002, 99, 10865-10869.	3.3	385
216	Regulatory Role of SGT1 in Early R Gene-Mediated Plant Defenses. <i>Science</i> , 2002, 295, 2077-2080.	6.0	385

#	ARTICLE	IF	CITATIONS
217	A Tomato Cysteine Protease Required for Cf-2-Dependent Disease Resistance and Suppression of Autonecrosis. <i>Science</i> , 2002, 296, 744-747.	6.0	365
218	Arabidopsis gp91phox homologues AtrbohD and AtrbohF are required for accumulation of reactive oxygen intermediates in the plant defense response. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2002, 99, 517-522.	3.3	1,488
219	Developmental Control of Promoter Activity Is Not Responsible for Mature Onset of Cf-9B-Mediated Resistance to Leaf Mold in Tomato. <i>Molecular Plant-Microbe Interactions</i> , 2002, 15, 1099-1107.	1.4	66
220	Arabidopsis RAR1 Exerts Rate-Limiting Control of R Gene-Mediated Defenses against Multiple Pathogens. <i>Plant Cell</i> , 2002, 14, 979-992.	3.1	197
221	GARNet, the Genomic Arabidopsis Resource Network. <i>Trends in Plant Science</i> , 2002, 7, 145-147.	4.3	10
222	Transposition patterns of unlinked transposed Ds elements from two T-DNA loci on tomato chromosomes 7 and 8. <i>Molecular Genetics and Genomics</i> , 2002, 266, 882-890.	1.0	4
223	Arabidopsis RPP4 is a member of the RPP5 multigene family of TIR-NB-LRR genes and confers downy mildew resistance through multiple signalling components. <i>Plant Journal</i> , 2002, 29, 439-451.	2.8	256
224	Trehalose-6-phosphate synthase 1, which catalyses the first step in trehalose synthesis, is essential for Arabidopsis embryo maturation. <i>Plant Journal</i> , 2002, 29, 225-235.	2.8	333
225	No Evidence for Binding Between Resistance Gene Product Cf-9 of Tomato and Avirulence Gene Product AVR9 of <i>Cladosporium fulvum</i> . <i>Molecular Plant-Microbe Interactions</i> , 2001, 14, 867-876.	1.4	78
226	Domain Swapping and Gene Shuffling Identify Sequences Required for Induction of an Avr-Dependent Hypersensitive Response by the Tomato Cf-4 and Cf-9 Proteins. <i>Plant Cell</i> , 2001, 13, 255.	3.1	2
227	Calcium-dependent protein kinases play an essential role in a plant defence response. <i>EMBO Journal</i> , 2001, 20, 5556-5567.	3.5	476
228	Plant pathogens and integrated defence responses to infection. <i>Nature</i> , 2001, 411, 826-833.	13.7	3,460
229	Putting knowledge of plant disease resistance genes to work. <i>Current Opinion in Plant Biology</i> , 2001, 4, 281-287.	3.5	151
230	Domain Swapping and Gene Shuffling Identify Sequences Required for Induction of an Avr-Dependent Hypersensitive Response by the Tomato Cf-4 and Cf-9 Proteins. <i>Plant Cell</i> , 2001, 13, 255-272.	3.1	116
231	Comparison of the Hypersensitive Response Induced by the Tomato Cf-4 and Cf-9 Genes in <i>Nicotiana spp.</i> . <i>Molecular Plant-Microbe Interactions</i> , 2000, 13, 465-469.	1.4	43
232	Resistance Gene-Dependent Activation of a Calcium-Dependent Protein Kinase in the Plant Defense Response. <i>Plant Cell</i> , 2000, 12, 803.	3.1	5
233	Early signalling events in the Avr9/Cf-9-dependent plant defence response. <i>Molecular Plant Pathology</i> , 2000, 1, 3-8.	2.0	12
234	Unravelling R gene-mediated disease resistance pathways in Arabidopsis. <i>Molecular Plant Pathology</i> , 2000, 1, 17-24.	2.0	35

#	ARTICLE	IF	CITATIONS
235	Dispersion of the Cf-4 disease resistance gene in Lycopersicon germplasm. <i>Heredity</i> , 2000, 85, 266-270.	1.2	10
236	Functional, c-myc-tagged Cf-9 resistance gene products are plasma-membrane localized and glycosylated. <i>Plant Journal</i> , 2000, 21, 529-536.	2.8	51
237	Salicylic acid is not required for Cf-2- and Cf-9-dependent resistance of tomato to <i>Cladosporium fulvum</i> . <i>Plant Journal</i> , 2000, 23, 305-318.	2.8	139
238	Transcriptional repression by AtMYB4 controls production of UV-protecting sunscreens in <i>Arabidopsis</i> . <i>EMBO Journal</i> , 2000, 19, 6150-6161.	3.5	797
239	Resistance Gene-Dependent Activation of a Calcium-Dependent Protein Kinase in the Plant Defense Response. <i>Plant Cell</i> , 2000, 12, 803-815.	3.1	253
240	<i>Arabidopsis</i> Research 2000. <i>Plant Cell</i> , 2000, 12, 2302.	3.1	0
241	cDNA-AFLP Reveals a Striking Overlap in Race-Specific Resistance and Wound Response Gene Expression Profiles. <i>Plant Cell</i> , 2000, 12, 963.	3.1	2
242	cDNA-AFLP Display for the Isolation of <i>Peronospora parasitica</i> Genes Expressed During Infection in <i>Arabidopsis thaliana</i> . <i>Molecular Plant-Microbe Interactions</i> , 2000, 13, 895-898.	1.4	58
243	cDNA-AFLP Reveals a Striking Overlap in Race-Specific Resistance and Wound Response Gene Expression Profiles. <i>Plant Cell</i> , 2000, 12, 963-977.	3.1	387
244	Genetic complexity of pathogen perception by plants: The example of Rcr3, a tomato gene required specifically by Cf-2. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2000, 97, 8807-8814.	3.3	151
245	<i>Arabidopsis</i> RelA/SpoT homologs implicate (p)ppGpp in plant signaling. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2000, 97, 3747-52.	3.3	85
246	Highlights from the Ninth International Congress on Molecular Plant-Microbe Interactions. <i>Plant Cell</i> , 1999, 11, 2063.	3.1	0
247	A Chromodomain Protein Encoded by the <i>Arabidopsis</i> CAO Gene Is a Plant-Specific Component of the Chloroplast Signal Recognition Particle Pathway That Is Involved in LHCP Targeting. <i>Plant Cell</i> , 1999, 11, 87.	3.1	2
248	Rapid Avr9- and Cf-9-Dependent Activation of MAP Kinases in Tobacco Cell Cultures and Leaves: Convergence of Resistance Gene, Elicitor, Wound, and Salicylate Responses. <i>Plant Cell</i> , 1999, 11, 273.	3.1	10
249	Multiple Independent Defective Suppressor-mutator Transposon Insertions in <i>Arabidopsis</i> : A Tool for Functional Genomics. <i>Plant Cell</i> , 1999, 11, 1841-1852.	3.1	353
250	Pronounced Intraspecific Haplotype Divergence at the RPP5 Complex Disease Resistance Locus of <i>Arabidopsis</i> . <i>Plant Cell</i> , 1999, 11, 2099-2111.	3.1	312
251	The tomato DWARF enzyme catalyses C-6 oxidation in brassinosteroid biosynthesis. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1999, 96, 1761-1766.	3.3	306
252	Recombination between diverged clusters of the tomato Cf-9 plant disease resistance gene family. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1999, 96, 5850-5855.	3.3	113

#	ARTICLE	IF	CITATIONS
253	EDS1, an essential component of R gene-mediated disease resistance in Arabidopsis has homology to eukaryotic lipases. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1999, 96, 3292-3297.	3.3	571
254	A Chromodomain Protein Encoded by the Arabidopsis CAO Gene Is a Plant-Specific Component of the Chloroplast Signal Recognition Particle Pathway That Is Involved in LHCP Targeting. <i>Plant Cell</i> , 1999, 11, 87-99.	3.1	142
255	Rapid Avr9- and Cf-9-Dependent Activation of MAP Kinases in Tobacco Cell Cultures and Leaves: Convergence of Resistance Gene, Elicitor, Wound, and Salicylate Responses. <i>Plant Cell</i> , 1999, 11, 273-287.	3.1	458
256	Homologues of the Cf-9 Disease Resistance Gene (Hcr9s) Are Present at Multiple Loci on the Short Arm of Tomato Chromosome 1. <i>Molecular Plant-Microbe Interactions</i> , 1999, 12, 93-102.	1.4	53
257	K <sup>+</sup> channels of Cf-9 transgenic tobacco guard cells as targets for <i>Cladosporium fulvum</i> Avr9 elicitor-dependent signal transduction. <i>Plant Journal</i> , 1999, 19, 453-462.	2.8	79
258	Multiple Independent Defective Suppressor-mutator Transposon Insertions in Arabidopsis: A Tool for Functional Genomics. <i>Plant Cell</i> , 1999, 11, 1841.	3.1	35
259	Pronounced Intraspecific Haplotype Divergence at the RPP5 Complex Disease Resistance Locus of Arabidopsis. <i>Plant Cell</i> , 1999, 11, 2099.	3.1	5
260	A second gene at the tomato Cf-4 locus confers resistance to <i>Cladosporium fulvum</i> through recognition of a novel avirulence determinant. <i>Plant Journal</i> , 1999, 20, 279-288.	2.8	73
261	Analysis of 1.9-Mb of contiguous sequence from chromosome 4 of Arabidopsis thaliana. <i>Nature</i> , 1998, 391, 485-488.	13.7	844
262	The NB-ARC domain: a novel signalling motif shared by plant resistance gene products and regulators of cell death in animals. <i>Current Biology</i> , 1998, 8, R226-R228.	1.8	539
263	Plant disease-resistance proteins and the gene-for-gene concept. <i>Trends in Biochemical Sciences</i> , 1998, 23, 454-456.	3.7	694
264	Systemin triggers an increase of cytoplasmic calcium in tomato mesophyll cells: Ca <sup>2+</sup> mobilization from intra- and extracellular compartments. <i>Plant, Cell and Environment</i> , 1998, 21, 1101-1111.	2.8	89
265	Six Arabidopsis thaliana homologues of the human respiratory burst oxidase (gp91phox). <i>Plant Journal</i> , 1998, 14, 365-370.	2.8	403
266	Genetic and molecular analysis of tomato Cf genes for resistance to <i>Cladosporium fulvum</i> . <i>Philosophical Transactions of the Royal Society B: Biological Sciences</i> , 1998, 353, 1413-1424.	1.8	81
267	The Tomato Cf-9 Disease Resistance Gene Functions in Tobacco and Potato to Confer Responsiveness to the Fungal Avirulence Gene Product Avr9. <i>Plant Cell</i> , 1998, 10, 1251-1266.	3.1	138
268	The Tomato Cf-5 Disease Resistance Gene and Six Homologs Show Pronounced Allelic Variation in Leucine-Rich Repeat Copy Number. <i>Plant Cell</i> , 1998, 10, 1915-1925.	3.1	286
269	Three Genes of the Arabidopsis RPP1 Complex Resistance Locus Recognize Distinct <i>Peronospora parasitica</i> Avirulence Determinants. <i>Plant Cell</i> , 1998, 10, 1847.	3.1	10
270	Transposon Tagging of the Defective embryo and meristems Gene of Tomato. <i>Plant Cell</i> , 1998, 10, 877-887.	3.1	34



#	ARTICLE	IF	CITATIONS
271	Three Genes of the Arabidopsis RPP1 Complex Resistance Locus Recognize Distinct Peronospora parasitica Avirulence Determinants. <i>Plant Cell</i> , 1998, 10, 1847-1860.	3.1	351
272	Transposon Tagging of the Defective embryo and meristems Gene of Tomato. <i>Plant Cell</i> , 1998, 10, 877.	3.1	2
273	The Tomato Cf-5 Disease Resistance Gene and Six Homologs Show Pronounced Allelic Variation in Leucine-Rich Repeat Copy Number. <i>Plant Cell</i> , 1998, 10, 1915.	3.1	17
274	The Tomato Cf-9 Disease Resistance Gene Functions in Tobacco and Potato to Confer Responsiveness to the Fungal Avirulence Gene Product Avr9. <i>Plant Cell</i> , 1998, 10, 1251.	3.1	11
275	Rapid, Cf-9- and Avr9-Dependent Production of Active Oxygen Species in Tobacco Suspension Cultures. <i>Molecular Plant-Microbe Interactions</i> , 1998, 11, 1155-1166.	1.4	118
276	Epigenetic Instability and Trans-Silencing Interactions Associated With an SPT::Ac T-DNA Locus in Tobacco. <i>Genetics</i> , 1998, 148, 457-469.	1.2	14
277	The Arabidopsis downy mildew resistance gene RPP5 shares similarity to the toll and interleukin-1 receptors with N and L6.. <i>Plant Cell</i> , 1997, 9, 879-894.	3.1	434
278	Characterization of the Tomato Cf-4 Gene for Resistance to <i>Cladosporium fulvum</i> Identifies Sequences That Determine Recognition Specificity in Cf-4 and Cf-9. <i>Plant Cell</i> , 1997, 9, 2209.	3.1	67
279	PLANT DISEASE RESISTANCE GENES. <i>Annual Review of Plant Biology</i> , 1997, 48, 575-607.	14.2	965
280	Novel Disease Resistance Specificities Result from Sequence Exchange between Tandemly Repeated Genes at the Cf-4/9 Locus of Tomato. <i>Cell</i> , 1997, 91, 821-832.	13.5	562
281	A kinase with keen eyes. <i>Nature</i> , 1997, 385, 397-398.	13.7	14
282	AtDMC1, the Arabidopsis homologue of the yeast DMC1 gene: characterization, transposon-induced allelic variation and meiosis-associated expression. <i>Plant Journal</i> , 1997, 11, 1-14.	2.8	235
283	Map positions of 47 Arabidopsis sequences with sequence similarity to disease resistance genes. <i>Plant Journal</i> , 1997, 12, 1197-1211.	2.8	102
284	The Tomato Cf-2 Disease Resistance Locus Comprises Two Functional Genes Encoding Leucine-Rich Repeat Proteins. <i>Cell</i> , 1996, 84, 451-459.	13.5	591
285	Resistance Gene-Dependent Plant Defense Responses. <i>Plant Cell</i> , 1996, 8, 1773.	3.1	37
286	Plant disease resistance genes: structure, function and evolution. <i>Current Opinion in Biotechnology</i> , 1996, 7, 155-160.	3.3	37
287	rbohA, a rice homologue of the mammalian gp91phox respiratory burst oxidase gene. <i>Plant Journal</i> , 1996, 10, 515-522.	2.8	294
288	Repression of the Ac-transposase gene promoter by Ac transposase. <i>Plant Journal</i> , 1996, 9, 911-917.	2.8	11

#	ARTICLE	IF	CITATIONS
289	Ensnaring microbes: the components of plant disease resistance. <i>New Phytologist</i> , 1996, 133, 11-34.	3.5	14
290	The Tomato Dwarf Gene Isolated by Heterologous Transposon Tagging Encodes the First Member of a New Cytochrome P450 Family. <i>Plant Cell</i> , 1996, 8, 959.	3.1	35
291	Four <i>Arabidopsis</i> RPP Loci Controlling Resistance to the <i>Noco2</i> Isolate of <i>Peronospora parasitica</i> Map to Regions Known to Contain Other RPP Recognition Specificities. <i>Molecular Plant-Microbe Interactions</i> , 1996, 9, 464.	1.4	44
292	Somatic and germinal activities of maize Activator (Ac) transposase mutants in transgenic tobacco. <i>Plant Journal</i> , 1995, 8, 45-54.	2.8	15
293	Identification of amplified restriction fragment polymorphism (AFLP) markers tightly linked to the tomato Cf-9 gene for resistance to <i>Cladosporium fulvum</i> . <i>Plant Journal</i> , 1995, 8, 785-794.	2.8	215
294	Altered regulation of tomato and tobacco pigmentation genes caused by the delila gene of <i>Antirrhinum</i> . <i>Plant Journal</i> , 1995, 7, 333-339.	2.8	136
295	The maize transposable element Ac is mobile in the legume <i>Lotus japonicus</i> . <i>Plant Molecular Biology</i> , 1995, 27, 981-993.	2.0	67
296	Analysis of splice donor and acceptor site function in a transposable gene trap derived from the maize element Activator. <i>Molecular Genetics and Genomics</i> , 1995, 249, 91-101.	2.4	16
297	Plant disease resistance genes: unravelling how they work. <i>Canadian Journal of Botany</i> , 1995, 73, 495-505.	1.2	10
298	Patterns of gene action in plant development revealed by enhancer trap and gene trap transposable elements.. <i>Genes and Development</i> , 1995, 9, 1797-1810.	2.7	671
299	Aberrant Transpositions of Maize Double Ds-Like Elements Usually Involve Ds Ends on Sister Chromatids. <i>Plant Cell</i> , 1995, 7, 1235.	3.1	18
300	Molecular genetics of plant disease resistance. <i>Science</i> , 1995, 268, 661-667.	6.0	866
301	Functional Expression of a Fungal Avirulence Gene from a Modified Potato Virus X Genome. <i>Molecular Plant-Microbe Interactions</i> , 1995, 8, 181.	1.4	73
302	High-Resolution Mapping of the Physical Location of the Tomato Cf-2 Gene. <i>Molecular Plant-Microbe Interactions</i> , 1995, 8, 200.	1.4	36
303	Identification of Two Genes Required in Tomato for Full Cf-9: Dependent Resistance to <i>Cladosporium fulvum</i> . <i>Plant Cell</i> , 1994, 6, 361.	3.1	48
304	RFLP linkage analysis of the Cf-4 and Cf-9 genes for resistance to <i>Cladosporium fulvum</i> in tomato. <i>Theoretical and Applied Genetics</i> , 1994, 88, 691-700.	1.8	62
305	Chloroplast targeting of spectinomycin adenyltransferase provides a cell-autonomous marker for monitoring transposon excision in tomato and tobacco. <i>Molecular Genetics and Genomics</i> , 1994, 244, 189-196.	2.4	6
306	Analysis of the chromosomal distribution of transposon-carrying T-DNAs in tomato using the inverse polymerase chain reaction. <i>Molecular Genetics and Genomics</i> , 1994, 242, 573-585.	2.4	82

#	ARTICLE	IF	CITATIONS
307	Plant Pathology: Resistance crumbles?. <i>Current Biology</i> , 1994, 4, 67-69.	1.8	11
308	Plant Pathology: Paranoid plants have their genes examined. <i>Current Biology</i> , 1994, 4, 749-751.	1.8	28
309	Isolation of the tomato Cf-9 gene for resistance to <i>Cladosporium fulvum</i> by transposon tagging. <i>Science</i> , 1994, 266, 789-793.	6.0	885
310	Developmentally regulated cell death on expression of the fungal avirulence gene Avr9 in tomato seedlings carrying the disease-resistance gene Cf-9.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1994, 91, 10445-10449.	3.3	121
311	Incomplete Dominance of Tomato Cf Genes for Resistance to <i>Cladosporium fulvum</i> . <i>Molecular Plant-Microbe Interactions</i> , 1994, 7, 58.	1.4	65
312	CfGene-Dependent Induction of a $\beta$ -1,3-Glucanase Promoter in Tomato Plants Infected with <i>Cladosporium fulvum</i> . <i>Molecular Plant-Microbe Interactions</i> , 1994, 7, 645.	1.4	24
313	Phenotypic characterization and molecular mapping of the <i>Arabidopsis thaliana</i> locus RPP5, determining disease resistance to <i>Peronospora parasitica</i> . <i>Plant Journal</i> , 1993, 4, 821-831.	2.8	83
314	Transactivation of Ds elements in plants of lettuce ( <i>Lactuca sativa</i> ). <i>Molecular Genetics and Genomics</i> , 1993, 241-241, 389-398.	2.4	16
315	Effects of gene dosage and sequence modification on the frequency and timing of transposition of the maize element Activator (Ac) in tobacco. <i>Plant Molecular Biology</i> , 1993, 21, 157-170.	2.0	36
316	Alkali treatment for rapid preparation of plant material for reliable PCR analysis. <i>Plant Journal</i> , 1993, 3, 493-494.	2.8	274
317	Use of the maize transposons Activator and Dissociation to show that phosphinothricin and spectinomycin resistance genes act non-cell-autonomously in tobacco and tomato seedlings. <i>Transgenic Research</i> , 1993, 2, 63-78.	1.3	19
318	High level expression of the Activator transposase gene inhibits the excision of Dissociation in tobacco cotyledons. <i>Cell</i> , 1993, 75, 507-517.	13.5	72
319	A Genetic Analysis of DNA Sequence Requirements for Dissociation State I Activity in Tobacco. <i>Plant Cell</i> , 1993, 5, 501.	3.1	11
320	Heterologous Transposon Tagging of the DRL1 Locus in <i>Arabidopsis</i> . <i>Plant Cell</i> , 1993, 5, 631.	3.1	29
321	Studies on the Mechanism by Which Tomato Cf ( <i>Cladosporium fulvum</i> ) Resistance Genes Activate Plant Defence. <i>Current Plant Science and Biotechnology in Agriculture</i> , 1993, , 457-461.	0.0	3
322	Close Linkage Between the Cf-2/Cf-5 and Mi Resistance Loci in Tomato. <i>Molecular Plant-Microbe Interactions</i> , 1993, 6, 341.	1.4	102
323	Two Complex Resistance Loci Revealed in Tomato by Classical and RFLP Mapping of the Cf-2, Cf-4, Cf-5, and Cf-9 Genes for Resistance to <i>Cladosporium fulvum</i> . <i>Molecular Plant-Microbe Interactions</i> , 1993, 6, 348.	1.4	113
324	Research Notes Use of Fungal Transformants Expressing $\beta$ -Glucuronidase Activity to Detect Infection and Measure Hyphal Biomass in Infected Plant Tissues. <i>Molecular Plant-Microbe Interactions</i> , 1993, 6, 521.	1.4	57

#	ARTICLE	IF	CITATIONS
325	Promoter Fusions to the Activator Transposase Gene Cause Distinct Patterns of Dissociation Excision in Tobacco Cotyledons. <i>Plant Cell</i> , 1992, 4, 573.	3.1	0
326	Elevated Levels of Activator Transposase mRNA Are Associated with High Frequencies of Dissociation Excision in Arabidopsis. <i>Plant Cell</i> , 1992, 4, 583.	3.1	0
327	Development of an efficient two-element transposon tagging system in Arabidopsis thaliana. <i>Molecular Genetics and Genomics</i> , 1992, 233, 449-61.	2.4	99
328	Behaviour of the maize transposable element Ac in Arabidopsis thaliana. <i>Plant Journal</i> , 1992, 2, 69-81.	2.8	113
329	Effective vectors for transformation, expression of heterologous genes, and assaying transposon excision in transgenic plants. <i>Transgenic Research</i> , 1992, 1, 285-297.	1.3	301
330	Expression and stability of amplified genes encoding 5-enolpyruvylshikimate-3-phosphate synthase in glyphosate-tolerant tobacco cells. <i>Plant Molecular Biology</i> , 1991, 17, 1127-1138.	2.0	47
331	The Mechanism and Control of Tam3 Transposition. , 1991, , 317-332.		0
332	Aminoglycoside-3'-adenyltransferase confers resistance to spectinomycin and streptomycin in Nicotiana tabacum. <i>Plant Molecular Biology</i> , 1990, 14, 197-205.	2.0	59
333	Preferential Transposition of the Maize Element Activator to Linked Chromosomal Locations in Tobacco. <i>Plant Cell</i> , 1990, 2, 701.	3.1	32
334	Visual Detection of Transposition of the Maize Element Activator (Ac) in Tobacco Seedlings. <i>Science</i> , 1989, 244, 204-207.	6.0	187
335	Relative strengths of the 35S califlower mosaic virus, 1â€², 2â€², and nopaline synthase promoters in transformed tobacco sugarbeet and oilseed rape callus tissue. <i>Molecular Genetics and Genomics</i> , 1988, 212, 182-190.	2.4	113
336	Improved expression of streptomycin resistance in plants due to a deletion in the streptomycin phosphotransferase coding sequence. <i>Molecular Genetics and Genomics</i> , 1988, 214, 456-459.	2.4	39
337	Expression of bacterial chitinase protein in tobacco leaves using two photosynthetic gene promoters. <i>Molecular Genetics and Genomics</i> , 1988, 212, 536-542.	2.4	55
338	Coordinated expression between two photosynthetic petunia genes in transgenic plants. <i>Molecular Genetics and Genomics</i> , 1988, 211, 507-514.	2.4	33
339	Influence of flanking sequences on variability in expression levels of an introduced gene in transgenic tobacco plants. <i>Nucleic Acids Research</i> , 1988, 16, 9267-9283.	6.5	141
340	An efficient mobilizable cosmid vector, pRK7813, and its use in a rapid method for marker exchange in Pseudomonas fluorescens strain HV37a. <i>Gene</i> , 1987, 61, 299-306.	1.0	145
341	Optimizing the expression of chimeric genes in plant cells. <i>Molecular Genetics and Genomics</i> , 1987, 210, 572-577.	2.4	56
342	T-DNA is organized predominantly in inverted repeat structures in plants transformed with Agrobacterium tumefaciens C58 derivatives. <i>Molecular Genetics and Genomics</i> , 1987, 207, 471-477.	2.4	158

#	ARTICLE	IF	CITATIONS
343	T-DNA structure and gene expression in petunia plants transformed by <i>Agrobacterium tumefaciens</i> C58 derivatives. <i>Molecular Genetics and Genomics</i> , 1987, 207, 478-485.	2.4	135
344	A dominant nuclear streptomycin resistance marker for plant cell transformation. <i>Molecular Genetics and Genomics</i> , 1987, 210, 86-91.	2.4	42
345	The Expression of Introduced Genes in Regenerated Plants. , 1987, , 45-59.		5
346	Isolation and characterization of genes encoding two chitinase enzymes from <i>Serratia marcescens</i> . <i>EMBO Journal</i> , 1986, 5, 467-473.	3.5	228
347	High level expression of introduced chimaeric genes in regenerated transformed plants. <i>EMBO Journal</i> , 1985, 4, 2411-2418.	3.5	438
348	<i>Klebsiella pneumoniae</i> nifA product activates the <i>Rhizobium meliloti</i> nitrogenase promoter. <i>Nature</i> , 1983, 301, 728-732.	13.7	130
349	The mapping of highly-repeated DNA families and their relationship to C-bands in chromosomes of <i>Secale cereale</i> . <i>Chromosoma</i> , 1982, 86, 595-612.	1.0	120
350	The structure, amount and chromosomal localisation of defined repeated DNA sequences in species of the genus <i>Secale</i> . <i>Chromosoma</i> , 1982, 86, 613-641.	1.0	106
351	A molecular description of telomeric heterochromatin in <i>secale</i> species. <i>Cell</i> , 1980, 19, 545-560.	13.5	610
352	SMRT RenSeq protocol. <i>Protocol Exchange</i> , 0, , .	0.3	5