List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/478231/publications.pdf Version: 2024-02-01

		61984	43889
101	13,133	43	91
papers	citations	h-index	g-index
111	111	111	14666
all docs	docs citations	times ranked	citing authors

ENANAN HALL

#	Article	IF	CITATIONS
1	DNA-Repair Defects and Olaparib in Metastatic Prostate Cancer. New England Journal of Medicine, 2015, 373, 1697-1708.	27.0	1,796
2	A Randomized Trial of Exemestane after Two to Three Years of Tamoxifen Therapy in Postmenopausal Women with Primary Breast Cancer. New England Journal of Medicine, 2004, 350, 1081-1092.	27.0	1,694
3	Parotid-sparing intensity modulated versus conventional radiotherapy in head and neck cancer (PARSPORT): a phase 3 multicentre randomised controlled trial. Lancet Oncology, The, 2011, 12, 127-136.	10.7	1,407
4	Conventional versus hypofractionated high-dose intensity-modulated radiotherapy for prostate cancer: 5-year outcomes of the randomised, non-inferiority, phase 3 CHHiP trial. Lancet Oncology, The, 2016, 17, 1047-1060.	10.7	941
5	Survival and safety of exemestane versus tamoxifen after 2–3 years' tamoxifen treatment (Intergroup) Tj ETQq1	1.0,7843	14 rgBT /0 805
6	Radiotherapy with or without Chemotherapy in Muscle-Invasive Bladder Cancer. New England Journal of Medicine, 2012, 366, 1477-1488.	27.0	788
7	Mortality in relation to consumption of alcohol: 13 years' observations on male British doctors. BMJ: British Medical Journal, 1994, 309, 911-918.	2.3	637
8	Intermittent Androgen Suppression for Rising PSA Level after Radiotherapy. New England Journal of Medicine, 2012, 367, 895-903.	27.0	428
9	Skeletal effects of exemestane on bone-mineral density, bone biomarkers, and fracture incidence in postmenopausal women with early breast cancer participating in the Intergroup Exemestane Study (IES): a randomised controlled study. Lancet Oncology, The, 2007, 8, 119-127.	10.7	404
10	Conventional versus hypofractionated high-dose intensity-modulated radiotherapy for prostate cancer: preliminary safety results from the CHHiP randomised controlled trial. Lancet Oncology, The, 2012, 13, 43-54.	10.7	303
11	Clinical development of new drug–radiotherapy combinations. Nature Reviews Clinical Oncology, 2016, 13, 627-642.	27.6	230
12	Independent validation of genes and polymorphisms reported to be associated with radiation toxicity: a prospective analysis study. Lancet Oncology, The, 2012, 13, 65-77.	10.7	202
13	Quality of Life in the Intergroup Exemestane Study: A Randomized Trial of Exemestane Versus Continued Tamoxifen After 2 to 3 Years of Tamoxifen in Postmenopausal Women With Primary Breast Cancer. Journal of Clinical Oncology, 2006, 24, 910-917.	1.6	180
14	Sequential docetaxel as adjuvant chemotherapy for early breast cancer (TACT): an open-label, phase III, randomised controlled trial. Lancet, The, 2009, 373, 1681-1692.	13.7	168
15	A genome wide association study (GWAS) providing evidence of an association between common genetic variants and late radiotherapy toxicity. Radiotherapy and Oncology, 2014, 111, 178-185.	0.6	128
16	Hypofractionated radiotherapy versus conventionally fractionated radiotherapy for patients with intermediate-risk localised prostate cancer: 2-year patient-reported outcomes of the randomised, non-inferiority, phase 3 CHHiP trial. Lancet Oncology, The, 2015, 16, 1605-1616.	10.7	126
17	Randomized Noninferiority Trial of Reduced High-Dose Volume Versus Standard Volume Radiation Therapy for Muscle-Invasive Bladder Cancer: Results of the BC2001 Trial (CRUK/01/004). International Journal of Radiation Oncology Biology Physics, 2013, 87, 261-269.	0.8	115
18	A three-stage genome-wide association study identifies a susceptibility locus for late radiotherapy toxicity at 2q24.1. Nature Genetics, 2014, 46, 891-894.	21.4	114

#	Article	IF	CITATIONS
19	Key issues in recruitment to randomised controlled trials with very different interventions: a qualitative investigation of recruitment to the SPARE trial (CRUK/07/011). Trials, 2011, 12, 78.	1.6	112
20	LIFE AND DEATH OF SPARE (SELECTIVE BLADDER PRESERVATION AGAINST RADICAL EXCISION): REFLECTIONS ON WHY THE SPARE TRIAL CLOSED. BJU International, 2010, 106, 753-755.	2.5	108
21	Clinical trial designs for rare diseases: Studies developed and discussed by the International Rare Cancers Initiative. European Journal of Cancer, 2015, 51, 271-281.	2.8	108
22	Clinical and patientâ€reported outcomes of <scp>SPARE</scp> – a randomised feasibility study of selective bladder preservation versus radical cystectomy. BJU International, 2017, 120, 639-650.	2.5	105
23	Double-blind placebo-controlled randomised trial of vitamin E and pentoxifylline in patients with chronic arm lymphoedema and fibrosis after surgery and radiotherapy for breast cancer. Radiotherapy and Oncology, 2004, 73, 133-139.	0.6	99
24	Individual patient data meta-analysis shows a significant association between the ATM rs1801516 SNP and toxicity after radiotherapy in 5456 breast and prostate cancer patients. Radiotherapy and Oncology, 2016, 121, 431-439.	0.6	98
25	Randomized controlled trial of the effect of selenium supplementation on thyroid function in the elderly in the United Kingdom. American Journal of Clinical Nutrition, 2008, 87, 370-378.	4.7	97
26	Impact of Selenium on Mood and Quality of Life: A Randomized, Controlled Trial. Biological Psychiatry, 2006, 59, 147-154.	1.3	91
27	R-IDEAL: A Framework for Systematic Clinical Evaluation of Technical Innovations in Radiation Oncology. Frontiers in Oncology, 2017, 7, 59.	2.8	90
28	Second cancer risk and mortality in men treated with radiotherapy for stage I seminoma. British Journal of Cancer, 2014, 110, 256-263.	6.4	85
29	Consensus Guidelines and Contouring Atlas for Pelvic Node Delineation in Prostate and Pelvic Node Intensity Modulated Radiation Therapy. International Journal of Radiation Oncology Biology Physics, 2015, 92, 874-883.	0.8	85
30	DARS: a phase III randomised multicentre study of dysphagia- optimised intensity- modulated radiotherapy (Do-IMRT) versus standard intensity- modulated radiotherapy (S-IMRT) in head and neck cancer. BMC Cancer, 2016, 16, 770.	2.6	83
31	Phase I Trial of the PARP Inhibitor Olaparib and AKT Inhibitor Capivasertib in Patients with <i>BRCA1/2</i> - and Non– <i>BRCA1/2</i> -Mutant Cancers. Cancer Discovery, 2020, 10, 1528-1543.	9.4	82
32	Non-randomised phase II trial of hyperbaric oxygen therapy in patients with chronic arm lymphoedema and tissue fibrosis after radiotherapy for early breast cancer. Radiotherapy and Oncology, 2004, 70, 217-224.	0.6	81
33	Diffusion-weighted Imaging as a Treatment Response Biomarker for Evaluating Bone Metastases in Prostate Cancer: A Pilot Study. Radiology, 2017, 283, 168-177.	7.3	81
34	Biomarkers Associating with PARP Inhibitor Benefit in Prostate Cancer in the TOPARP-B Trial. Cancer Discovery, 2021, 11, 2812-2827.	9.4	78
35	Dosimetric explanations of fatigue in head and neck radiotherapy: An analysis from the PARSPORT Phase III trial. Radiotherapy and Oncology, 2012, 104, 205-212.	0.6	73
36	Meta-analysis of Genome Wide Association Studies Identifies Genetic Markers of Late Toxicity Following Radiotherapy for Prostate Cancer. EBioMedicine, 2016, 10, 150-163.	6.1	69

#	Article	IF	CITATIONS
37	Human Epidermal Growth Factor Receptor 2 Status Correlates With Lymph Node Involvement in Patients With Estrogen Receptor (ER) –Negative, but With Grade in Those With ER-Positive Early-Stage Breast Cancer Suitable for Cytotoxic Chemotherapy. Journal of Clinical Oncology, 2007, 25, 4423-4430.	1.6	66
38	Validation and Reliability Testing of the EORTC QLQ-NMIBC24 Questionnaire Module to Assess Patient-reported Outcomes in Non–Muscle-invasive Bladder Cancer. European Urology, 2014, 66, 1148-1156.	1.9	66
39	Novel approaches to improve the therapeutic index of head and neck radiotherapy: An analysis of data from the PARSPORT randomised phase III trial. Radiotherapy and Oncology, 2012, 103, 82-87.	0.6	65
40	Persistence of bone marrow micrometastases in patients receiving adjuvant therapy for breast cancer: Results at 4 years. International Journal of Cancer, 2005, 114, 94-100.	5.1	63
41	Patient-reported Quality of Life Outcomes in Patients Treated for Muscle-invasive Bladder Cancer with Radiotherapy ± Chemotherapy in the BC2001 Phase III Randomised Controlled Trial. European Urology, 2020, 77, 260-268.	1.9	58
42	ls High Dose Therapy Superior to Conventional Dose Therapy as Initial Treatment for Relapsed Germ Cell Tumors? The TIGER Trial. Journal of Cancer, 2011, 2, 374-377.	2.5	55
43	Phase II study of vitamin E and pentoxifylline in patients with late side effects of pelvic radiotherapy. Radiotherapy and Oncology, 2005, 75, 334-341.	0.6	51
44	Double-blind, placebo-controlled, randomised phase II trial of IH636 grape seed proanthocyanidin extract (GSPE) in patients with radiation-induced breast induration. Radiotherapy and Oncology, 2006, 79, 45-51.	0.6	43
45	Pre-trial quality assurance processes for an intensity-modulated radiation therapy (IMRT) trial: PARSPORT, a UK multicentre Phase III trial comparing conventional radiotherapy and parotid-sparing IMRT for locally advanced head and neck cancer. British Journal of Radiology, 2009, 82, 585-594.	2.2	43
46	The 111 Study: A Single-arm, Phase 3 Trial Evaluating One Cycle of Bleomycin, Etoposide, and Cisplatin as Adjuvant Chemotherapy in High-risk, Stage 1 Nonseminomatous or Combined Germ Cell Tumours of the Testis. European Urology, 2020, 77, 344-351.	1.9	41
47	Time to define an international standard of postoperative care for resected upper urinary tract transitional cell carcinoma (TCC) – opening of the periâ€operative chemotherapy versus surveillance in upper tract urothelial cancer (POUT) Trial. BJU International, 2012, 110, 919-921.	2.5	37
48	Comparison of patient reported quality of life and impact of treatment side effects experienced with a taxane-containing regimen and standard anthracycline based chemotherapy for early breast cancer: 6year results from the UK TACT trial (CRUK/01/001). European Journal of Cancer, 2014, 50, 2375-2389.	2.8	36
49	Triplet Therapy with Palbociclib, Taselisib, and Fulvestrant in <i>PIK3CA</i> -Mutant Breast Cancer and Doublet Palbociclib and Taselisib in Pathway-Mutant Solid Cancers. Cancer Discovery, 2021, 11, 92-107.	9.4	36
50	Long-term endometrial effects in postmenopausal women with early breast cancer participating in the Intergroup Exemestane Study (IES)—a randomised controlled trial of exemestane versus continued tamoxifen after 2–3 years tamoxifen. Annals of Oncology, 2010, 21, 498-505.	1.2	34
51	The Patient Deficit Model Overturned: a qualitative study of patients' perceptions of invitation to participate in a randomized controlled trial comparing selective bladder preservation against surgery in muscle invasive bladder cancer (SPARE, CRUK/07/011). Trials, 2012, 13, 228.	1.6	28
52	Two-Year and Lifetime Cost-Effectiveness of Intensity Modulated Radiation Therapy Versus 3-Dimensional Conformal Radiation Therapy for Head-and-Neck Cancer. International Journal of Radiation Oncology Biology Physics, 2013, 87, 683-689.	0.8	27
53	CALIBER: a phase II randomized feasibility trial of chemoablation with mitomycinâ€C vs surgical management in lowâ€risk nonâ€muscleâ€invasive bladder cancer. BJU International, 2020, 125, 817-826. 	2.5	27
54	Diffusion-weighted MRI in Advanced Epithelial Ovarian Cancer: Apparent Diffusion Coefficient as a Response Marker. Radiology, 2019, 293, 374-383.	7.3	25

#	Article	IF	CITATIONS
55	Alcohol and coronary heart disease reduction among British doctors: confounding or causality?. European Heart Journal, 1997, 18, 23-25.	2.2	24
56	Aromatase inhibitors as adjuvant therapies in patients with breast cancer. Journal of Steroid Biochemistry and Molecular Biology, 2003, 86, 309-311.	2.5	21
57	Outcomes in Patients with Muscle-invasive Bladder Cancer Treated with Neoadjuvant Chemotherapy Followed by (Chemo)radiotherapy in the BC2001 Trial. European Urology, 2021, 79, 307-315.	1.9	20
58	Randomised pilot study of dose escalation using conformal radiotherapy in prostate cancer: long-term follow-up. British Journal of Cancer, 2013, 109, 651-657.	6.4	19
59	Defining Bowel Dose Volume Constraints for Bladder Radiotherapy Treatment Planning. Clinical Oncology, 2015, 27, 22-29.	1.4	18
60	Common genetic variation associated with increased susceptibility to prostate cancer does not increase risk of radiotherapy toxicity. British Journal of Cancer, 2016, 114, 1165-1174.	6.4	17
61	Quality assuring "Plan of the day―selection in a multicentre adaptive bladder trial: Implementation of a pre-accrual IGRT guidance and assessment module. Clinical and Translational Radiation Oncology, 2019, 19, 27-32.	1.7	14
62	Abstract CT323: Accelerated phase I trial of two schedules of the combination of the PARP inhibitor olaparib and AKT inhibitor AZD5363 using a novel intrapatient dose escalation design in advanced cancer patients. Cancer Research, 2015, 75, CT323-CT323.	0.9	12
63	Results of a 2 x 2 Phase III Randomized Trial of Synchronous Chemo-radiotherapy (CRT) Compared to Radiotherapy (RT) Alone and Standard vs. Reduced High Volume RT in Muscle Invasive Bladder Cancer (MIBC) (BC2001 CRUK/01/004). International Journal of Radiation Oncology Biology Physics, 2010, 78, \$2.53	0.8	10
64	The predictive and prognostic value of tumour necrosis in muscle invasive bladder cancer patients receiving radiotherapy with or without chemotherapy in the BC2001 trial (CRUK/01/004). British Journal of Cancer, 2017, 116, 649-657.	6.4	9
65	Benefits and challenges of electronic data capture (EDC) systems versus paper case report forms. Trials, 2015, 16, .	1.6	8
66	Boxing Bladder Cancer with COX-2–specific Inhibition. Cancer Prevention Research, 2011, 4, 1534-1535.	1.5	6
67	Current attitudes of head and neck oncologists in the United Kingdom to induction chemotherapy for locally advanced head and neck cancer: A survey of centres participating in a national randomised controlled trial. Oral Oncology, 2014, 50, 141-146.	1.5	6
68	Prostate cancer and hypofractionation: reflections on recent randomised phase III clinical trial results. Translational Andrology and Urology, 2017, 6, 134-136.	1.4	5
69	HYBRID — Evaluating New Radiation Technology in Patients with Unmet Needs. Clinical Oncology, 2013, 25, 546-548.	1.4	4
70	Implementing electronic data capture (EDC) training for site staff. Trials, 2015, 16, .	1.6	4
71	EP-1566 MR-guided online adaptive radiotherapy: First experience in the UK. Radiotherapy and Oncology, 2019, 133, S845.	0.6	4
72	Letter in response to the Wedlake et al. paper â€~Evaluating the efficacy of statins and ACE-inhibitors in reducing gastrointestinal toxicity in patients receiving radiotherapy for pelvic malignancies'. European Journal of Cancer, 2013, 49, 1783-1786.	2.8	3

#	Article	IF	CITATIONS
73	VinCaP: a phase II trial of vinflunine in locally advanced and metastatic squamous carcinoma of the penis. British Journal of Cancer, 2021, , .	6.4	3
74	Phase III trial of conformal radiotherapy following neoadjuvant hormone treatment in early prostate cancer. International Journal of Radiation Oncology Biology Physics, 2002, 54, 134-135.	0.8	2
75	First Toxicity Report of the BC2001 Trial: A Multicentre Phase III Randomised Trial of Radiotherapy with and without Synchronous Chemotherapy in Muscle Invasive Bladder Cancer ISCRTN No. 68324339, EUDRACT No. 2004-000164-26. International Journal of Radiation Oncology Biology Physics, 2008, 72, S7.	0.8	2
76	Results of a Phase III Multi-centre Randomised Controlled Trial of Intensity Modulated (IMRT) vs Conventional Radiotherapy (RT) in HeadÂand Neck Cancer (PARSPORT: ISRCTN48243537; CRUK/03/005). Clinical Oncology, 2010, 22, 899.	1.4	2
77	343 Translational phase I trial combining the AKT inhibitor AZD5363 (AZD) and PARP inhibitor Olaparib (Ola) in advanced cancer patients (pts). European Journal of Cancer, 2015, 51, S68.	2.8	2
78	Design considerations when transitioning from paper case report forms (CRFS) to electronic data capture (EDC). Trials, 2015, 16, .	1.6	2
79	Abstract CT322: DNA repair defects and antitumor activity with PARP inhibition: TOPARP, a phase II trial of olaparib in metastatic castration resistant prostate cancer. , 2015, , .		2
80	A Road Map for Designing Phase I Clinical Trials of Radiotherapy–Novel Agent Combinations. Clinical Cancer Research, 2022, 28, 3639-3651.	7.0	2
81	430 Cross-sectional study of Quality of Life (QL) 6 years after start of treatment in the UK Taxotere as Adjuvant Chemotherapy Trial (TACT; CRUK01/001). European Journal of Cancer, Supplement, 2010, 8, 184.	2.2	1
82	497 oral PHASE III PILOT STUDY OF DOSE ESCALATION USING CON-FORMAL RADIOTHERAPY IN PROSTATE CANCER: LONG TERM FOLLOW UP. Radiotherapy and Oncology, 2011, 99, S201.	0.6	1
83	Phase II investigation of a PARP inhibitor (olaparib) in castration resistant prostate cancer (CRPC) which incorporates the possibility that treatment effect may be restricted to biomarker defined subgroups. Trials, 2011, 12, .	1.6	1
84	Using a Bayesian approach with reverse philosophy to design clinical trials in rare diseases. Trials, 2013, 14, .	1.6	1
85	2507 Prognostic factors in muscle invasive bladder cancer patients receiving radiotherapy with or without chemotherapy in the BC2001 trial (CRUK/01/004). European Journal of Cancer, 2015, 51, S475-S476.	2.8	1
86	Remote training solutions for complex intervention trials. Trials, 2015, 16, .	1.6	1
87	Methodology for tissue sample collection within a translational sub-study of the CHHiP trial (CRUK/06/016), a large randomised phase III trial in localised prostate cancer. Clinical and Translational Radiation Oncology, 2018, 10, 1-6.	1.7	1
88	Abstract CT138: Translating preclinical observations to the clinic: Combination of the dual m-TORC1/2 inhibitor AZD2014 and paclitaxel in ovarian and lung cancer. , 2015, , .		1
89	A phase I trial of selective PI3K inhibitor taselisib (tas) plus palbociclib (palb) with and without endocrine therapy incorporating pharmacodynamic (PD) studies in patients (pts) with advanced cancers Journal of Clinical Oncology, 2017, 35, 2573-2573.	1.6	1
90	Quantitative Estimates of the Effects of Concomitant Chemotherapy on Acute Dysphagia in Patients Receiving Radical Treatment for Head and Neck Cancer. International Journal of Radiation Oncology Biology Physics, 2009, 75, S612.	0.8	0

#	Article	IF	CITATIONS
91	Re: Marco Roscigno, Maurizio Brausi, Axel Heidenreich, et al. Lymphadenectomy at the Time of Nephroureterectomy for Upper Tract Urothelial Cancer. Eur Urol 2011;60:776–83. European Urology, 2012, 61, e16.	1.9	0
92	OC-0432: Evaluating the feasibility of sparing CNS structures to reduce acute fatigue from IMRT for head and neck cancer. Radiotherapy and Oncology, 2013, 106, S166.	0.6	0
93	OC-0154: IGRT for prostate cancer $\tilde{A}\pm$ results from the CHHiP IGRT phase II sub-study. Radiotherapy and Oncology, 2014, 111, S59.	0.6	0
94	2642 The predictive and prognostic value of tumour necrosis in muscle invasive bladder cancer patients receiving radiotherapy with or without chemotherapy in the BC2001 (CRUK/01/004) trial. European Journal of Cancer, 2015, 51, S527-S528.	2.8	0
95	Dynamic prediction methods in the BC2001 clinical trial. Trials, 2015, 16, .	1.6	0
96	2579 Evaluation of clinical decision-making and the use of circulating tumor cells (CTCs) by physicians treating castration-resistant prostate cancer (CRPC). European Journal of Cancer, 2015, 51, S504.	2.8	0
97	OC-0340: Effect of dose and image guided radiotherapy (IGRT) on erectile potency (EP) in prostate radiotherapy. Radiotherapy and Oncology, 2016, 119, S156-S157.	0.6	0
98	Clinical management and research priorities for high-risk prostate cancer in the UK: Meeting report of a multidisciplinary panel in conjunction with the NCRI Prostate Cancer Clinical Studies Localised Subgroup. Journal of Clinical Urology, 2016, 9, 369-379.	0.1	0
99	OC-0634 Implementation of plan of the day adaptive radiotherapy: Compliance to guidelines. Radiotherapy and Oncology, 2019, 133, S337-S338.	0.6	0
100	Reply to Wei Liu, Xiaoping Liu, Sheng Li's Letter to the Editor, re: Robert A. Huddart, Emma Hall, Rebecca Lewis, et al. Patient-reported Quality of Life Outcomes in Patients Treated for Muscle-invasive Bladder Cancer with Radiotherapy ± Chemotherapy in the BC2001 Phase III Randomised Controlled Trial. Eur Urol 2020;77:260–8. European Urology, 2020, 77, e156-e157.	1.9	0
101	TAX-TORC: A phase I trial of vistusertib (AZD2014) in combination with weekly paclitaxel with integrated pharmacodynamic (PD) and molecular characterization (MC) studies Journal of Clinical	1.6	0