## Luis Miguel Ortega Mora

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/4782910/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Epidemiology and Control of Neosporosis and Neospora caninum. Clinical Microbiology Reviews, 2007, 20, 323-367.	5.7	825
2	A review of the importance of cryptosporidiosis in farm animals. International Journal for Parasitology, 1999, 29, 1269-1287.	1.3	355
3	Toxoplasma gondii infection and toxoplasmosis in farm animals: Risk factors and economic impact. Food and Waterborne Parasitology, 2019, 15, e00037.	1.1	206
4	Quantitative Detection of Neospora caninum in Bovine Aborted Fetuses and Experimentally Infected Mice by Real-Time PCR. Journal of Clinical Microbiology, 2002, 40, 1194-1198.	1.8	134
5	Detection of Infectious Cryptosporidium parvum Oocysts in Mussels ( Mytilus galloprovincialis ) and Cockles ( Cerastoderma edule ). Applied and Environmental Microbiology, 2000, 66, 1866-1870.	1.4	121
6	Occurrence of Neospora caninum and Toxoplasma gondii infections in ovine and caprine abortions. Veterinary Parasitology, 2012, 187, 312-318.	0.7	115
7	A century of bovine besnoitiosis: an unknown disease re-emerging in Europe. Trends in Parasitology, 2013, 29, 407-415.	1.5	114
8	Supranational comparison of Neospora caninum seroprevalences in cattle in Germany, The Netherlands, Spain and Sweden. Veterinary Parasitology, 2006, 137, 17-27.	0.7	96
9	In Vitro Metacestodicidal Activities of Genistein and Other Isoflavones against Echinococcus multilocularis and Echinococcus granulosus. Antimicrobial Agents and Chemotherapy, 2006, 50, 3770-3778.	1.4	87
10	In Vitro and In Vivo Treatments of <i>Echinococcus</i> Protoscoleces and Metacestodes with Artemisinin and Artemisinin Derivatives. Antimicrobial Agents and Chemotherapy, 2008, 52, 3447-3450.	1.4	86
11	Molecular approaches to diversity of populations of apicomplexan parasites. International Journal for Parasitology, 2009, 39, 175-189.	1.3	85
12	Influence of age and purpose for testing on the cut-off selection of serological methods in bovine neosporosis. Veterinary Research, 2003, 34, 341-352.	1.1	81
13	Comparison and standardisation of serological methods for the diagnosis of Neospora caninum infection in bovines. Veterinary Parasitology, 2004, 120, 11-22.	0.7	76
14	Isolation and genetic characterization of <i>Neospora caninum</i> from asymptomatic calves in Spain. Parasitology, 2008, 135, 1651-1659.	0.7	76
15	Pattern of recognition of Neospora caninum tachyzoite antigens by naturally infected pregnant cattle and aborted foetuses. Veterinary Parasitology, 2002, 107, 15-27.	0.7	75
16	Dynamics of <i>Besnoitia besnoiti</i> infection in cattle. Parasitology, 2014, 141, 1419-1435.	0.7	75
17	Neospora caninum infection during early pregnancy in cattle: how the isolate influences infection dynamics, clinical outcome and peripheral and local immune responses. Veterinary Research, 2014, 45, 10.	1.1	75
18	<i>In Vitro</i> and <i>In Vivo</i> Effects of the Bumped Kinase Inhibitor 1294 in the Related Cyst-Forming Apicomplexans Toxoplasma gondii and Neospora caninum. Antimicrobial Agents and Chemotherapy, 2015, 59, 6361-6374.	1.4	72

#	Article	IF	CITATIONS
19	Chronic bovine besnoitiosis: Intra-organ parasite distribution, parasite loads and parasite-associated lesions in subclinical cases. Veterinary Parasitology, 2013, 197, 95-103.	0.7	71
20	First Isolation of Besnoitia besnoiti from a Chronically Infected Cow in Spain. Journal of Parasitology, 2009, 95, 474-476.	0.3	69
21	Experimental infection with a low virulence isolate of <i>Neospora caninum</i> at 70 days gestation in cattle did not result in foetopathy. Veterinary Research, 2009, 40, 49.	1.1	68
22	Isolation and characterization of a bovine isolate of Neospora caninum with low virulence. Veterinary Parasitology, 2009, 159, 7-16.	0.7	66
23	Seroprevalence of Neospora caninum infection in dairy and beef cattle in Spain. International Journal for Parasitology, 1999, 29, 1201-1208.	1.3	65
24	In vitro invasion efficiency and intracellular proliferation rate comprise virulence-related phenotypic traits of Neospora caninum. Veterinary Research, 2011, 42, 41.	1.1	65
25	Evaluation of ovine abortion associated with Toxoplasma gondii in Spain by different diagnostic techniques. Veterinary Parasitology, 2004, 121, 33-43.	0.7	63
26	Influence of <i>Neospora caninum</i> intra-specific variability in the outcome of infection in a pregnant BALB/c mouse model. Veterinary Research, 2010, 41, 52.	1.1	63
27	ADAPTATION OF NEOSPORA CANINUM ISOLATES TO CELL-CULTURE CHANGES: AN ARGUMENT IN FAVOR OF ITS CLONAL POPULATION STRUCTURE. Journal of Parasitology, 2005, 91, 507-510.	0.3	62
28	MULTILOCUS MICROSATELLITE ANALYSIS REVEALS EXTENSIVE GENETIC DIVERSITY IN NEOSPORA CANINUM. Journal of Parasitology, 2006, 92, 517-524.	0.3	60
29	Temporal Distribution and Parasite Load Kinetics in Blood and Tissues during Neospora caninum Infection in Mice. Infection and Immunity, 2006, 74, 2491-2494.	1.0	60
30	Development and use of an indirect ELISA in an outbreak of bovine besnoitiosis in Spain. Veterinary Record, 2010, 166, 818-822.	0.2	60
31	Infected Dendritic Cells Facilitate Systemic Dissemination and Transplacental Passage of the Obligate Intracellular Parasite Neospora caninum in Mice. PLoS ONE, 2012, 7, e32123.	1.1	60
32	An Inter-Laboratory Comparative Study of Serological Tools Employed in the Diagnosis of <i>Besnoitia besnoiti</i> Infection in Bovines. Transboundary and Emerging Diseases, 2013, 60, 59-68.	1.3	60
33	Neospora caninum infection as a cause of reproductive failure in a sheep flock. Veterinary Research, 2014, 45, 88.	1.1	57
34	In vitro and in vivo effects of 2-methoxyestradiol, either alone or combined with albendazole, against Echinococcus metacestodes. Experimental Parasitology, 2008, 119, 475-482.	0.5	56
35	Molecular characterisation of Cryptosporidium isolates from pet reptiles. Veterinary Parasitology, 2009, 160, 204-210.	0.7	56
36	Detection of Neospora caninum in semen of bulls. Veterinary Parasitology, 2003, 117, 301-308.	0.7	55

#	Article	IF	CITATIONS
37	Evaluation by different diagnostic techniques of bovine abortion associated with Neospora caninum in Spain. Veterinary Parasitology, 2003, 111, 143-152.	0.7	54
38	Development of a murine vertical transmission model for Toxoplasma gondii oocyst infection and studies on the efficacy of bumped kinase inhibitor (BKI)-1294 and the naphthoquinone buparvaquone against congenital toxoplasmosis. Journal of Antimicrobial Chemotherapy, 2017, 72, 2334-2341.	1.3	52
39	Transcriptome modulation of bovine trophoblast cells in vitro by Neospora caninum. International Journal for Parasitology, 2017, 47, 791-799.	1.3	52
40	An interlaboratory comparison of immunohistochemistry and PCR methods for detection of Neospora caninum in bovine foetal tissues. Veterinary Parasitology, 2004, 126, 351-364.	0.7	51
41	The Neospora caninum-Spain 7 isolate induces placental damage, fetal death and abortion in cattle when inoculated in early gestation. Veterinary Parasitology, 2012, 189, 171-181.	0.7	50
42	Placental thrombosis in acute phase abortions during experimental Toxoplasma gondii infection in sheep. Veterinary Research, 2014, 45, 9.	1.1	50
43	Identification and molecular cloning of the Neospora caninum SAG4 gene specifically expressed at bradyzoite stageâ <sup>-+</sup> . Molecular and Biochemical Parasitology, 2006, 146, 89-97.	0.5	49
44	Serological diagnosis of bovine neosporosis: A comparative study of commercially available ELISA tests. Veterinary Parasitology, 2013, 198, 85-95.	0.7	49
45	Genetic Diversity and Geographic Population Structure of Bovine Neospora caninum Determined by Microsatellite Genotyping Analysis. PLoS ONE, 2013, 8, e72678.	1.1	49
46	Usefulness of rNcGRA7- and rNcSAG4-based ELISA tests for distinguishing primo-infection, recrudescence, and chronic bovine neosporosis. Veterinary Parasitology, 2008, 157, 182-195.	0.7	48
47	Vaccines for bovine neosporosis: current status and key aspects for development. Parasite Immunology, 2016, 38, 709-723.	0.7	48
48	Anthelmintic and nutritional effects of heather supplementation on Cashmere goats grazing perennial ryegrass-white clover pastures1. Journal of Animal Science, 2007, 85, 861-870.	0.2	47
49	Influence of the gestational stage on the clinical course, lesional development and parasite distribution in experimental ovine neosporosis. Veterinary Research, 2015, 46, 19.	1.1	45
50	Endogenous transplacental transmission of Neospora caninum during successive pregnancies across three generations of naturally infected sheep. Veterinary Research, 2018, 49, 106.	1.1	45
51	Seroprevalence ofFasciola hepatica infection in sheep in northwestern Spain. Zeitschrift Für Parasitenkunde (Berlin, Germany), 1995, 81, 137-142.	0.8	43
52	Control options for <i>Neospora caninum</i> – is there anything new or are we going backwards?. Parasitology, 2014, 141, 1455-1470.	0.7	43
53	Age-related resistance in ovine cryptosporidiosis: patterns of infection and humoral immune response. Infection and Immunity, 1994, 62, 5003-5009.	1.0	43
54	Detection of Neospora caninum in the semen and blood of naturally infected bulls. Theriogenology, 2005, 63, 1504-1518.	0.9	42

#	Article	IF	CITATIONS
55	The NcGRA7gene encodes the immunodominant 17 kDa antigen ofNeospora caninum. Parasitology, 2007, 134, 41-50.	0.7	42
56	Role of adult sheep in transmission of infection by Cryptosporidium parvum to lambs: confirmation of periparturient rise. International Journal for Parasitology, 1999, 29, 1261-1268.	1.3	41
57	Herd-level risk factors associated with Leptospira spp. seroprevalence in dairy and beef cattle in Spain. Preventive Veterinary Medicine, 2001, 52, 109-117.	0.7	41
58	Experimental ovine toxoplasmosis: influence of the gestational stage on the clinical course, lesion development and parasite distribution. Veterinary Research, 2016, 47, 43.	1.1	40
59	In vitro efficacy of bumped kinase inhibitors against Besnoitia besnoiti tachyzoites. International Journal for Parasitology, 2017, 47, 811-821.	1.3	40
60	Treatment of Toxoplasmosis and Neosporosis in Farm Ruminants: State of Knowledge and Future Trends. Current Topics in Medicinal Chemistry, 2018, 18, 1304-1323.	1.0	40
61	Pattern of recognition of Besnoitia besnoiti tachyzoite and bradyzoite antigens by naturally infected cattle. Veterinary Parasitology, 2009, 164, 104-110.	0.7	39
62	Influence of the stage of pregnancy on Neospora caninum distribution, parasite loads and lesions in aborted bovine foetuses. Theriogenology, 2006, 65, 629-641.	0.9	38
63	Grazing behaviour and performance of lactating suckler cows, ewes and goats on partially improved heathlands. Animal, 2008, 2, 1818-1831.	1.3	38
64	Besnoitia besnoiti lytic cycle in vitro and differences in invasion and intracellular proliferation among isolates. Parasites and Vectors, 2016, 9, 115.	1.0	37
65	Impact of human-associated Escherichia coli clonal groups in Antarctic pinnipeds: presence of ST73, ST95, ST141 and ST131. Scientific Reports, 2018, 8, 4678.	1.6	37
66	Bumped Kinase Inhibitors as therapy for apicomplexan parasitic diseases: lessons learned. International Journal for Parasitology, 2020, 50, 413-422.	1.3	37
67	Diagnosis of bovine neosporosis: Recent advances and perspectives. Acta Parasitologica, 2006, 51, 1-14.	0.4	36
68	Dose-dependent effects of experimental infection with the virulent Neospora caninum Nc-Spain7 isolate in a pregnant mouse model. Veterinary Parasitology, 2015, 211, 133-140.	0.7	36
69	Buparvaquone is active against Neospora caninum in vitro and in experimentally infected mice. International Journal for Parasitology: Drugs and Drug Resistance, 2015, 5, 16-25.	1.4	36
70	COMPARATIVE ANALYSIS OF STRESS AGENTS IN A SIMPLIFIED IN VITRO SYSTEM OF NEOSPORA CANINUM BRADYZOITE PRODUCTION. Journal of Parasitology, 2004, 90, 466-470.	0.3	35
71	COMPARATIVE EFFECT OF NEOSPORA CANINUM INFECTION IN BALB/c MICE AT THREE DIFFERENT GESTATION PERIODS. Journal of Parasitology, 2006, 92, 1286-1291.	0.3	35
72	Failure of a vaccine using immunogenic recombinant proteins rNcSAG4 and rNcGRA7 against neosporosis in mice. Vaccine, 2009, 27, 7331-7338.	1.7	35

#	Article	IF	CITATIONS
73	Virulence in Mice of a Toxoplasma gondii Type II Isolate Does Not Correlate With the Outcome of Experimental Infection in Pregnant Sheep. Frontiers in Cellular and Infection Microbiology, 2018, 8, 436.	1.8	35
74	Seroprevalence and risk factors associated with Neospora caninum infection in different dog populations in Spain. Veterinary Parasitology, 2008, 152, 148-151.	0.7	34
75	Microsatellite markers for the molecular characterization of Neospora caninum: Application to clinical samples. Veterinary Parasitology, 2009, 166, 38-46.	0.7	34
76	Neospora caninum seroprevalence in dairy and beef cattle from the northwest region of Spain, Galicia. Preventive Veterinary Medicine, 2011, 98, 128-132.	0.7	34
77	Proteome expression changes among virulent and attenuated Neospora caninum isolates. Journal of Proteomics, 2012, 75, 2306-2318.	1.2	34
78	Differential Responses of Bovine Monocyte-Derived Macrophages to Infection by Neospora caninum Isolates of High and Low Virulence. Frontiers in Immunology, 2019, 10, 915.	2.2	34
79	Immune response and protection provided by live tachyzoites and native antigens from the NC-6 Argentina strain of Neospora caninum in pregnant heifers. Veterinary Parasitology, 2013, 197, 436-446.	0.7	33
80	Ovine Toxoplasmosis: A New Look at its Pathogenesis. Journal of Comparative Pathology, 2017, 157, 34-38.	0.1	33
81	Toxoplasma gondii Genotyping: A Closer Look Into Europe. Frontiers in Cellular and Infection Microbiology, 2022, 12, 842595.	1.8	33
82	Molecular characterisation of BSR4, a novel bradyzoite-specific gene from Neospora caninum. International Journal for Parasitology, 2007, 37, 887-896.	1.3	32
83	Influence of Neospora caninum infection in BALB/c mice during pregnancy in post-natal development. Veterinary Parasitology, 2008, 155, 175-183.	0.7	32
84	Pathogenic characterization in mice of <i>Neospora caninum</i> isolates obtained from asymptomatic calves. Parasitology, 2010, 137, 1057-1068.	0.7	32
85	Serological evidence of Besnoitia spp. infection in Canadian wild ruminants and strong cross-reaction between Besnoitia besnoiti and Besnoitia tarandi. Veterinary Parasitology, 2012, 190, 19-28.	0.7	32
86	Systemic and local immune responses in sheep after Neospora caninum experimental infection at early, mid and late gestation. Veterinary Research, 2016, 47, 2.	1.1	32
87	SARS-CoV-2 Infection in One Cat and Three Dogs Living in COVID-19-Positive Households in Madrid, Spain. Frontiers in Veterinary Science, 2021, 8, 779341.	0.9	32
88	First description of naturally acquired Tritrichomonas foetus infection in a Persian cattery in Spain. Parasitology Research, 2011, 109, 1151-1154.	0.6	31
89	The first report of Cryptosporidium bovis, C. ryanae and Giardia duodenalis sub-assemblage A-II in roe deer (Capreolus capreolus) in Spain. Veterinary Parasitology, 2013, 197, 658-664.	0.7	31
90	First Report of Neospora caninum Infection in Adult Alpacas (Vicugna pacos) and Llamas (Lama glama). Journal of Parasitology, 2004, 90, 864-866.	0.3	30

#	Article	IF	CITATIONS
91	Neospora caninum infection in sheep and goats from north-eastern Italy and associated risk factors. Small Ruminant Research, 2016, 140, 7-12.	0.6	30
92	Differential susceptibility of bovine caruncular and trophoblast cell lines to infection with high and low virulence isolates of Neospora caninum. Parasites and Vectors, 2017, 10, 463.	1.0	30
93	High prevalence of Tritrichomonas foetus infection in Asturiana de la Montaña beef cattle kept in extensive conditions in Northern Spain. Veterinary Journal, 2012, 193, 146-151.	0.6	29
94	Effect of vaccination of cattle with the low virulence Nc-Spain 1H isolate of Neospora caninum against a heterologous challenge in early and mid-gestation. Veterinary Research, 2013, 44, 106.	1.1	29
95	First 2-DE approach towards characterising the proteome and immunome of Besnoitia besnoiti in the tachyzoite stage. Veterinary Parasitology, 2013, 195, 24-34.	0.7	29
96	Experimental ruminant models for bovine neosporosis: what is known and what is needed. Parasitology, 2014, 141, 1471-1488.	0.7	29
97	A live vaccine against Neospora caninum abortions in cattle. Vaccine, 2015, 33, 1299-1301.	1.7	29
98	Use of Avidity Enzyme-Linked Immunosorbent Assay and Avidity Western Blot to Discriminate between Acute and Chronic Neospora Caninum Infection in Cattle. Journal of Veterinary Diagnostic Investigation, 2005, 17, 442-450.	0.5	28
99	Presence of Cryptosporidium scrofarum, C. suis and C. parvum subtypes IIaA16G2R1 and IIaA13G1R1 in Eurasian wild boars (Sus scrofa). Veterinary Parasitology, 2013, 196, 497-502.	0.7	28
100	First serosurvey of Besnoitia spp. infection in wild European ruminants in Spain. Veterinary Parasitology, 2013, 197, 557-564.	0.7	28
101	Advances in the diagnosis of bovine besnoitiosis: current options and applications for control. International Journal for Parasitology, 2017, 47, 737-751.	1.3	28
102	Safety and efficacy of the bumped kinase inhibitor BKI-1553 in pregnant sheep experimentally infected with Neospora caninum tachyzoites. International Journal for Parasitology: Drugs and Drug Resistance, 2018, 8, 112-124.	1.4	28
103	Serum antibody response in lambs naturally and experimentally infected with Cryptosporidium parvum. Veterinary Parasitology, 1993, 50, 45-54.	0.7	27
104	Occasional detection of Neospora caninum DNA in frozen extended semen from naturally infected bulls. Theriogenology, 2004, 62, 1329-1336.	0.9	27
105	Comparison of host cell invasion and proliferation among Neospora caninum isolates obtained from oocysts and from clinical cases of naturally infected dogs. Experimental Parasitology, 2014, 145, 22-28.	0.5	27
106	Anti-Neospora caninum and anti-Sarcocystis spp. specific antibodies cross-react with Besnoitia besnoiti and influence the serological diagnosis of bovine besnoitiosis. Veterinary Parasitology, 2015, 214, 49-54.	0.7	27
107	The role of wild ruminants as reservoirs of Besnoitia besnoiti infection in cattle. Veterinary Parasitology, 2016, 223, 7-13.	0.7	27
108	CHARACTERIZATION OF PATHOLOGY AND PARASITE LOAD IN OUTBRED AND INBRED MOUSE MODELS OF CHRONIC NEOSPORA CANINUM INFECTION. Journal of Parasitology, 2004, 90, 579-583.	0.3	26

#	Article	IF	CITATIONS
109	Intrauterine Neospora caninum inoculation of heifers. Veterinary Parasitology, 2006, 135, 197-203.	0.7	26
110	Evaluation of Neospora caninum and Toxoplasma gondii infections in alpaca (Vicugna pacos) and llama (Lama glama) aborted foetuses from Peru. Veterinary Parasitology, 2007, 150, 39-45.	0.7	26
111	Identification of <i>Besnoitia besnoiti</i> proteins that showed differences in abundance between tachyzoite and bradyzoite stages by difference gel electrophoresis. Parasitology, 2013, 140, 999-1008.	0.7	26
112	Experimental caprine neosporosis: the influence of gestational stage on the outcome of infection. Veterinary Research, 2016, 47, 29.	1.1	26
113	The tandemly repeated NTPase (NTPDase) from Neospora caninum is a canonical dense granule protein whose RNA expression, protein secretion and phosphorylation coincides with the tachyzoite egress. Parasites and Vectors, 2016, 9, 352.	1.0	26
114	A survey of the prevalence of canine filariasis in Spain. Preventive Veterinary Medicine, 1991, 11, 63-68.	0.7	25
115	Toxoplasma gondii infection in adult llamas (Lama glama) and vicunas (Vicugna vicugna) in the Peruvian Andean region. Veterinary Parasitology, 2005, 130, 93-97.	0.7	25
116	Identification of <i>Neospora caninum</i> proteins regulated during the differentiation process from tachyzoite to bradyzoite stage by DIGE. Proteomics, 2010, 10, 1740-1750.	1.3	25
117	A vaccine formulation combining rhoptry proteins NcROP40 and NcROP2 improves pup survival in a pregnant mouse model of neosporosis. Veterinary Parasitology, 2015, 207, 203-215.	0.7	25
118	Experimental neosporosis in bulls: Parasite detection in semen and blood and specific antibody and interferon-gamma responses. Theriogenology, 2007, 67, 1175-1184.	0.9	24
119	Transgenic Neospora caninum strains constitutively expressing the bradyzoite NcSAG4 protein proved to be safe and conferred significant levels of protection against vertical transmission when used as live vaccines in mice. Vaccine, 2011, 29, 7867-7874.	1.7	24
120	Proteomics reveals differences in protein abundance and highly similar antigenic profiles between Besnoitia besnoiti and Besnoitia tarandi. Veterinary Parasitology, 2014, 205, 434-443.	0.7	24
121	Two Novel Calcium-Dependent Protein Kinase 1 Inhibitors Interfere with Vertical Transmission in Mice Infected with Neospora caninum Tachyzoites. Antimicrobial Agents and Chemotherapy, 2017, 61, .	1.4	24
122	Immune response profile of caruncular and trophoblast cell lines infected by high- (Nc-Spain7) and low-virulence (Nc-Spain1H) isolates of Neospora caninum. Parasites and Vectors, 2019, 12, 218.	1.0	24
123	Influence of adjuvant and antigen dose on protection induced by an inactivated whole vaccine against Neospora caninum infection in mice. Veterinary Parasitology, 2011, 175, 220-229.	0.7	23
124	Detection of Toxoplasma gondii antibodies in Antarctic pinnipeds. Veterinary Parasitology, 2012, 190, 259-262.	0.7	23
125	Integrative transcriptome and proteome analyses define marked differences between Neospora caninum isolates throughout the tachyzoite lytic cycle. Journal of Proteomics, 2018, 180, 108-119.	1.2	23
126	Treatment with Bumped Kinase Inhibitor 1294 Is Safe and Leads to Significant Protection against Abortion and Vertical Transmission in Sheep Experimentally Infected with Toxoplasma gondii during Pregnancy. Antimicrobial Agents and Chemotherapy, 2019, 63, .	1.4	23

#	Article	IF	CITATIONS
127	Low efficacy of NcGRA7, NcSAG4, NcBSR4 and NcSRS9 formulated in poly-É>-caprolactone against Neospora caninum infection in mice. Vaccine, 2012, 30, 4983-4992.	1.7	22
128	Presence and molecular characterisation of Giardia and Cryptosporidium in alpacas (Vicugna pacos) from Peru. Veterinary Parasitology, 2012, 187, 414-420.	0.7	22
129	Clinical outcome and vertical transmission variability among canine <i>Neospora caninum</i> isolates in a pregnant mouse model of infection. Parasitology, 2014, 141, 356-366.	0.7	22
130	Anthelmintic resistance in nematode parasites from goats in Spain. Veterinary Parasitology, 1997, 73, 83-88.	0.7	21
131	Is the anthelmintic effect of heather supplementation to grazing goats always accompanied by anti-nutritional effects?. Animal, 2008, 2, 1449-1456.	1.3	21
132	Experimental inoculation of Neospora caninum in pregnant water buffalo. Veterinary Parasitology, 2012, 187, 72-78.	0.7	21
133	Combination of monoclonal antibodies improves immunohistochemical diagnosis of Neospora caninum. Veterinary Parasitology, 2013, 197, 477-486.	0.7	21
134	Serological dynamics and risk factors of Besnoitia besnoiti infection in breeding bulls from an endemically infected purebred beef herd. Parasitology Research, 2017, 116, 1383-1393.	0.6	21
135	Seroprevalence of Toxoplasma gondii in Iberian pig sows. Parasitology Research, 2018, 117, 1419-1424.	0.6	21
136	Early Neospora caninum infection dynamics in cattle after inoculation at mid-gestation with high (Nc-Spain7)- or low (Nc-Spain1H)-virulence isolates. Veterinary Research, 2019, 50, 72.	1.1	21
137	Foetal death in naive heifers inoculated with Neospora caninum isolate Nc-Spain7 at 110 days of pregnancy. Experimental Parasitology, 2016, 168, 62-69.	0.5	20
138	A new lyophilized tachyzoite based ELISA to diagnose Besnoitia spp. infection in bovids and wild ruminants improves specificity. Veterinary Parasitology, 2017, 244, 176-182.	0.7	20
139	Global selective sweep of a highly inbred genome of the cattle parasite Neospora caninum. Proceedings of the National Academy of Sciences of the United States of America, 2019, 116, 22764-22773.	3.3	20
140	Lytic cycle of Besnoitia besnoiti tachyzoites displays similar features in primary bovine endothelial cells and fibroblasts. Parasites and Vectors, 2019, 12, 517.	1.0	20
141	Isolation and genetic characterization of Toxoplasma gondii in Spanish sheep flocks. Parasites and Vectors, 2020, 13, 396.	1.0	20
142	Serum and bile antibody responses (IgG and IgA) during subclinical Fasciola hepatica infection in sheep. Veterinary Parasitology, 1997, 68, 261-267.	0.7	19
143	Stage-specific expression of Nc <i>SAG4</i> as a marker of chronic <i>Neospora caninum</i> infection in a mouse model. Parasitology, 2009, 136, 757-764.	0.7	19
144	Evaluation of the protection conferred by a naturally attenuated Neospora caninum isolate against congenital and cerebral neosporosis in mice. Veterinary Research, 2012, 43, 62.	1.1	19

#	Article	IF	CITATIONS
145	Prevalence of Besnoitia besnoiti infection in beef cattle from the Spanish Pyrenees. Veterinary Journal, 2014, 200, 468-470.	0.6	19
146	Seroprevalence of Besnoitia besnoiti infection and associated risk factors in cattle from an endemic region in Europe. Veterinary Journal, 2014, 200, 328-331.	0.6	19
147	Cell mediated immune responses in the placenta following challenge of vaccinated pregnant heifers with Neospora caninum. Veterinary Parasitology, 2015, 214, 247-254.	0.7	19
148	Immunization with a cocktail of antigens fused with OprI reduces Neospora caninum vertical transmission and postnatal mortality in mice. Vaccine, 2019, 37, 473-483.	1.7	19
149	Comparison of Neospora caninum distribution, parasite loads and lesions between epidemic and endemic bovine abortion cases. Veterinary Parasitology, 2006, 142, 187-191.	0.7	18
150	Identification of novel rhoptry proteins in Neospora caninum by LC/MS-MS analysis of subcellular fractions. Journal of Proteomics, 2011, 74, 629-642.	1.2	18
151	Detection and Characterization of a Cryptosporidium Isolate from a Southern Elephant Seal () Tj ETQq1 1 0.7843 1524-1527.	314 rgBT / 1.4	Overlock 10 18
152	In vitro effect of heather (Ericaceae) extracts on different development stages of Teladorsagia circumcincta and Haemonchus contortus. Veterinary Parasitology, 2013, 197, 235-243.	0.7	18
153	Coxiella burnetii in dairy goats with a history of reproductive disorders in Brazil. Acta Tropica, 2018, 183, 19-22.	0.9	18
154	Neospora caninum IgG avidity tests: An interlaboratory comparison. Veterinary Parasitology, 2006, 140, 273-280.	0.7	17
155	Potential use of heather to control gastrointestinal nematodes in goats. Small Ruminant Research, 2012, 103, 60-68.	0.6	17
156	Clinical and Serological Dynamics of <i>Besnoitia besnoiti</i> Infection in Three Endemically Infected Beef Cattle Herds. Transboundary and Emerging Diseases, 2017, 64, 538-546.	1.3	17
157	Neospora caninum: Structure and Fate of Multinucleated Complexes Induced by the Bumped Kinase Inhibitor BKI-1294. Pathogens, 2020, 9, 382.	1.2	17
158	In vitro activity, safety and in vivo efficacy of the novel bumped kinase inhibitor BKI-1748 in non-pregnant and pregnant mice experimentally infected with Neospora caninum tachyzoites and Toxoplasma gondii oocysts. International Journal for Parasitology: Drugs and Drug Resistance, 2021, 16–90-101	1.4	17
159	Identification of Cryptosporidium parvum oocyst/sporozoite antigens recognized by infected and hyperimmune lambs. Veterinary Parasitology, 1994, 53, 159-166.	0.7	16
160	Intrauterine Neospora caninum inoculation of heifers and cows using contaminated semen with different numbers of tachyzoites. Theriogenology, 2007, 67, 729-737.	0.9	16
161	The effect of heather supplementation on gastrointestinal nematode infections and performance in Cashmere and local Celtiberic goats on pasture. Small Ruminant Research, 2007, 67, 184-191.	0.6	16
162	Effects of heather and oat supplementation on gastrointestinal nematode infections and performance of grazing Cashmere goats. Small Ruminant Research, 2010, 91, 186-192.	0.6	16

#	Article	IF	CITATIONS
163	Isolation and Characterization of <i>Campylobacter</i> spp. from Antarctic Fur Seals () Tj ETQq1 1 0.784314 rgB	[ /Overlock 1.4	2 10 Tf 50
	Microbiology, 2010, 76, 6013-6016.		
164	Effects of Neospora caninum Infection at Mid-Gestation on Placenta in a Pregnant Mouse Model. Journal of Parasitology, 2010, 96, 1017-1020.	0.3	16
165	EFFECT OF DIFFERENT ECOSYSTEMS AND MANAGEMENT PRACTICES ON <i>TOXOPLASMA GONDII </i> AND <i>NEOSPORA CANINUM </i> INFECTIONS IN WILD RUMINANTS IN SPAIN. Journal of Wildlife Diseases, 2016, 52, 293-300.	0.3	16
166	Toxoplasma CRISPR/Cas9 constructs are functional for gene disruption in Neospora caninum. International Journal for Parasitology, 2018, 48, 597-600.	1.3	16
167	One health therapeutics: Target-Based drug development for cryptosporidiosis and other apicomplexa diseases. Veterinary Parasitology, 2021, 289, 109336.	0.7	16
168	The role of Neospora caninum and Toxoplasma gondii in spontaneous bovine abortion in Argentina. Veterinary Parasitology, 2008, 156, 163-167.	0.7	15
169	An Ibero-American inter-laboratory trial to evaluate serological tests for the detection of anti-Neospora caninum antibodies in cattle. Tropical Animal Health and Production, 2018, 50, 75-84.	0.5	15
170	Isolation, Genotyping, and Mouse Virulence Characterization of Toxoplasma gondii From Free Ranging Iberian Pigs. Frontiers in Veterinary Science, 2020, 7, 604782.	0.9	15
171	Neospora caninum: Differential Proteome of Multinucleated Complexes Induced by the Bumped Kinase Inhibitor BKI-1294. Microorganisms, 2020, 8, 801.	1.6	15
172	The acid phosphatase activity and morphological characteristics of Dipetalonema dracunculoides (cobbold, 1870) microfilariae. Veterinary Parasitology, 1989, 33, 187-190.	0.7	14
173	Prevalance of Dicrocoelium dendriticum infection in sheep in LeÃ <sup>3</sup> n province (NW Spain). Preventive Veterinary Medicine, 1994, 21, 147-154.	0.7	14
174	Prevalence of Tritrichomonas foetus infection in beef bulls in northwestern Spain. Veterinary Parasitology, 1998, 75, 265-268.	0.7	14
175	Neospora caninum infection in breeder bulls: seroprevalence and comparison of serological methods used for diagnosis. Veterinary Parasitology, 2004, 124, 19-24.	0.7	14
176	Effects of re-infection with Neospora caninum in bulls on parasite detection in semen and blood and immunological responses. Theriogenology, 2008, 69, 905-911.	0.9	14
177	Characterisation of NcGRA7 and NcSAG4 proteins: Immunolocalisation and their role in the host cell invasion by Neospora caninum tachyzoites. Acta Parasitologica, 2010, 55, .	0.4	14
178	In vitro effect of heather extracts on Trichostrongylus colubriformis eggs, larvae and adults. Veterinary Parasitology, 2013, 197, 586-594.	0.7	14
179	Anthelmintic effect of heather in goats experimentally infected with Trichostrongylus colubriformis. Parasitology Research, 2014, 113, 693-699.	0.6	14
180	Tritrichomonas foetus infection in cats with diarrhea from densely housed origins. Veterinary Parasitology, 2016, 221, 118-122.	0.7	14

#	Article	IF	CITATIONS
181	Systemic Besnoitiosis in a Juvenile Roe Deer ( <i>Capreolus capreolus</i> ). Transboundary and Emerging Diseases, 2017, 64, e8-e14.	1.3	14
182	A serosurvey of selected cystogenic coccidia in Spanish equids: first detection of anti-Besnoitia spp. specific antibodies in Europe. BMC Veterinary Research, 2017, 13, 128.	0.7	14
183	Peripheral and placental immune responses in sheep after experimental infection with Toxoplasma gondii at the three terms of gestation. Veterinary Research, 2019, 50, 66.	1.1	14
184	In vivo and in vitro models show unexpected degrees of virulence among Toxoplasma gondii type II and III isolates from sheep. Veterinary Research, 2021, 52, 82.	1.1	14
185	Identification of a gene cluster for cell-surface genes of the SRS superfamily in <i>Neospora caninum</i> and characterization of the novel <i>SRS9</i> gene. Parasitology, 2011, 138, 1832-1842.	0.7	13
186	Detection of a novel genotype of Cryptosporidium in Antarctic pinnipeds. Veterinary Parasitology, 2013, 191, 112-118.	0.7	13
187	Repurposing of commercially available anti-coccidials identifies diclazuril and decoquinate as potential therapeutic candidates against Besnoitia besnoiti infection. Veterinary Parasitology, 2018, 261, 77-85.	0.7	13
188	Absence of <i>Neospora caninum</i> DNA in Human Clinical Samples, Spain. Emerging Infectious Diseases, 2019, 25, 1226-1227.	2.0	13
189	Comparative efficacy of immunization with inactivated whole tachyzoites versus a tachyzoite-bradyzoite mixture against neosporosis in mice. Parasitology, 2011, 138, 1372-1383.	0.7	12
190	Efficacy of a control program for bovine trichomonosis based on testing and culling infected bulls in beef cattle managed under mountain pastoral systems of Northern Spain. Veterinary Journal, 2014, 200, 140-145.	0.6	12
191	Characterization of the <i>Neospora caninum</i> NcROP40 and NcROP2Fam-1 rhoptry proteins during the tachyzoite lytic cycle. Parasitology, 2016, 143, 97-113.	0.7	12
192	Gene Expression Profiling of Neospora caninum in Bovine Macrophages Reveals Differences Between Isolates Associated With Key Parasite Functions. Frontiers in Cellular and Infection Microbiology, 2019, 9, 354.	1.8	12
193	Crosstalk between Neospora caninum and the bovine host at the maternal-foetal interface determines the outcome of infection. Veterinary Research, 2020, 51, 83.	1.1	12
194	Molecular survey for cyst-forming coccidia (Toxoplasma gondii, Neospora caninum, Sarcocystis spp.) in Mediterranean periurban micromammals. Parasitology Research, 2020, 119, 2679-2686.	0.6	12
195	Dynamics of Neospora caninum-Associated Abortions in a Dairy Sheep Flock and Results of a Test-and-Cull Control Programme. Pathogens, 2021, 10, 1518.	1.2	12
196	Contamination of Soil, Water, Fresh Produce, and Bivalve Mollusks with Toxoplasma gondii Oocysts: A Systematic Review. Microorganisms, 2022, 10, 517.	1.6	12
197	Effect of different decoquinate treatments on cryptosporidiosis in naturally infected cashmere goat kids. Veterinary Record, 2005, 157, 261-262.	0.2	11
198	Isolation and biological characterisation of a new isolate of Neospora caninum from an asymptomatic calf in Brazil. Acta Parasitologica, 2009, 54, .	0.4	11

#	Article	IF	CITATIONS
199	Genetic manipulation of Neospora caninum to express the bradyzoite-specific protein NcSAG4 in tachyzoites. Parasitology, 2011, 138, 472-480.	0.7	11
200	Genetic characterisation of Neospora caninum strains from clinical samples of zebuine foetuses obtained in abattoirs in GoiÃ <sub>i</sub> s, Brazil. Veterinary Parasitology, 2014, 204, 381-387.	0.7	11
201	Genetic characterization of Neospora caninum from aborted bovine foetuses in Aguascalientes, Mexico. Veterinary Parasitology, 2016, 228, 183-187.	0.7	11
202	Bovine chronic besnoitiosis in a calf: Characterization of a novel B. besnoiti isolate from an unusual case report. Veterinary Parasitology, 2017, 247, 10-18.	0.7	11
203	Microsatellite pattern analysis of Neospora caninum from a naturally infected goat fetus. Veterinary Parasitology, 2018, 255, 58-60.	0.7	11
204	Effects of challenge dose and inoculation route of the virulent Neospora caninum Nc-Spain7 isolate in pregnant cattle at mid-gestation. Veterinary Research, 2019, 50, 68.	1.1	11
205	Neospora caninum infection in stray and farm dogs: Seroepidemiological study and oocyst shedding. Veterinary Parasitology, 2010, 174, 332-335.	0.7	10
206	Specific antibody responses against Neospora caninum recombinant rNcGRA7, rNcSAG4, rNcBSR4 and rNcSRS9 proteins are correlated with virulence in mice. Parasitology, 2013, 140, 569-579.	0.7	10
207	Low rates of <i>Neospora caninum</i> infection reactivation during gestation are observed in both chronically and congenitally infected mice. Parasitology, 2013, 140, 220-228.	0.7	10
208	Neospora caninum infection induces an isolate virulence-dependent pro-inflammatory gene expression profile in bovine monocyte-derived macrophages. Parasites and Vectors, 2020, 13, 374.	1.0	10
209	Comparative tachyzoite proteome analyses among six Neospora caninum isolates with different virulence. International Journal for Parasitology, 2020, 50, 377-388.	1.3	10
210	Vascular wall injury and inflammation are key pathogenic mechanisms responsible for early testicular degeneration during acute besnoitiosis in bulls. Parasites and Vectors, 2020, 13, 113.	1.0	10
211	Modeling the Ruminant Placenta-Pathogen Interactions in Apicomplexan Parasites: Current and Future Perspectives. Frontiers in Veterinary Science, 2020, 7, 634458.	0.9	10
212	Unifying Virulence Evaluation in Toxoplasma gondii: A Timely Task. Frontiers in Cellular and Infection Microbiology, 2022, 12, 868727.	1.8	10
213	Effects of Stocking Rate and Heather Supplementation on Gastrointestinal Nematode Infections and Host Performance in Naturally-Infected Cashmere Goats. Rangeland Ecology and Management, 2009, 62, 127-135.	1.1	9
214	Effect of the consumption of heather on incoming larvae and established population of Teladorsagia circumcincta in experimentally infected Cashmere goats. Veterinary Parasitology, 2013, 196, 124-129.	0.7	9
215	Presence of Ostertagia ostertagi antibodies in bulk tank milk from cattle herds in northern Spain. Veterinary Parasitology, 2013, 197, 388-392.	0.7	9
216	Mice congenitally infected with low-to-moderate virulence Neospora caninum isolates exhibited clinical reactivation during the mating period without transmission to the next generation. Experimental Parasitology, 2013, 134, 244-248.	0.5	9

#	Article	IF	CITATIONS
217	Seroprevalence of Leptospirosis, Brucellosis, and Q Fever in a Wild Red Deer ( <i>Cervus elaphus</i> ) Population Kept in a Fenced Reserve in Absence of Contact with Livestock. Vector-Borne and Zoonotic Diseases, 2017, 17, 692-697.	0.6	9
218	Characterization of Fetal Brain Damage in Early Abortions of Ovine Toxoplasmosis. Veterinary Pathology, 2020, 57, 535-544.	0.8	9
219	Neospora caninum tachyzoite immunome study reveals differences among three biologically different isolates. Veterinary Parasitology, 2015, 212, 92-99.	0.7	8
220	First description of clonal lineage type II (genotype #1) of Toxoplasma gondii in abortion outbreak in goats. Experimental Parasitology, 2018, 188, 21-25.	0.5	8
221	Exposure to Neospora spp. and Besnoitia spp. in wildlife from Israel. International Journal for Parasitology: Parasites and Wildlife, 2018, 7, 317-321.	0.6	8
222	Influence of dose and route of administration on the outcome of infection with the virulent Neospora caninum isolate Nc-Spain7 in pregnant sheep at mid-gestation. Veterinary Research, 2018, 49, 42.	1.1	8
223	Macrophages and T Lymphocytes in the Ovine Placenta After Experimental Infection With <i>Toxoplasma gondii</i> . Veterinary Pathology, 2020, 57, 545-549.	0.8	8
224	Genetic characterization of Neospora caninum from Northern Italian cattle reveals high diversity in European N. caninum populations. Parasitology Research, 2020, 119, 1353-1362.	0.6	8
225	Identification of molecular biomarkers associated with disease progression in the testis of bulls infected with Besnoitia besnoiti. Veterinary Research, 2021, 52, 106.	1.1	8
226	Use of an immunodominant p17 antigenic fraction of Neospora caninum in detection of antibody response in cattle. Memorias Do Instituto Oswaldo Cruz, 2006, 101, 529-534.	0.8	7
227	Natural breeding with bulls experimentally infected with Neospora caninum failed to induce seroconversion in dams. Theriogenology, 2009, 71, 639-642.	0.9	7
228	Helminth parasites found in faecal samples of phocids from the Antarctic Peninsula. Polar Biology, 2014, 37, 685-695.	0.5	7
229	Isolation and biological and molecular characterization of Neospora caninum (NC-SP1) from a naturally infected adult asymptomatic cattle (Bos taurus) in the state of São Paulo, Brazil. Parasitology, 2017, 144, 707-711.	0.7	7
230	Prevalence of bovine trichomonosis and associated risk factors in bulls from Spanish beef herds. Theriogenology, 2019, 128, 116-121.	0.9	7
231	Proteomic Characterization of Host-Pathogen Interactions during Bovine Trophoblast Cell Line Infection by Neospora caninum. Pathogens, 2020, 9, 749.	1.2	7
232	The Impact of BKI-1294 Therapy in Mice Infected With the Apicomplexan Parasite Neospora caninum and Re-infected During Pregnancy. Frontiers in Veterinary Science, 2020, 7, 587570.	0.9	7
233	Endochin-like quinolones (ELQs) and bumped kinase inhibitors (BKIs): Synergistic and additive effects of combined treatments against Neospora caninum infection in vitro and in vivo. International Journal for Parasitology: Drugs and Drug Resistance, 2021, 17, 92-106.	1.4	7
234	Abortions in bovines and Neospora caninum transmission in an embryo transfer center. Veterinary Parasitology, 2010, 173, 206-210.	0.7	6

#	Article	IF	CITATIONS
235	Differences in the prevalence of Tritrichomonas foetus infection in beef cattle farmed under extensive conditions in northern Spain. Veterinary Journal, 2013, 196, 547-549.	0.6	6
236	Neospora caninum tachyzoites inoculated by the conjunctival route are not vertically transmitted in pregnant cattle: A descriptive study. Veterinary Parasitology, 2014, 199, 1-7.	0.7	6
237	Health impact evaluation of alternative management systems in vicuña (Vicugna vicugna mensalis) populations in Peru. Tropical Animal Health and Production, 2014, 46, 641-646.	0.5	6
238	Immunohistochemical study and mRNA cytokine profile of the local immune response in cattle naturally infected with Calicophoron daubneyi. Veterinary Parasitology, 2015, 214, 178-183.	0.7	6
239	Absence of antibodies specific to Besnoitia spp. in European sheep and goats from areas in Spain where bovine besnoitiosis is endemic. Parasitology Research, 2017, 116, 445-448.	0.6	6
240	Isolation of Neospora caninum from kidney and brain of a bovine foetus and molecular characterization in Brazil. Experimental Parasitology, 2018, 185, 10-16.	0.5	6
241	Effect of parasite dose and host age on the infection with Besnoitia besnoiti tachyzoites in cattle. Transboundary and Emerging Diseases, 2018, 65, 1979-1990.	1.3	6
242	Isolation and genetic characterization of Neospora caninum from naturally infected sheep. Veterinary Parasitology, 2020, 280, 109091.	0.7	6
243	Assessment of the Activity of Decoquinate and Its Quinoline-O-Carbamate Derivatives against Toxoplasma gondii In Vitro and in Pregnant Mice Infected with T. gondii Oocysts. Molecules, 2021, 26, 6393.	1.7	6
244	Prevalence of Bovine Genital Campylobacteriosis, Associated Risk Factors and Spatial Distribution in Spanish Beef Cattle Based on Veterinary Laboratory Database Records. Frontiers in Veterinary Science, 2021, 8, 750183.	0.9	6
245	Apparent absence of <i>Cryptosporidium, Giardia</i> and <i>Toxoplasma gondii</i> in three species of penguins along the Antarctic Peninsula. Antarctic Science, 2010, 22, 265-270.	0.5	5
246	IS Q fever a significant cause of reproductive failure in cattle?. Veterinary Record, 2012, 170, 257-258.	0.2	5
247	Immune response to Neospora caninum live tachyzoites in prepubertal female calves. Parasitology Research, 2019, 118, 2945-2955.	0.6	5
248	A model for chronic bovine besnoitiosis: Parasite stage and inoculation route are key factors. Transboundary and Emerging Diseases, 2020, 67, 234-249.	1.3	5
249	Maternal and Foetal Cellular Immune Responses in Dams Infected With High- and Low- Virulence Isolates of Neospora caninum at Mid-Gestation. Frontiers in Cellular and Infection Microbiology, 2021, 11, 684670.	1.8	5
250	Direct economic losses of Toxoplasma gondii abortion outbreaks in two Spanish sheep flocks. Veterinary Parasitology: Regional Studies and Reports, 2021, 26, 100623.	0.3	5
251	A short-term treatment with BKI-1294 does not protect foetuses from sheep experimentally infected with Neospora caninum tachyzoites during pregnancy. International Journal for Parasitology: Drugs and Drug Resistance, 2021, 17, 176-185.	1.4	5
252	Prevalence of intestinal parasite infections in stray and farm dogs from Spain. Brazilian Journal of Veterinary Parasitology, 2020, 29, e014920.	0.2	5

#	Article	IF	CITATIONS
253	Common Molecular Targets of a Quinolone Based Bumped Kinase Inhibitor in Neospora caninum and Danio rerio. International Journal of Molecular Sciences, 2022, 23, 2381.	1.8	5
254	Peripheral and placental immune responses in goats after primoinfection with Neospora caninum at early, mid and late gestation. Veterinary Parasitology, 2017, 242, 38-43.	0.7	4
255	Trichomonas. , 2018, , 313-388.		4
256	The route of Besnoitia besnoiti tachyzoites inoculation does not influence the clinical outcome of the infection in calves. Veterinary Parasitology, 2019, 267, 21-25.	0.7	4
257	Microsatellite genotyping reveals extensive genetic diversity in bovine Neospora caninum from the humid Pampa region in Argentina. Parasitology Research, 2020, 119, 4049-4059.	0.6	4
258	Toxoplasma gondii and Neospora caninum seroprevalences in domestic South American camelids of the Peruvian Andes. Tropical Animal Health and Production, 2014, 46, 1141-1147.	0.5	3
259	Added value of IgM detection and low avidity index as markers of acute bovine besnoitiosis. Veterinary Parasitology, 2020, 277, 109012.	0.7	3
260	Histological findings in experimentally infected male calves with chronic besnoitiosis. Veterinary Parasitology, 2020, 281, 109120.	0.7	3
261	Development and characterization of monoclonal antibodies against <i>Besnoitia besnoiti</i> tachyzoites. Parasitology, 2019, 146, 187-196.	0.7	2
262	Multilocus analysis reveals further genetic differences between Tritrichomonas foetus from cats and cattle. Veterinary Parasitology, 2019, 276, 108965.	0.7	2
263	Morphometric study of encephalic lesions in aborted bovine fetuses naturally infected by two subpopulations of Neospora caninum. Parasitology Research, 2021, 120, 2995-3000.	0.6	2
264	Changes in serum biomarkers of inflammation in bovine besnoitiosis. Parasites and Vectors, 2021, 14, 488.	1.0	2
265	Vaccine-Linked Chemotherapy Approach: Additive Effects of Combining the Listeria monocytogenes-Based Vaccine Lm3Dx_NcSAG1 With the Bumped Kinase Inhibitor BKI-1748 Against Neospora caninum Infection in Mice. Frontiers in Veterinary Science, 0, 9, .	0.9	2
266	Controlled field efficacy of injectable moxidectin against naturally acquired psoroptic mange in sheep. Small Ruminant Research, 1998, 29, 271-276.	0.6	1
267	Pathological and immunological findings in placentas from pregnant BALB/c mice infected with Neospora caninum at early and late stages of gestation. Acta Parasitologica, 2011, 56, .	0.4	1
268	Parasitemia and Associated Immune Response in Pregnant and Non-Pregnant Beef Cows Naturally Infected With Neospora caninum. Frontiers in Veterinary Science, 0, 9, .	0.9	1