

Qingguang Liu

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/479459/publications.pdf>

Version: 2024-02-01

12
papers

981
citations

759233

12
h-index

1125743

13
g-index

13
all docs

13
docs citations

13
times ranked

1013
citing authors

#	ARTICLE	IF	CITATIONS
1	Long non-coding RNA MAPKAPK5-AS1/PLAGL2/HIF-1 \pm signaling loop promotes hepatocellular carcinoma progression. <i>Journal of Experimental and Clinical Cancer Research</i> , 2021, 40, 72.	8.6	50
2	Histone citrullination by PADI4 is required for HIF-dependent transcriptional responses to hypoxia and tumor vascularization. <i>Science Advances</i> , 2021, 7, .	10.3	31
3	HIF-1 \pm -activated long non-coding RNA KDM4A-AS1 promotes hepatocellular carcinoma progression via the miR-411-5p/KPNA2/AKT pathway. <i>Cell Death and Disease</i> , 2021, 12, 1152.	6.3	39
4	Hypoxia-inducible long noncoding RNA NPSR1-AS1 promotes the proliferation and glycolysis of hepatocellular carcinoma cells by regulating the MAPK/ERK pathway. <i>Biochemical and Biophysical Research Communications</i> , 2020, 533, 886-892.	2.1	17
5	Hypoxia-induced miR-3677-3p promotes the proliferation, migration and invasion of hepatocellular carcinoma cells by suppressing SIRT5. <i>Journal of Cellular and Molecular Medicine</i> , 2020, 24, 8718-8731.	3.6	22
6	LncRNA RUNX1-IT1 which is downregulated by hypoxia-driven histone deacetylase 3 represses proliferation and cancer stem-like properties in hepatocellular carcinoma cells. <i>Cell Death and Disease</i> , 2020, 11, 95.	6.3	67
7	MicroRNA-875-5p inhibits tumor growth and metastasis of hepatocellular carcinoma by targeting eukaryotic translation initiation factor 3 subunit a. <i>Oncology Reports</i> , 2020, 44, 2067-2079.	2.6	8
8	Hypoxia-induced TUFT1 promotes the growth and metastasis of hepatocellular carcinoma by activating the Ca ²⁺ /PI3K/AKT pathway. <i>Oncogene</i> , 2019, 38, 1239-1255.	5.9	108
9	Hypoxia-induced up-regulation of VASP promotes invasiveness and metastasis of hepatocellular carcinoma. <i>Theranostics</i> , 2018, 8, 4649-4663.	10.0	120
10	Long non-coding RNA DSCR8 acts as a molecular sponge for miR-485-5p to activate Wnt/ β -catenin signal pathway in hepatocellular carcinoma. <i>Cell Death and Disease</i> , 2018, 9, 851.	6.3	110
11	Long non-coding RNA CASC2 suppresses epithelial-mesenchymal transition of hepatocellular carcinoma cells through CASC2/miR-367/FBXW7 axis. <i>Molecular Cancer</i> , 2017, 16, 123.	19.2	200
12	miR-187-3p inhibits the metastasis and epithelial-mesenchymal transition of hepatocellular carcinoma by targeting S100A4. <i>Cancer Letters</i> , 2016, 381, 380-390.	7.2	104