

Dinglan Wu

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4817064/publications.pdf>

Version: 2024-02-01

24
papers

702
citations

516710

16
h-index

610901

24
g-index

25
all docs

25
docs citations

25
times ranked

1162
citing authors

#	ARTICLE	IF	CITATIONS
1	Endothelial nitric oxide synthase (eNOS)-NO signaling axis functions to promote the growth of prostate cancer stem-like cells. <i>Stem Cell Research and Therapy</i> , 2022, 13, 188.	5.5	7
2	Orphan nuclear receptors as regulators of intratumoral androgen biosynthesis in castration-resistant prostate cancer. <i>Oncogene</i> , 2021, 40, 2625-2634.	5.9	19
3	Orphan nuclear receptor TLX promotes immunosuppression via its transcriptional activation of PD-L1 in glioma. <i>Stem Cell Research and Therapy</i> , 2021, 9, e001937.		15
4	Nuclear receptor HNF4 α performs a tumor suppressor function in prostate cancer via its induction of p21-driven cellular senescence. <i>Oncogene</i> , 2020, 39, 1572-1589.	5.9	27
5	Targeting prostate cancer stem-like cells by an immunotherapeutic platform based on immunogenic peptide-sensitized dendritic cells-cytokine-induced killer cells. <i>Stem Cell Research and Therapy</i> , 2020, 11, 123.	5.5	16
6	Nuclear factor IX promotes glioblastoma development through transcriptional activation of Ezrin. <i>Oncogenesis</i> , 2020, 9, 39.	4.9	11
7	In Vitro and In Vivo Antitumor Activity of Cucurbitacin C, a Novel Natural Product From Cucumber. <i>Frontiers in Pharmacology</i> , 2019, 10, 1287.	3.5	32
8	miR-182-5p affects human bladder cancer cell proliferation, migration and invasion through regulating Cofilin 1. <i>Cancer Cell International</i> , 2019, 19, 42.	4.1	32
9	Long non-coding RNA HOXA-AS2 promotes the migration, invasion and stemness of bladder cancer via regulating miR-125b/Smad2 axis. <i>Experimental Cell Research</i> , 2019, 375, 1-10.	2.6	41
10	Orphan nuclear receptor TLX contributes to androgen insensitivity in castration-resistant prostate cancer via its repression of androgen receptor transcription. <i>Oncogene</i> , 2018, 37, 3340-3355.	5.9	20
11	Nuclear Receptor LRH-1 Functions to Promote Castration-Resistant Growth of Prostate Cancer via Its Promotion of Intratumoral Androgen Biosynthesis. <i>Cancer Research</i> , 2018, 78, 2205-2218.	0.9	36
12	Hyperglycaemia-induced miR-301a promotes cell proliferation by repressing p21 and Smad4 in prostate cancer. <i>Cancer Letters</i> , 2018, 418, 211-220.	7.2	40
13	Nuclear receptor profiling in prostatospheroids and castration-resistant prostate cancer. <i>Endocrine-Related Cancer</i> , 2018, 25, 35-50.	3.1	24
14	Development of a novel and economical agar-based non-adherent three-dimensional culture method for enrichment of cancer stem-like cells. <i>Stem Cell Research and Therapy</i> , 2018, 9, 243.	5.5	48
15	Nuclear receptor ERR α and transcription factor ERG form a reciprocal loop in the regulation of TMPRSS2:ERG fusion gene in prostate cancer. <i>Oncogene</i> , 2018, 37, 6259-6274.	5.9	36
16	Collapsin response mediator protein-1 (CRMP1) acts as an invasion and metastasis suppressor of prostate cancer via its suppression of epithelial-mesenchymal transition and remodeling of actin cytoskeleton organization. <i>Oncogene</i> , 2017, 36, 546-558.	5.9	38
17	Study on the Significance of Cofilin 1 Overexpression in Human Bladder Cancer. <i>Tumori</i> , 2017, 103, 537-542.	1.1	7
18	<i>IL1</i> genes polymorphism and the risk of renal cell carcinoma in Chinese Han population. <i>Oncotarget</i> , 2017, 8, 56021-56029.	1.8	7

#	ARTICLE	IF	CITATIONS
19	Cofilin 1 promotes bladder cancer and is regulated by TCF7L2. <i>Oncotarget</i> , 2017, 8, 92043-92054.	1.8	12
20	The emerging roles of orphan nuclear receptors in prostate cancer. <i>Biochimica Et Biophysica Acta: Reviews on Cancer</i> , 2016, 1866, 23-36.	7.4	23
21	Orphan nuclear receptor <sc>TLX</sc> functions as a potent suppressor of oncogene-induced senescence in prostate cancer via its transcriptional co-regulation of the <i><sc>CDKN1A</sc></i> (<sc>p21^{WAF1}</sc>[/]<sc>^{CIP1}</sc>) and <i><sc>SIRT1</sc></i> genes. <i>Journal of Pathology</i> , 2015, 236, 103-115.	4.5	40
22	<sc>ERR</sc><i>1</i> augments <sc>HIF</sc><i>1</i> signalling by directly interacting with <sc>HIF</sc><i>1</i> in normoxic and hypoxic prostate cancer cells. <i>Journal of Pathology</i> , 2014, 233, 61-73.	4.5	72
23	Ion channel <sc>TRPM8</sc> promotes hypoxic growth of prostate cancer cells via an <sc>O₂</sc>-independent and <sc>RACK1</sc>-mediated mechanism of <sc>HIF</sc><i>1</i> stabilization. <i>Journal of Pathology</i> , 2014, 234, 514-525.	4.5	53
24	Increased expression of activated endothelial nitric oxide synthase contributes to antiandrogen resistance in prostate cancer cells by suppressing androgen receptor transactivation. <i>Cancer Letters</i> , 2013, 328, 83-94.	7.2	34