

# Zbigniew Dominski

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/48325/publications.pdf>

Version: 2024-02-01

33  
papers

1,704  
citations

361413  
20  
h-index

395702  
33  
g-index

33  
all docs

33  
docs citations

33  
times ranked

1308  
citing authors

| #  | ARTICLE  | IF   | CITATIONS |
|----|--|------|-----------|
| 1  | Reconstitution and biochemical assays of an active human histone pre-mRNA 3' end processing machinery. <i>Methods in Enzymology</i> , 2021, 655, 291-324.  | 1.0  | 7         |
| 2  | Superresolution light microscopy of the <i>Drosophila</i> histone locus body reveals a core-shell organization associated with expression of replication-dependent histone genes. <i>Molecular Biology of the Cell</i> , 2021, 32, 942-955.      | 2.1  | 15        |
| 3  | U7 deciphered: the mechanism that forms the unusual 3' end of metazoan replication-dependent histone mRNAs. <i>Biochemical Society Transactions</i> , 2021, 49, 2229-2240.   | 3.4  | 8         |
| 4  | Composition and processing activity of a semi-recombinant holo U7 snRNP. <i>Nucleic Acids Research</i> , 2020, 48, 1508-1530.  | 14.5 | 13        |
| 5  | Structural Analysis of the SANT/Myb Domain of FLASH and YARP Proteins and Their Complex with the C-Terminal Fragment of NPAT by NMR Spectroscopy and Computer Simulations. <i>International Journal of Molecular Sciences</i> , 2020, 21, 5268.  | 4.1  | 5         |
| 6  | Studies with recombinant U7 snRNP demonstrate that CPSF73 is both an endonuclease and a 5' endonuclease. <i>Rna</i> , 2020, 26, 1345-1359.   | 3.5  | 20        |
| 7  | Structure of an active human histone pre-mRNA 3' end processing machinery. <i>Science</i> , 2020, 367, 700-703.  | 12.6 | 76        |
| 8  | Single-step Purification of Macromolecular Complexes Using RNA Attached to Biotin and a Photo-cleavable Linker. <i>Journal of Visualized Experiments</i> , 2019, , .   | 0.3  | 3         |
| 9  | Protein composition of catalytically active U7-dependent processing complexes assembled on histone pre-mRNA containing biotin and a photo-cleavable linker. <i>Nucleic Acids Research</i> , 2018, 46, 4752-4770.                                 | 14.5 | 21        |
| 10 | U7 snRNP is recruited to histone pre-mRNA in a FLASH-dependent manner by two separate regions of the stem-loop binding protein. <i>Rna</i> , 2017, 23, 938-951.  | 3.5  | 26        |
| 11 | The N-terminal domains of FLASH and Lsm11 form a 2:1 heterotrimer for histone pre-mRNA 3' end processing. <i>PLoS ONE</i> , 2017, 12, e0186034.  | 2.5  | 12        |
| 12 | Mapping the Interaction Network of Key Proteins Involved in Histone mRNA Generation: A Hydrogen/Deuterium Exchange Study. <i>Journal of Molecular Biology</i> , 2016, 428, 1180-1196.  | 4.2  | 8         |
| 13 | Concentrating pre-mRNA processing factors in the histone locus body facilitates efficient histone mRNA biogenesis. <i>Journal of Cell Biology</i> , 2016, 213, 557-570.  | 5.2  | 75        |
| 14 | A Conserved Interaction That Is Essential for the Biogenesis of Histone Locus Bodies. <i>Journal of Biological Chemistry</i> , 2014, 289, 33767-33782.   | 3.4  | 35        |
| 15 | Molecular mechanisms for the regulation of histone mRNA stem-loop binding protein by phosphorylation. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2014, 111, E2937-46.                              | 7.1  | 29        |
| 16 | Structure of Histone mRNA Stem-Loop, Human Stem-Loop Binding Protein, and hExo Ternary Complex. <i>Science</i> , 2013, 339, 318-321.   | 12.6 | 101       |
| 17 | Emergence of the $\hat{I}^2$ -CASP ribonucleases: Highly conserved and ubiquitous metallo-enzymes involved in messenger RNA maturation and degradation. <i>Biochimica Et Biophysica Acta - Gene Regulatory Mechanisms</i> , 2013, 1829, 532-551. | 1.9  | 48        |
| 18 | A Complex Containing the CPSF73 Endonuclease and Other Polyadenylation Factors Associates with U7 snRNP and Is Recruited to Histone Pre-mRNA for 3' End Processing. <i>Molecular and Cellular Biology</i> , 2013, 33, 28-37.                     | 2.3  | 67        |

| #  | ARTICLE   | IF   | CITATIONS |
|----|---|------|-----------|
| 19 | 3' End processing of histone pre-mRNAs in <i>Drosophila</i> : U7 snRNP is associated with FLASH and polyadenylation factors. <i>Rna</i> , 2013, 19, 1726-1744.  | 3.5  | 47        |
| 20 | FLASH Is Required for the Endonucleolytic Cleavage of Histone Pre-mRNAs but Is Dispensable for the 5' Exonucleolytic Degradation of the Downstream Cleavage Product. <i>Molecular and Cellular Biology</i> , 2011, 31, 1492-1502.   | 2.3  | 34        |
| 21 | The hunt for the 3' endonuclease. <i>Wiley Interdisciplinary Reviews RNA</i> , 2010, 1, 325-340.  | 6.4  | 21        |
| 22 | An RNA end tied to the cell cycle: New ties to apoptosis and microRNA formation?. <i>Cell Cycle</i> , 2010, 9, 1308-1312.   | 2.6  | 7         |
| 23 | Studies of the 5' Exonuclease and Endonuclease Activities of CPSF-73 in Histone Pre-mRNA Processing. <i>Molecular and Cellular Biology</i> , 2009, 29, 31-42.   | 2.3  | 69        |
| 24 | FLASH, a Proapoptotic Protein Involved in Activation of Caspase-8, Is Essential for 3' End Processing of Histone Pre-mRNAs. <i>Molecular Cell</i> , 2009, 36, 267-278.  | 9.7  | 113       |
| 25 | Formation of the 3' end of histone mRNA: Getting closer to the end. <i>Gene</i> , 2007, 396, 373-390.   | 2.2  | 157       |
| 26 | Nucleases of the Metallo- $\beta$ -lactamase Family and Their Role in DNA and RNA Metabolism. <i>Critical Reviews in Biochemistry and Molecular Biology</i> , 2007, 42, 67-93.  | 5.2  | 117       |
| 27 | Differences and similarities between <i>Drosophila</i> and mammalian 3' end processing of histone pre-mRNAs. <i>Rna</i> , 2005, 11, 1835-1847.  | 3.5  | 23        |
| 28 | The Polyadenylation Factor CPSF-73 Is Involved in Histone-Pre-mRNA Processing. <i>Cell</i> , 2005, 123, 37-48.  | 28.9 | 180       |
| 29 | A 3' Exonuclease that Specifically Interacts with the 3' End of Histone mRNA. <i>Molecular Cell</i> , 2003, 12, 295-305.  | 9.7  | 106       |
| 30 | 3' End Processing of <i>Drosophila melanogaster</i> Histone Pre-mRNAs: Requirement for Phosphorylated <i>Drosophila</i> Stem-Loop Binding Protein and Coevolution of the Histone Pre-mRNA Processing System. <i>Molecular and Cellular Biology</i> , 2002, 22, 6648-6660. | 2.3  | 48        |
| 31 | A novel zinc finger protein is associated with U7 snRNP and interacts with the stem-loop binding protein in the histone pre-mRNP to stimulate 3'-end processing. <i>Genes and Development</i> , 2002, 16, 58-71.  | 5.9  | 73        |
| 32 | Dual role for the RNA-binding domain of <i>Xenopus laevis</i> SLBP1 in histone pre-mRNA processing. <i>Rna</i> , 2000, 6, 1635-1648.  | 3.5  | 15        |
| 33 | Stem-Loop Binding Protein Facilitates 3' End Formation by Stabilizing U7 snRNP Binding to Histone Pre-mRNA. <i>Molecular and Cellular Biology</i> , 1999, 19, 3561-3570.  | 2.3  | 125       |